

# Cell culture developing and the imaging of total protein product changing with SDS-PAGE in *Saccharomyces cerevisiae*

*Abdullah Aslan*

Department of Molecular Biology and Genetics Faculty of Science Firat University, Turkey- Email: aaslan@firat.edu.tr

**Summary.** Five groups were created in this work. i: Control group, ii: H<sub>2</sub>O<sub>2</sub> group, iii: H<sub>2</sub>O<sub>2</sub>+ %10 Mulberry Juice (MBJ) group, iv: H<sub>2</sub>O<sub>2</sub>+ %15 MBJ, v: H<sub>2</sub>O<sub>2</sub>+ %25 MBJ group. After sterilization, H<sub>2</sub>O<sub>2</sub> and fruit juice were inserted different concentration to *Saccharomyces cerevisiae* (*S. cerevisiae*) cultures and the cultures were developed at 30°C for 1h, 3h, 5h and 24 hours (overnight). *S. cerevisiae* cell growth was computed by spectrophotometer, total protein alteration was analysed by SDS-PAGE electrophoresis and reckoned with Bradford method. Our studies results indicated that; cell developing increased in MBJ groups in proportion to the positive control (H<sub>2</sub>O<sub>2</sub>) group at different growing times (1, 3, 5 and 24 hours) (p<0,05). As a result MBJ has a preservative role for reduce the oxidative damage and expanded cell developing and encourage protein synthesis in *S. cerevisiae*.

**Key words:** *S. cerevisiae*, mulberry juice, hydrogen peroxide, protein expression, SDS-PAGE

## Introduction

*Saccharomyces cerevisiae* (*S. cerevisiae*) is important yeast and it has been employed recent researchs (1). The uptake of H<sub>2</sub>O<sub>2</sub> by *S. cerevisiae* is to change the production of total protein and fatty acid in plasma membrane (1). ROS can oxidize protein, nucleic acid, fat and carbohydrates. For example, the oxidative abuse to proteins bring about to collapse of amino acid shacles diminishing the biologic activity (1, 3-5). Many works executed assert that unlike fruit content expands cellular growth in yeasts, supports protein expression and shows preservative properties towards oxidative stress (6-8). In reference to a study it has been detected that the intake of H<sub>2</sub>O<sub>2</sub> at lower dose, lead to lethal stress in *S. cerevisiae* and bring on negative effect on the expression of significant proteins (1, 3-5). Native antimicrobials can be used with varied new conservation technologies to make easy the modification of conventional attitudes in food

prevention (9). In the last years, new kind of fruit juice products, including pomegranate, strawberry, mulberry, grapefruit, lemon juice, etc. have become very important for human health (10, 11). Fruit and vegetable juices are useful for the people live every time. Low sodium, cholesterol, fat; rich polyphenol, flavonoids and vitamin C acting essential roles in the salutary lives of people (12) in addition for example almond very distinguished for human health with regard to its protein and fatty acid contents (13, 14). Mulberry (MB) is one of the most consumed fruit in the world and it has a nice color, aroma and its leaves have been used as treatment of different illness. In addition, MB is also the source of quercetin, rutin, isoquercetin, and astragaloside and significant phenolic compounds, this compounds has preservative effect against H<sub>2</sub>O<sub>2</sub>-induced oxidative damage, antidiabetic, anti-inflammatory activity and inhibit oxidative injury (15-18). In this work we studied the effect of MB on the rate of the cell developing, total protein expres-

sion and cell proliferation that the induced with  $H_2O_2$  against to oxidative injury growing at  $30^\circ C$  temperature of adding to MB in *S. cerevisiae* culture.

## Material and Methods

### *Research groups and growth conditions*

In this research five groups were composed. i: Control group, ii:  $H_2O_2$  group, iii:  $H_2O_2$  + %10 Mulberry Juice (MBJ) group, iv:  $H_2O_2$  + %15 MBJ, v:  $H_2O_2$  + %25 MBJ group. After sterilization,  $H_2O_2$  and fruit juice were inserted different concentration to *Saccharomyces cerevisiae* (*S. cerevisiae*) cultures and the cultures were developed at  $30^\circ C$  for 1h, 3h, 5h and 24 hours (overnight). *S. cerevisiae* cell proliferation was calculated by spectrophotometer, total protein expression was indicated by SDS-PAGE electrophoresis and reckoned with bradford method for the developed and reproduce of yeast, YEPD (for 50 mL 1,5 g yeast extract, 1 g trypton, 1,5 g glucose) in addition, for the developing and reproduce of *S. cerevisiae*, mulberry fruit juices was added and cultivated. After sterilization, yeasts were cultured into media and the samples were incubated for 1h, 3h, 5h, 24 h (overnight, h: hour) at  $30^\circ C$  (7).

### *Mulberry juice extract and $H_2O_2$ Chemical*

Fruit (From center county of Elazığ city) was squashed in water and added in to *S. cerevisiae* media cultures and added 20% (v/v) ratio in at the reproducing for  $30^\circ C$ .  $H_2O_2$  was inserted in  $H_2O_2$  and MBJ +  $H_2O_2$  groups.

### *Cell concentration measurements*

In these measurements, culture samples that were examined at  $30^\circ C$  for 1, 3, 5 hours and overnight (24 hours) have been analyzed. The calculation has been accomplished using a spectrophotometer at 600 nm ( $OD_{600}$ ).

### *SDS-PAGE (Sodium dodecyl sulfate polyacrylamide gel electrophoresis)*

SDS-PAGE was made using BIO-RAD Mini-PROTEAN® 3 Cell gel electrophoresis system. The samples of *S. cerevisiae* cultures were organized for SDS-PAGE after which they were loaded to sample

loading wells to be subject to electrical current and after this process, their images were taken and the intergroup protein bandings were used as data in the study (19).

### *Protein density measurements*

The measurement has been accomplished using a spectrophotometer at 600 nm ( $OD_{600}$ ) with regard to bradford method. BSA (bovine serum albumin) protein standards at different concentrations were obtained using BSA protein. Accordingly, the total protein amount in *S. cerevisiae* groups corresponding to this standard valuation was calculated (Figures 3, 4).

### *Statistical Analysis*

SPSS 20.0 software was used. The comparison between experimental groups and the control group was made using one way ANOVA and Post Hoc Duncan and Games howell tests. Statistically important differentiation among groups have been stated as  $p < 0.05$  and the statistically non-significant differences have been specified as  $p > 0.05$ . Standard deviations were point out as  $\pm$ .

## Results and Discussion

We think that the results of this study will provide important contributions to the present literature. The results of table 1 and figure 1 show that mulberry has essential effects on *S. cerevisiae* proliferation. It is indicated that mulberry juice (MBJ) maintains its live cell amount in spite of the growing hydrogen peroxide densities. A dissimilarity is detected between the yeast proliferation amounts for 1h in comparison with the control ( $p < 0.05$ ). It is observed that MBJ preserves the cell almost as much as the control opposite hydrogen peroxide which is the great radical origin in the 25% MBJ +  $H_2O_2$  group and 15% MBJ +  $H_2O_2$  group. When 3h values are investigated; it is obtained that MBJ has increased yeast development in the 25% MBJ +  $H_2O_2$  group, in spite of the inverse effects of the hydrogen peroxide radical comparatively the control and  $H_2O_2$  group ( $p < 0.05$ ). When the 5h values are investigated; it is obtained that MBJ has risen yeast improving at a maximum level in the 10% MBJ +  $H_2O_2$  group despite the inverse effects of the hydrogen peroxide radical


**Table 1.** *Saccharomyces cerevisiae* cell growth in mulberry juices

OD <sub>600</sub> 30°C	1h	3h	5h	Overnight
Control	1,488±0,00 <sup>b</sup>	1,395±0,00 <sup>c</sup>	1,755±0,00 <sup>c</sup>	2,096±0,00 <sup>c</sup>
H <sub>2</sub> O <sub>2</sub>	1,510±0,00 <sup>c</sup>	1,413±0,00 <sup>d</sup>	1,776±0,00 <sup>d</sup>	1,876±0,00 <sup>a</sup>
H <sub>2</sub> O <sub>2</sub> + 10% mulberry	1,073±0,00 <sup>a</sup>	1,286±0,00 <sup>a</sup>	1,774±0,00 <sup>d</sup>	2,092±0,00 <sup>c</sup>
H <sub>2</sub> O <sub>2</sub> + 15% mulberry	1,577±0,00 <sup>d</sup>	1,364±0,00 <sup>b</sup>	1,556±0,00 <sup>a</sup>	2,102±0,00 <sup>d</sup>
H <sub>2</sub> O <sub>2</sub> + 25% mulberry	1,605±0,00 <sup>c</sup>	1,464±0,00 <sup>c</sup>	1,741±0,00 <sup>b</sup>	2,071±0,00 <sup>b</sup>


\*\*a,b,c,d,e; among the groups which bearing of different letter are significant ( $p < 0.05$ ). one way ANOVA and Post Hoc Duncan and Games howell tests

comparatively the control ( $p < 0.05$ ). When the overnight (24 h) values are investigated; it is obtained that MBJ has risen yeast growth in the 10% MBJ + H<sub>2</sub>O<sub>2</sub>, 15% MBJ + H<sub>2</sub>O<sub>2</sub> and 25% MBJ + H<sub>2</sub>O<sub>2</sub> groups, in spite of the opposite effects of the hydrogen peroxide radical in comparison with the control and H<sub>2</sub>O<sub>2</sub> group; besides it can also be obtained that yeast growth has risen at a statistically significant degree in all other groups comparatively the control and H<sub>2</sub>O<sub>2</sub> groups ( $p < 0.05$ ) (Table 1). Stinco et al (2015) indicated that orange juice activates the antioxidant defensive system towards free radicals for yeast development (20). Aslan et al (2014a) have indicated that pomegranate juice is protective against oxidative injury in *S. cerevisiae* (1). Again Aslan (2015) indicated that as a result of the work performed with several fruit juices and their mixtures that different fruit juices and their mixtures are preservative against oxidative injury in *S. cerevisiae* and that they rise yeast development (10). Tserennadmid et al (2011) have stated that apple juice has a protective role for growing in yeasts (21). Krivoruchko and

Nielsen (2015) have stated that resveratrol and flavonoids act protective roles towards oxidative injury in bacteria and yeasts (22). Zhang et al (2017) showed that the mulberry extract has protective effect in human cell culture against oxidative stress (17). Rynko et al (2016) indicated that the leave of mulberry has antidiabetic activity, anticancer activity, antibacterial activity (15). Riche et al (2017) demonstrated that mulberry leaf extract decrease the human blood glucose level (18). Gregorio et al (2011) have stated that mulberry extract has antioxidant activity in *S. cerevisiae* (23). Chen et al (2015) have put forth that mulberry fruit has antioxidant and hyperglycemic activity in vitro (16). When the SDS-PAGE results are investigated; it is obtained that protein band intensity rise in pellet gel images is greater in groups to which MBJ is applied



**Figure 1.** The growing of *Saccharomyces cerevisiae* in mulberry juices at different hours.



**Figure 2.** SDS-PAGE pelet total protein bands profiles for development at 30°C. Lanes 1: Marker; 2: Control; 3: H<sub>2</sub>O<sub>2</sub>; 4: H<sub>2</sub>O<sub>2</sub> +10% MBJ; 5: H<sub>2</sub>O<sub>2</sub> +15% MBJ; 6: H<sub>2</sub>O<sub>2</sub> +25% MBJ

in comparison with the control (Figure 2). Aslan et al (2014b) have stated that pomegranate juice has a preservative effect in *S. cerevisiae* towards oxidative injury reasened by the applying of hydrogen peroxide and that protein band intensity rise is bigger in pomegranate applied groups comparatively hydrogen peroxide applied groups (3). When the bradford results in Figure 3 and figure 4 are analyzed; large protein quantity has been calculated in MBJ ( $H_2O_2$ +10% MBJ,  $H_2O_2$ +15% MBJ,  $H_2O_2$ +25% MBJ) applied groups comparatively to control and  $H_2O_2$  groups ( Figure 3,4). However, there are a lotof research in vivo on rat about fruit and vegetable mechanism. For example these, Aslan et al (2014c) and Aslan et al (2016a) have indicated that the milk thistle extract is preservative towards lung damage in rats (24, 25), Aslan and Can (2017a) have stated that lemon juice has a protective effect for diminish the oxidative injury, increased cell growing and protein synthesis in *S. cerevisiae* culture (26), Aslan et al (2016b) demonsrated that black cumin extract may be a drug for lung damage in rats (27), Aslan et al (2015) indicated that *Nigella sativa* extracts has a preservative effects against to rats

lung damage (28), Ozsahin et al (2009) expressed that different sugar extracts induce fatty acid biosynthesis in the *S. cerevisiae* cell culture (29), Aslan et al (2017b) indicated that kiwi fruit juice has a protective effect against to hydrogen peroxide damage in *S. cerevisiae* (30).. With respect to these results, MBJ has a positive effect on *S. cerevisiae* cell proliferation and decreased the oxidative injury effect.

## Conclusion

When these results are evaluated; we can said that MBJ is quite effective towards the hydrogen peroxide induced oxidative injury in *S. cerevisiae*, that it safeguards cell improve and even rises cell thrive; thus supporting protein expression in yeast cells. With respect to these findings, we expect that our study will support other studies to try MBJ in animal experiments and that in this respect MBJ will be digested more by people based on the positive results that will be gained.




Figure 3. Protein density at between groups




Figure 4. Bradford BSA (bovine serum albumin) standart graph

## References

- Aslan A, Can MI, Boydak D. Anti-oxidant effects of pomegranate juice on *Saccharomyces cerevisiae* cell growth. Afr J Tradit Complement Altern Med. 2014; 11(4):14-18
- Folmer V, Pedroso N, Matias AC, Lopes SCDN and F Antunes. H<sub>2</sub>O<sub>2</sub> induces rapid biophysical and permeability changes in the plasma membrane of *Saccharomyces cerevisiae*. Biochim Biophys Acta. 2008; 1778: 1141-1147.
- Aslan A, Baspinar S, Yilmaz O. Is Pomegranate juice has a vital role for protective effect on *Saccharomyces cerevisiae* growth? Progress in Nutrition 2014; 16(3): 212-217.
- Zupan J, Mavri J and P. Raspor. Quantitative cell wall protein profiling of invasive and non-invasive *Saccharomyces cerevisiae* strains. J Microbiol Meth. 2009; 79: 260-265.
- Chondrogianni N, Petropoulos I, Grimm S, Georgila K, Catalgol B, Friguet B, Grune T, and Gonos, ES. Protein damage, repair and proteolysis. Mol Aspects Med. 2014; 35: 1-71.
- Tyagi AK, Gottardi D, Malik A, Guerzoni ME. Chemical composition, in vitro anti-yeast activity and fruit juice preservation potential of lemon grass oil. LWT Food Sci Technol 2014; 57: 731-737. 13.
- Aslan A, Can MI. The inhibition of chromium effect in *Saccharomyces cerevisiae* thrive from grapefruit. Progress in Nutrition 2015b; 17(4): 339-342.
- Aslan A and Can MI. The effect of orange juice against to H<sub>2</sub>O<sub>2</sub> stress in *Saccharomyces cerevisiae*. Progress in Nutri-

- on 2015c; 17(3): 250-254.
9. Bukvicki D, Gottardi D, Tyagi AK, Veljic M, Marin Petar D, Vujisic L, Guerzoni ME, Vannini L. Scapania nemorea liverwort extracts: Investigation on volatile compounds, in vitro antimicrobial activity and control of *Saccharomyces cerevisiae* in fruit juice, *LWT - Food Science and Technology* 2014; 55: 452-458.
  10. Aslan A. The effects of different essential fruit juice and their combination on *Saccharomyces cerevisiae* cell growth. *Progress in Nutrition* 2015;17: 36-40.
  11. Aguirre DB, Barbosa-Canovas GV. Inactivation of *Saccharomyces cerevisiae* in pineapple, grape and cranberry juices under pulsed and continuous thermo-sonication treatments, *Journal of Food Engineering* 2012; 108: 383-392.
  12. Patrignani F, Vannini L, Kamdem SLS, Lanciotti R, Guerzoni ME. Effect of high pressure homogenization on *Saccharomyces cerevisiae* inactivation and physico-chemical features in apricot and carrot juices, *International Journal of Food Microbiology* 2009; 136: 26-31.
  13. Karatay H, Sahin A, Yilmaz O, Aslan A. Major Fatty Acids Composition of 32 Almond (*Prunus dulcis* (Mill.) D. A. Webb.) Genotypes Distributed in East and Southeast of Anatolia, *Turkish Journal of Biochemistry* 2014a; 39(3); 307-316.
  14. Karatay H, Sahin A, Aslan A. Determining seed storage proteins of some almond (*Prunus dulcis* (Mill.) D.A. Webb.) genotypes distributed in east and southeast of Anatolia, *Progress in Nutrition* 2014b; 16: 116-125.
  15. Rynko AG, Bazylak G, Slonina DO. New potential phytotherapeutics obtained from white mulberry (*Morus alba* L.) leaves, *Biomedicine & Pharmacotherapy* 2016; 84: 628-636.
  16. Chen C, You LJ, Abbasi AM, Fua X, Liua RH. Optimization for ultrasound extraction of polysaccharides from mulberry fruits with antioxidant and hyperglycemic activity *in vitro*, *Carbohydrate Polymers* 2015; 130: 122-132.
  17. Zhang L, Xu Y, Li Y, Bao T, Gowd V, Chen W. Protective property of mulberry digest against oxidative stress – A potential approach to ameliorate dietary acrylamide-induced cytotoxicity, *Food Chemistry* 2017; 230: 306-315.
  18. Richea DM, Riche KD, Eastc HE, Barretta EK, May WL. Impact of mulberry leaf extract on type 2 diabetes (Mul-DM): A randomized, placebo-controlled pilot study, *Complementary Therapies in Medicine* 2017; 32: 105-108.
  19. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227: 680-685.
  20. Stinco CM, Baroni MV, Naranjo RDDP, Wunderlin DA, Heredia FJ, Melendez-Martinez AJ, Vicario IM. Hydrophilic antioxidant compounds in orange juice from different fruit cultivars: Composition and antioxidant activity evaluated by chemical and cellular based (*Saccharomyces cerevisiae*) assays, *Journal of Food Composition and Analysis* 2015; 37: 1-10.
  21. Tserennadmid R, Tako M, Galgoczy L, Papp T, Pesti M, Vagvolgyi C, Almassy K, Krisch J. Anti yeast activities of some essential oils in growth medium, fruit juices and milk, *International Journal of Food Microbiology* 2011; 144 480-486.
  22. Krivoruchko A and Nielsen J. Production of natural products through metabolic engineering of *Saccharomyces cerevisiae*, *Current Opinion in Biotechnology* 2015; 35:7-15.
  23. Gregorio MRP, Regueiro J, Gonzalez EA, Castro LMP, Gandara JS. Influence of alcoholic fermentation process on antioxidant activity and phenolic levels from mulberries (*Morus nigra* L.), *LWT - Food Science and Technology* 2011; 44: 1793-1801.
  24. Aslan A, Can MI. Milk thistle impedes the development of carbontetrachloride-induced liver damage in rats through suppression of bcl-2 and regulating caspase pathway. *Life Sciences* 2014c; 117: 13-18.
  25. Aslan A, Can MI, Kuloglu, T. Baspinar, S. Milk thistle may induce apoptosis in development of carbontetrachloride-induced liver DNA damage in rats. *Progress in Nutrition* 2016; 18, 146-151.
  26. Aslan A, Can MI. Protein expression product alterations in *Saccharomyces cerevisiae*. *Progress in Nutrition* 2017a; 19, 81-85.
  27. Aslan A. Boydak D. Can MI. Kuloglu, T. Baspinar, S. Black cumin may be a potential drug for development of carbon-tetrachloride-induced lung damage in rats. *Progress in Nutrition* 2016; 18, 56-62.
  28. Aslan A, Boydak D, Can MI, Kuloglu T. Nigella sativa improves the carbon tetrachloride-induced lung damage in rats through repression of erk/akt pathway, *Bangladesh Journal of Pharmacology* 2015; 10: 654-659.
  29. Ozsahin AD, Guvenc M, Yilmaz O, Aslan A, Tuzcu M The effects of different sugar sources on fatty acid biosynthesis in the *Saccharomyces cerevisiae* cell culture. *J Anim Vet Adv.* 2009; 8: 424-429.
  30. Aslan A, Gok O, Erman O. The protective effect of kiwi fruit extract against to chromium effect on protein expression in *Saccharomyces cerevisiae*. *Progressin Nutrition* 2017b; 19: 472-478.

---

Correspondence:

Abdullah Aslan

Department of Molecular Biology and Genetics

Faculty of Science Firat University, Turkey

Tel. +90 (424) 2370000-3819

Fax +90 (424) 2330062

E-mail: aaslan@firat.edu.tr