

Intraoperative squash Cytology of diffuse glioma not otherwise specified, of the Cerebellum

Alexandra Kalogeraki¹, Dimitrios Tamiolakis¹, Iris Zoi¹, John Segredakis¹, Antonios Vakis²

Departments of Pathology-Cytopathology ¹ and Neurosurgery ², Medical School, University of Crete, University Hospital, Heraklion, Crete, Greece.

Abstract. *Objective:* Diffuse glioma arises anywhere in the CNS, but most frequent in the cerebral hemispheres. The tumor tends to be seen in children and in younger adults aged 20-30. We report one such case in an older female patient presenting the intraoperative cytology of the tumor. *Case report:* A 48-year-old female was diagnosed by MRI with a tumor of cerebellum. Cytologic material was obtained during the resection of the tumor and diagnosed cytologically as glioma. *Conclusion:* This case is presented to focus the ability of the intraoperative cytology in diagnosis of the glioma, using immunocytology and confirmed by histo- immunohistology. (www.actabiomedica.it)

Key words: Glioma cytology, histology.

Introduction

The updated 2016 edition of the World Health Organization (WHO) Classification of Tumours of the Central Nervous System (CNS) uses molecular parameters and the histology to define the main tumor categories. Major reclassification was made with regard to diffuse gliomas, medulloblastomas and other embryonal tumors. In this aspect after the histological confirmation of astrocytoma, the second indicator for adult patients is the presence or absence of isocitrate dehydrogenase (IDH1 or IDH2) mutations and 1p/19q status. A tumor (in adults) with oligodendroglial morphology, showing an IDH mutation but no 1p/19q loss, will be designated astrocytoma, IDH mutated. For diffuse astrocytoma without IDH mutations, the term “IDH wild type” is used. If molecular testing for IDH status could not be completed or was inconclusive, the term “not otherwise specified” (NOS) is used. Recent studies have focused on other genes expression as well (1-4).

Diffuse gliomas tend to be seen in younger adults aged 20-30 and in children (5). They arise anywhere

in the CNS, but most frequent in the cerebral hemispheres. Some gliomas involve a large part of the brain or the entire CNS in a diffuse fashion (gliomatosis cerebri). Most pontine and medullary gliomas are diffuse. Histologically, the tumor cells can be stellate, spindle-shaped with fiber like processes, or plump with a large eosinophilic mass (6).

The cytology of diffuse glioma is characterized by slight hypercellularity, mild nuclear enlargement, and absence of mitotic activity and necrosis (7).

Here we report an intraoperative cytologic diagnosis of a diffuse glioma confirmed by histology.

Case Report

A 48-year-old female patient hospitalized at University Hospital of Heraklion Crete suffering from cranial pain and unsteadiness. She was subsequently diagnosed by MRI with a tumor of cerebellum. Past, personal and family history was free.

The hematological and biochemical parameters were within normal limits.

Squash cytology

Intraoperative squash smears were prepared as follows: 1-2 mm³ of fresh tissue from a specific area after gross evaluation was crushed between two slides to prepare smears as at first described by Adams et al (8). Intraoperative cytological consultation of lesions by squash smear method is an axillary technique to help the neurosurgeon in the management of CNS tumors. It is efficient when applied upon minimal tissue pieces, provides the cytopathologist with cellular pattern (cohesive or non cohesive), cell (nuclear and cytoplasmic) morphology, and allows for immunocytochemical testing. More over it lacks the disadvantage of ice artifacts by frozen section analysis (9). Squash cytology smear method has improved diagnostic efficacy and is an adjunct, reliable simple and cheap tool for the neurosurgeon in the operating theatre (10).

Results

Cytology

The air dried smears were used for Giemsa stain and immunocytochemistry, while the alcohol (60%) fixed for routine Papanicolaou stain (Fig. 1). Isolated elongated, epithelial-like neoplastic cells with oval nuclei and scanty cytoplasm were demonstrated. Mitoses and necrosis were not found. Glioma, ependymoma, medulloblastoma and teratoid/rhaboid tumor, were considered in the differential diagnosis.




Figure 1. Grade-II Astrocytoma of the Cerebellum. Cytological squash smear. Papanicolaou stain x 400.

Immunocytochemistry

Rapid immunocytochemistry for GFAP and cytokeratin on air dried smears took 22 minutes, and showed that the tumor cells were of glial origin: GFAP cytoplasmic positive (Fig. 2)

Additional antibodies for conventional ICC, included, S-100, synaptophysin, smooth muscle actin (SMA), desmin, and EMA. IDH-1 immunostain was not performed

Gross total resection of the tumor was performed by the neurosurgeon.

Histology: In histological specimens of the tumor, fixed in 10% formalin, the H&E stain was performed (Fig. 3).




Figure 2. Grade-II Astrocytoma of the Cerebellum. Cytological squash smear. GFAP immunostain x 400.




Figure 3. Grade-II Astrocytoma of the Cerebellum. Histopathology section. Hematoxyline-Eosin stain x 400.

Abundant material with cells of medium size with elongated or oval nuclei, extremely rare mitoses and neo-angiogenesis was observed..

Immunohistochemistry: The tumor cells expressed cytoplasmic GFAP (Fig. 4), S-100 and Vimentin but were negative for Synaptophysin, NF, EMA, CD34 and p53 markers. The proliferation index MIB-1 was found to be positive in 5% of tumor cells.

IDH(isocitrate dehydrogenase) and 1p/19q status were determined. Our case was a diffuse glioma not otherwise specified as IDH status was inconclusive. The patient received RT (54 Gy/30 fractions (fr)) followed by 6 courses of PCV (procarbazine, CCNU - lomustine, vincristine) chemotherapy. Five months after surgery she is disease free.

Discussion

Gliomas constitute a large and heterogenous group of tumors and notorious for wide differences in clinical incidence gross, microscopic features and biologic behavior.

Morphology cannot be accurate prognostic indicator where pathologists receive small biopsy samples but the gold standard remains histopathological examination of an abundantly sampled tissue (11).

The history of intraoperative cytology dates back to 1930, when Eisenhardt and Cushing introduced this technique by using supravital staining (12). The

technique has recently gained importance because of advent of CT and MRI guided stereotactic biopsies. Cytology has been shown to be a great value in intraoperative consultation of CNS pathology (13) but the diagnosis of low-grade astrocytoma, especially grade-I fibrillary type, may be difficult to diagnose cytologically as well as histologically (14). The differentiation of normal, reactive and neoplastic astrocytes depends of identification of cytologic abnormalities. The astrocytes of reactive gliosis show cell processes and mild degree of cytologic abnormality that overlaps with slight degree of abnormality seen in low-grade astrocytoma (15). Low-grade astrocytomas show mild hypercellularity and slight cytologic atypia and pleomorphism. The biopsy from the central part of the tumor may be more representative but a biopsy taken from a peripheral area can be misdiagnosed cytologically due to sampling error (16).

Intraoperative cytology and frozen section are important in the diagnosis of neurosurgical samples. There are limitations in both procedures but understanding the errors and pitfalls may increase diagnostic accuracy. In a study by Chand P et.al. several technical errors were observed in the frozen sections, the principal being freezing process that changes the architecture, with sampling faults and misinterpretation coming next. In the crush smears, there was disagreement in diagnosis due to sampling and interpretative failure (17).

Intraoperative diagnosis of CNS lesions helps neurosurgical approach of these disorders. The perfect intraoperative process needs to be accurate, rapid, and must permit the maintenance of tissue for paraffin section analysis. Frozen sections provide structural details. Yet, the characteristic soft nature and high water content of the nervous tissue gives poor quality frozen sections. Consequently, different techniques such as squash smears and touch imprints are applied. Squash smears and touch imprints are inexpensive and no special skilfulness is required to be obtained. No particular supplies are needed and minute tissue pieces can be employed, thus enough tissue is available for paraffin section examination. Failure to manage thickness, crushing artefacts, and inappropriate smearing are the limitations of squash smears. Freezing artefacts can be prevented by most desirable temperature control and




Figure 4. Grade-II Astrocytoma of the Cerebellum. Histopathology section. GFAP immunostain x 400.

prompt freezing of tissue. Touch imprints show better structural details than squash smear as crush artefacts are restrained (18-24).

Rapid immunocytochemistry is limited to some antibodies well documented in the literature such as keratins (25,26)

IDH-1 immunostain has values and limitations. It has been reported that it does not reliably identifies infiltrating tumor cells when admixed with pre-existing or reactive glial cells and that it produces a non specific background staining (27).

In our case the use of intraoperative cytology not only helped the surgeon in rapid diagnosis of glioma, also it was ensured that minimum injury is caused to the normal brain structures surrounding the tumor.

In conclusion this report was undertaken to access utility of intraoperative smear cytology in diagnosing and grading glioma correlating and confirming by histopathological and immunohistological stains.

Conflicts of interest: Each author declares that he or she has no commercial associations (e.g. consultancies, stock ownership, equity interest, patent/licensing arrangement etc.) that might pose a conflict of interest in connection with the submitted article.

References

- Li Z, Guan Y, Liu, Q. et al. Astrocytoma progression scoring system based on the WHO 2016 criteria. *Sci Rep* 9, 96 (2019)
- Louis DN, Ohgaki H, Wiestler OD, Cavenee WK: WHO classification of tumours of the central nervous system, ed 4 Lyon, IARC Press, 2016
- Komori T. The 2016 WHO Classification of Tumours of the Central Nervous System: The Major Points of Revision *Neurol Med Chir (Tokyo)* 2017 Jul; 57(7): 301-311
- Van den Bent MJ, Weller M, Wen PY, Kros JM, Aldape K, Chang S. A clinical perspective on the 2016 WHO brain tumor classification and routine molecular diagnostics. *Neuro-Oncology* 2017, 19 (5): 614-624
- Chandana SR, Movva S, Arora M, Singh T. Primary brain tumors in adults. *Am Fam Physician.* 2008 May 15;77(10):1423-30.
- Rodriguez FJ, Lim KS, Bowers D, and Eberhart CG. Pathological and Molecular Advances in Pediatric Low Grade Astrocytoma. *Annu Rev Pathol.* 2013 Jan 24; 8: 361-379.
- Parakarama T, Chandrasoma PT, Apuzzo ML. Stereotactic brain biopsy: Astrocytic neoplasms. Los Angeles California, 1989: 96-97.
- Adams JH, Graham DI, Doyle D. London: Chapman and Hall; 1981. Brain Biopsy: the smear technique for neurosurgical biopsies; pp. 11-4
- Kishore S, Thakur B., Bhardwaj A, Kusum A Diagnostic accuracy of squash cytology and role of GFAP immunopression in glial tumors. *Indian Journal of Pathology and Oncology*, January-March, 2018;5(1):12-17
- Sarkar S, Sengupta M, Datta Ch, Chatterjee U, Ghosh SN. Evaluation of intraoperative cytological smears for diagnosis of brain tumors with special reference to immunohistochemistry. *Indian J Med Paediatr Oncol.* 2017 Jul-Sep; 38(3): 296-301.
- Sukheeraj D, Singhvi S, Rai NN, and Midya M. A Comparative Study of Histopathology of Astrocytomas with Intraoperative Cytology with Special Reference to MIB-1 Labelling Index. *J Clin Diagn Res.* 2015 Aug; 9(8): EC01-EC03.
- Eisenhardt L and Cushing H. Diagnosis of Intracranial Tumors by Supravital Technique. *Am J Pathol.* 1930 Sep; 6(5): 541-552.7.
- Savargaonkar P and Farmer PM. Utility of Intra-operative Consultations for the Diagnosis of Central Nervous System Lesions. *Ann Clin Lab Sci Spring* 2001 vol. 31 no. 2 133-139.
- Ranchod M. Intraoperative consultation in Surgical Pathology. In Weidner, Cote, Suster, Weiss. *Modern Surgical Pathology* Second Edition. Saunders, Elsevier 2009; Chapter 2: pp 13-26
- Robbins PD, Yu LL, Lee M, Stokes BA, Thomas GW, Watson P, Wong G. Stereotactic biopsy of 100 intracerebral lesions at Sir Charles Gairdner Hospital. *Pathology.* 1994 Oct;26(4):410-3.
- Nasir H, and Haque AUI. Value of Touch Preparation Cytology in Intraoperative Consultation Diagnosis of Astrocytomas. *International Journal of Pathology* 2003; 1: 8-12.
- Chand P, Amit S, Gupta R, Agarwal A. Errors, limitations, and pitfalls in the diagnosis of central and peripheral nervous system lesions in intraoperative cytology and frozen sections. *J Cytol.* 2016;33(2):93-97
- Narang V, Jacob S, Mahapatra D, Mathew JE. Intraoperative diagnosis of central nervous system lesions: Comparison of squash smear, touch imprint, and frozen section. *J Cytol.* 2015;32:153-8
- Jaiswal S, Vij M, Jaiswal AK, Behari S. et.al. Intraoperative squash cytology of central nervous system lesions: a single center study of 326 cases. *Diagnostic Cytopathology* 2012; 40(2):104 - 112
- Roessler K, Dietrich W, Kitz K. High Diagnostic Accuracy of Cytologic Smears of Central Nervous System Tumors. A 15-Year Experience Based on 4,172 Patients.. *Acta cytologica* 2002;46(4):667-74
- Cheunsuchon P, Punyashthira A, Tanboon J, Sangruchi T. Accuracy of Intraoperative Consultation of Central-Nervous System Lesions in Siriraj Hospital. *Siriraj Med J* 2014;66:113-119.
- Jindal A, Diwan H, Kaur K, and Sinha VD. Intraoperative Squash Smear in Central Nervous System Tumors and Its

- Correlation with Histopathology: 1 Year Study at a Tertiary Care Centre. *J Neurosci Rural Pract.* 2017 Apr-Jun; 8(2): 221-224.
23. Govindaraman PK, Arumugam N, Ramasamy C, Prakasam G. Role of squash smear in intraoperative consultation of central nervous system tumors. *J Sci Soc* 2017;44:7-14.
24. Khuroo MS, Hamdani SM, Alam SS, Sahaf BR, Dar NQ, Bhat RA. Accuracy and utility of intraoperative squash smear cytology in neurosurgical practice. *Int J Med Sci Public Health* 2019;8(2):130-135.
25. Zhang X, Liu J, Yan X. Rapid intraoperative immunocytochemistry of central nervous system tumors. *Int. J. Clin. Exp. Pathol* 2020;13(1):44-48).
26. Moriya J, Tanino MA, Takenami T, Ehdoh T et al; R-IHC Study Group. Rapid immunocytochemistry based an alternating current electric field using squash smear preparation of central nervous system tumors. *Brain Tumor Pathol* 2016;33:13-18
27. Preusser M, Wohrer A, Stary S, Hoftberger R et al Value and Limitations of immunohistochemistry and gene sequencing for detection of the IDH1-R132H mutation in diffuse glioma biopsy specimens. *J. Neuropathol Exp. Neurol.* 2011;70(8): 715-721.
-
- Received: 3 August 2020
Accepted: 17 August 2020
Correspondence:
Alexandra Kalogeraki, MD, PhD. University of Crete, Medical Faculty, PO Box 1393, Heraklion 71110, Crete Greece
Tel:+ 30 2810 394692
Fax:+30 2810 394694
E-mail: kalogerakimed@yahoo.gr