

C A S E R E P O R T

High energy trauma with complex fracture of the right tibial plateau, compartment syndrome and infection in a professional freestyler motorcyclist

Alessio Pedrazzini¹, Alice Garzia², Nicola Bertoni¹, Henry Yerwo Simo¹, Roberto Bisaschi¹, Vanni Medina¹, Francesco Pogliacom²

¹Orthopaedic Unit, Oglio Po Hospital, Vicomoscato (CR), Italy ; ²Orthopaedic and Traumatology Clinic. Department of Medicine and Surgery, University of Parma, Parma, Italy

Summary. High-energy tibial plateau fractures are complex lesions that may be associated with extensive soft tissue damages and severe complications, such as compartment syndrome and neurological injuries. This case report interests a professional motocross freestyler with complex tibial plateau fracture associated with compartment syndrome and partial common peroneal nerve impairment which complicated with a dehiscence of the surgical wound and infection after plate and screws fixation. One year later, despite the complexity of the fracture, the gravity of the soft tissue lesions and subsequent complications, the patient healed. This satisfactory result depended on the correct management in terms of type of treatment and timing. (www.actabiomedica.it)

Key Words: Knee; Tibial plateau; Fracture; Soft tissues; Fixation

Introduction

High-energy tibial plateau fractures are complex lesions that are often associated with extensive soft tissue damages and severe complications such as compartment syndrome and neurological injuries. The consequences of the severity of the injury and of an inadequate treatment can be serious and accompanied by unsatisfactory results. In this context it is important the correct timing of surgery, that inevitably adds further traumatism to soft tissues, thus augmenting the possibility of wound complication, especially if open reduction and internal fixation (ORIF) is performed (1).

For this reason, in the preoperative planning it is also important the distinction of tibial plateau fractures into complex and simple that allows to identify the more serious or potentially serious injuries, whose treatment is more difficult and more frequently characterized by complications. Zeltser in accordance with

Schatzker (2) considers type IV, V and VI fractures as complex. Type V (fractures of the medial and lateral hemiplates without metaphyseal involvement) and VI (fractures of the two hemiplates with associated metaphyseal involvement) are consequent in most cases, above all in young patients, of high-energy traumas and for definition are multi-fragmented. Also type IV fracture (medial hemiplate fracture), although not multi-fragmented, generally derives from high-energy traumas and their typical mechanism (stress in varus) can cause strain injuries of the common peroneal nerve.

Optimal treatment of the tibial plateau fractures is still open to debate and a wide choice of methods can be used (ORIF, minimally invasive plate osteosynthesis, arthroscopically assisted fixation, external fixation); open reduction with internal fixation seems to be the best therapeutic option in complex fractures. In any case the goal of treatment is to obtain limb axis restoration, joint stability and accurate intra-articular

reduction, avoiding further soft tissue damages (3,4); obtaining these aims is not simple and it requires excellent soft tissue handling and fracture reduction skills to avoid iatrogenic complications (5-10).

This case report refers of a professional motocross freestyler with complex tibial plateau fracture with compartment syndrome and partial impairment of the common peroneal nerve which complicated with a dehiscence of the surgical wound and infection of plate and screws.

Case report

A 32-year-old male, motocross freestyler, arrived at the emergency department of another hospital after an high-energy motorcycle accident with the diagnosis of complex fracture of the right tibial plateau (Figure 1) associated with acute compartment syndrome and partial impairment of the common peroneal nerve. The patient's only comorbidity was ulcerative rectocolitis. Colleagues performed in emergency a posterior and lateral fasciotomy, positioned the limb in transkeletal traction to the heel and applied a plaster cast. When 2 days later the patient arrived at our hospital (Figure 2), he had an ipomotility of the active dorsal extension of the foot and toes and a partial loss of sensibility of the dorsum of the foot.

A Computed Tomography CT scan with 3D reconstructions (Figure 3) was performed in order to study the characteristics of the fracture and to correctly plan osteosynthesis; according to Schatzker this fracture was classified as type VI (2). A bridging external fixator from the femur to the tibia was applied.

The patient underwent definitive fixation two weeks after trauma, thus allowing the soft tissues to heal. An isolated medial surgical approach was performed. Lateral and posterior surgical access were not executable due to the presence of the fasciotomy which were simultaneously closed. Multi-fragmented displaced fracture of the proximal tibial epiphysis associated with bone loss (Figure 4) was synthesized with ORIF with plate and screws. A cadaveric femoral head graft was positioned in order to fill the bone loss and to increase the stability of the fixation (Figure 5).

After surgery, the patient presented intermit-




Figure 1. Pre-operative radiographs.



Figure 2. Clinical views 2 days after fasciotomy.




Figure 3. CT scans with 3D reconstruction.

tent fever and raised white blood cells; uro- and blood culture and pharyngeal swab were performed (results negative for ongoing infections). The surgical wound initially appeared normal but its condition progressively worsened (Figure 6), with dehiscence and serum-purulent secretion. Ten days after surgery, culture swabs were performed from the exudate of the surgical wound. The culture test report demonstrated the in-



Figure 4. Intra-operative view which shows the bone loss.




Figure 5. Post-operative X-rays following ORIF with plate and screws and bone graft.

fection with *Staphylococcus Aureus* Methicillin Resistant. At the same time magnetic resonance imaging excluded the presence of abscess collections.

On the basis of the antibiogram, specific antibiotic therapy was started as prescribed by the infectious disease specialist. In the following days two surgical wound debridements were performed as well as hyperbaric therapy. These treatments were associated with plate and screws removal which was done 35 days after definitive fixation (Figure 7).




Figure 6. Dehiscence of the surgical wound.




Figure 7. X-rays after removal of plate and screws.

The patient was discharged from the Hospital 2 weeks following the last surgery with a brace blocked at 30 degrees of flexion for another 2 weeks, a Codivilla splint and specific oral antibiotic therapy. During the following weeks the brace was progressively unlocked and the patient continued with assisted physiotherapy. Partial weight bearing was allowed 2.5 months after ORIF. Nowadays, the patient has obtained full load and he has started again his sporting activity. The inflammation indexes have progressively improved until normalization, as well elettromyographic findings.

X-ray controls performed 12 months after trauma demonstrated good graft osteointegration and fracture healing (Figure 8) with restoration of normal range of motion (ROM) (Figure 9).




Figure 8. X-rays 12 months after trauma.



Figure 9. Clinical view 1 year after trauma.

Discussion

Tibial plateau fractures are common and can be due to high- or low energy traumas and can affect young adults or third-age patients (11-13). They represent 1% of all fractures (14). Their incidence progressively raised over the years mainly due to the increase of road, sports and work accidents and of high energy traumas (15).

The anatomy of the tibial plateau combined with high energy trauma produce complicated injury patterns with involvement of metaphysis, articular comminution and, frequently, with loss of integrity of the soft-tissue envelope (16). The severity of the trauma may also produce associated disruption of primary and secondary knee stabilizers (17,18). In a 2010 study, Stannard et al. reported an incidence of torn ligaments following tibial plateau fractures, as high as 85% and 79% in type V and VI respectively (17). Furthermore, this type of injury is also becoming more and more prevalent in the elderly, as consequence of low energy falls and osteoporosis (19). In such cases, soft tissue damage arises from the delicacy of the skin.

For these reasons complex fractures of the tibial plateau (Schatzker type IV, V and VI) are considered severe lesions even for experienced surgeons.

In 1916 Sever (20) was the first to describe three cases of condyle tibial fractures. Since then, numerous publications and case studies have been reported and all agree that the consequence of the severity of the trauma and of an inadequate treatment can be serious (21-25).

Soft tissue problems in fractures around the knee is of crucial importance. One should think that the fracture will not change but soft tissue will. In high-energy traumas, fractures should be considered as “substantial soft-tissue injuries with a broken bone inside” (12,26). As consequence, compartment syndrome can be a devastating complication of such traumas and its incidence can vary from 17% to 18.7% in closed and open fractures respectively (12). In particular, Schatzker V and VI fractures have a notoriously high incidence of compartment syndrome that can reach 30.4% for type VI in some studies (27,28) and are often accompanied by peripheral nerve lesions.

In this context it is very important the correct timing of surgery that inevitably adds further traumatism to soft tissues; therefore, the management in the early

stages of treatment should focus on preventing their further injury while waiting to repair definitively the fracture (12). The optimal temporizing treatment in initial damage control is spanning external fixation (29). Spanning external fixators reduce fracture fragments via ligamentotaxis, along with providing pain relief and a stable environment for soft tissue healing, as well as early mobilization of the patient.

It is also important the distinction of tibial plateau fractures into complex and simple that allows to identify the more serious or potentially serious injuries, whose treatment is more difficult and more frequently characterized by complications.

CT is of great value for determining the location and magnitude of the joint depression, enabling greater precision of preoperative planning, while 3D reconstructions provide an estimation of metaphyseal bone loss, of articular comminution and joint depression. The importance of CT evaluation was best demonstrated through the identification of postero-medial and postero-lateral shear fractures as a distinct subtype of complex bicondylar tibial fracture prone to be missed by antero-posterior radiograph (30).

Optimal treatment is still open to debate. Non-operative management has historically been the preferred treatment for such fractures. However, surgery is now the preferred method of treatment for displaced fractures (31). A wide choice of fixation techniques can be used (ORIF, minimally invasive plate osteosynthesis, arthroscopically assisted fixation, external fixation) but open reduction with internal fixation seems to be the best therapeutic option in complex fractures (Schatzker IV,V,VI). In any case the goal of treatment is to obtain limb axis restoration, joint stability and accurate intra-articular reduction, avoiding further soft tissue damages (3,4). One of the major concerns of traditional ORIF in tibial plateau fractures is the wound infection, especially in high-energy tibial plateau fractures, in which deep infection and wound dehiscence is described in up to 87.5% of cases (31,32). For minimizing this complication the use of minimally invasive osteosynthesis and a ring fixator (EFMO) has been advocated. Despite EFMO is usually related to a worse articular reduction results reported are encouraging. A randomized study comparing ORIF versus EFMO in type C fractures showed that EFMO was associated with less blood loss,

fewer unplanned re-operations and a shorter hospital stay without differences in 2 years functional results between groups (33). A recent review by McNamara comparing ORIF and EFMO did not find enough evidence to ascertain the best method of fixation but he concluded that current evidence does not contradict the idea of the best results obtained when using limited exposures to treat these fractures (34).

Furthermore, arthroscopy-assisted minimal invasive surgery is now an attractive option among available surgical treatments. The advantages are direct visualization of intra-articular fractures, more accurate reduction, reduced morbidity in comparison with arthrotomy, possibility to diagnose and treat associated meniscal and ligamentous injuries and removal of loose fragments (31,35,36). Arthroscopic assisted treatment has been widely accepted as a safe method for the treatment of Schatzker I-IV fractures, but still remains controversial regarding V and VI injuries for several authors (37-39) because of the high danger of iatrogenic compartment syndrome from fluid extravasation even though its occurrence is reported extremely rarely in the literature (38,40).

This case report summarizes all it has been explained in this discussion (high-energy trauma, complex fracture, compartment syndrome, nerve impairment and wound infection). Despite all these negative variables, the result was satisfactory. Authors believe that this positive outcome depended on the precise management of the lesions both for the therapeutic strategy and for its correct timing.

Conclusions

High-energy tibial plateau fractures are complex lesions that may be associated with extensive soft tissue damages and severe complications. The success of their treatment depends on the precise surgical procedure and on its correct timing. Soft tissues evaluation is as important as bone care. Minimally invasive techniques may improve results.

Conflict of interest: Each author declares that he or she has no commercial associations (e.g. consultancies, stock ownership, equity interest, patent/licensing arrangement etc.) that might pose a conflict of interest in connection with the submitted article.

References

1. Prasad GT, Kumar TS, Kumar RK, et al. Functional outcome of Schatzker type V and VI tibial plateau fractures treated with dual plates. *Indian J Orthop* 2013; 47: 188e194.
2. Zeltser DW, Leopold SS. Classifications in brief: Schatzker classification of tibial plateau fractures. *Clin Orthop Relat Res* 2013; 471(2): 371-4.
3. Sciadini MF, Sims SH. Proximal tibial intra-articular osteotomy for treatment of complex Schatzker type IV tibial plateau fractures with lateral joint line impaction: description of surgical technique and report of nine cases. *J Orthop Trauma* 2013; 27: e18e23.
4. Hill AD, Palmer MJ, Tanner SL, et al. Use of continuous passive motion in the postoperative treatment of intra-articular knee fractures. *J Bone Joint Surg Am* 2014; 96e118.
5. Ozkaya U, Parmaksizoglu AS. Dual locked plating of unstable bicondylar tibial plateau fractures. *Injury* 2015; 46(Suppl 2): S9eS13.
6. Ahearn N, Oppy A, Halliday R, et al. The outcome following fixation of bicondylar tibial plateau fractures. *Bone Jt J* 2014; 96-B: 956e962.
7. Zura RD, Adams SB Jr, Jeray KJ, et al. Timing of definitive fixation of severe tibial plateau fractures with compartment syndrome does not have an effect on the rate of infection. *J Trauma* 2010; 69: 1523.
8. Hak DJ, Lee M, Gotham DR. Influence of prior fasciotomy on infection after open reduction and internal fixation of tibial plateau fractures. *J Trauma* 2010; 69: 886.
9. Catagni MA, Ottaviani G, Maggioni M. Treatment strategies for complex fractures of the tibial plateau with external circular fixation and limited internal fixation. *J Trauma* 2007; 63: 1043-53.
10. Jeremy AH, Murray JB, Michael DM. The Canadian Orthopaedic Trauma Society. Open reduction and internal fixation compared with the circular fixator. Application for bicondylar tibial plateau fractures. *Surg Technol J Bone Jt Surg Am* 2009; 91: 74-88.
11. Bove F, Sala F, Capitani P, Thabet AM, Scita V, Spagnolo R. Treatment of fractures of the tibial plateau (Schatzker VI) with external fixators versus plate osteosynthesis. *Injury* 2018; 49 Suppl 3: S12-S18.
12. Prat-Fabregat S, Camacho-Carrasco P. Treatment strategy for tibial plateau fractures: an update. *EFORT Open Rev* 2017 Mar 13; 1(5): 225-32.
13. Frattini M, Vaienti E, Soncini G, Pogliacomini F. Tibial plateau fractures in elderly patients. *Chir Organi Mov* 2009 Dec; 93(3): 109-14.
14. Hung SS, Chao EK, Chan YS, Yuan LJ, Chung P-H, Chen CY. Arthroscopically assisted osteosynthesis for tibial plateau fractures. *J Trauma* 2003; 54: 356-63.
15. Bengner U, Johnell O, Redlund-Johnell I. Increasing incidence of tibia condyle and patella fractures. *Acta Orthop Scand* 1986; 57: 334-6.
16. Kokkalis ZT, Iliopoulos ID, Pantazis C, Panagiotopoulos E. What's new in the management of complex tibial plateau fractures? *Injury* 2016 Jun; 47(6): 1162-9.
17. Gardner MJ, Yacoubian S, Geller D, et al. The incidence of soft tissue injury in operative tibial plateau fractures: a magnetic resonance imaging analysis of 103 patients. *J Orthop Trauma* 2005; 19(2): 79-84.
18. Stannard JP, Lopez R, Volgas D. Soft tissue injury of the knee after tibial plateau fractures. *J Knee Surg* 2010; 23(4): 187-92.
19. Ebraheim NA, Sabry FF, Haman SP. Open reduction and internal fixation of 117 tibial plateau fractures. *Orthopedics* 2004; 27(12): 1281-7.
20. Sever JW. Fractures of tuberosities of the tibia: A report of three cases. *Am J Orthop Surg* 1916; 14: 299.
21. Luo CF, Sun H, Zhang B, Zeng BF. Three-column fixation for complex tibial plateau fractures. *J Orthop Trauma* 2010 Nov; 24(11): 683-92.
22. Kfuri M, Schatzker J, Castiglia MT, Giordano V, Fogagnolo F, Stannard JP. Extended Anterolateral Approach for Complex Lateral Tibial Plateau Fractures. *J Knee Surg* 2017 Mar; 30(3): 204-11.
23. Menghi A, Mazzitelli G, Marzetti E, Barberio F, D'Angelo E, Maccauro G. Complex tibial plateau fractures: a retrospective study and proposal of treatment algorithm. *Injury* 2017 Oct; 48 Suppl 3: S1-S6.
24. Subramanyam KN, Tammanaiah M, Mundargi AV, Bhoskar RN, Reddy PS. Outcome of complex tibial plateau fractures with Ilizarov external fixation with or without minimal internal fixation. *Chin J Traumatol* 2019 Jun; 22(3): 166-71.
25. Chouhan DK, Chand Saini U, Kumar Rajnish R, Prakash M. Complex bicondylar tibial plateau fractures with reversed tibial slope - Our experience with a fracture-specific correction strategy. *Trauma Case Rep* 2019 Nov 28; 25: 100256.
26. Lowe JA, Tejwani N, Yoo B, Wolinsky P. Surgical techniques for complex proximal tibial fractures. *J Bone Joint Surg Am* 2011 Aug 17; 93(16): 1548-59.
27. Chang YH, Tu YK, Yeh WL, Hsu RW. Tibial plateau fracture with compartment syndrome: a complication of higher incidence in Taiwan. *Chang Gung Med J* 2000; 23(3): 149-55.
28. Ziran BH, Becher SJ. Radiographic predictors of compartment syndrome in tibial plateau fractures. *J Orthop Trauma* 2013; 27(11): 612-5.
29. Zura RD, Browne Ja, Black MD, Olson SA. Current management of high-energy tibial plateau fractures. *Curr Orthop* 2007; 21(3): 229-35.
30. Berber R, Lewis CP, Copas D, Forward DP, Moran CG. Postero-medial approach for complex tibial plateau injuries with a postero-medial or postero-lateral shear fragment. *Injury* 2014; 45(4): 757-65.
31. Chan YS. Arthroscopy assisted surgery for tibial plateau fractures *Chang Gung Med J* May-Jun 2011; 34(3): 239-47.
32. Young MJ. Complications of internal fixation of tibial plateau fractures. *Orthop Rev* 1994; 23: 149-54.
33. Hall JA, Beuerlein MJ, McKee MD. Open reduction and internal fixation compared with circular fixator application for bicondylar tibial plateau fractures. *Surgical technique*;

- Canadian Orthopaedic Trauma Society. *J Bone Joint Surg Am* 2009 Mar 1; 91 Suppl 2 Pt 1:74-88.
34. McNamara IR, Smith TO, Shepherd K et al. Surgical fixation methods for tibial plateau fractures. *Cochrane Database Syst Rev* 2015 Sep 15; (9): CD009679.
35. Pogliacomi F, Verdano MA, Frattini M, Costantino C, Vaienti E, Soncini G. Combined arthroscopic and radioscopic management of tibial plateau fractures: report of 18 clinical cases. *Acta Biomed* 2005 Sep; 76(2): 107-14.
36. Leigheb M, Rusconi M, De Consoli A, et al. Arthroscopically-assisted Reduction and Internal Fixation (ARIF) of tibial plateau fractures: clinical and radiographic medium-term follow-up. *Acta Biomed* 2020 May 30; 91(4-S): 152-9.
37. Burdin G. Arthroscopic management of tibial plateau fractures: surgical technique. *Orthop Traumatol Surg Res* 2013; 99(1 Suppl): S208-18.
38. Herbert M, Domnick C, Petersen W. Arthroscopic treatment of tibial plateau fractures. *Oper Orthop Traumatol* 2014; 26(6): 573-88.
39. Siegler J, Galissier B, Marcheix PS, Charissoux JL, Mabit C, Arnaud JP. Percutaneous fixation of tibial plateau fractures under arthroscopy: a medium term perspective. *Orthop Traumatol Surg Res* 2011; 97(1): 44-50.
40. Kiefer H, Zivaljevic N, Imbriglia JE. Arthroscopic reduction and internal fixation (ARIF) of lateral tibial plateau fractures. *Knee Surg Sports Traumatol Arthrosc* 2001; 9(3): 167-72.

Received: 10 October 2020

Accepted: 20 November 2020

Correspondence:

Alessio Pedrazzini

Orthopaedic Unit, Oglio Po Hospital, Vicomoscato (CR), Italy
Tel. 0039 3478685689

Email: alessiopedrazzini@hotmail.com