

Recovery of gastric function in patients affected by chronic atrophic gastritis using L-cysteine (Acetium®): one year survey in comparison with a control group

Francesco Di Mario¹, Pellegrino Crafa¹, Simone Grillo¹, Stefano Landi¹, Marilisa Franceschi², Kryssia Rodriguez Castro², Antonio Tursi³, Giovanni Brandimarte⁴, Lorella Franzoni¹

¹ Department of Medicine and Surgery, University of Parma; Parma, Italy; ² Endoscopy Unit, Department of Medicine, ULSS7 Pedemontana, Hospital AltoVicentino, Santorso (VI), Italy; ³ Territorial Gastroenterology Service, Azienda Sanitaria Locale Barletta-Andria-Trani, Andria, Italy; ⁴ Division of Internal Medicine and Gastroenterology, Cristo Re Hospital, Rome, Italy

Abstract. *Background and aim:* Chronic Atrophic Gastritis (CAG) is a precancerous condition for gastric cancer (GC) as single risk factor, being a consequence of a previous *Helicobacter pylori* (Hp) infection or based on autoimmune mechanisms. Achlorhydria plays an important role towards the formation of a class I carcinogen, acetaldehyde, after food intake. L-cysteine has been claimed to be able to bind in a covalent way acetaldehyde when administered at meals. *Methods:* In this study we enrolled two CAG groups of patients, one treated with 300 mg/daily of L-cysteine for one year, the other one untreated. We assessed gastric function lasting the one-year follow-up by using noninvasive surrogates, i.e. Pepsinogen I (PGI) and gastrin 17 (G17). *Results:* In the group of 77 CAG on therapy we found a statistically significant increase in PGI values and a decrease in G17 levels, in comparison with unchanged values in control group. *Conclusions:* L-cysteine seems able to provide a recovery in gastric function when administered in CAG patients and could be proposed as a possible therapy in such patients. (www.actabiomedica.it)

Key words: Pepsinogen I, Gastrin 17, Chronic Atrophic Gastritis, L-cysteine, Acetaldehyde

Introduction

Chronic atrophic gastritis (CAG) is a precancerous condition for gastric cancer (GC) as single risk factor (1-7). *Helicobacter pylori* (H.p.) infection is claimed to be the etiological factor in the development of chronic gastritis leading, lasting the years in some subjects, to CAG and then GC. (8-12). A different mechanism to develop CAG is based on autoimmune response targeted against parietal cells and intrinsic factor, typically located only in the fundus and in the corpus (13-15). The risk of GC increases with the extension of Intestinal Metaplasia (IM) as well as the severity of CAG (16). Among the patients with severe atrophy in the antrum, this risk is higher than in healthy subjects and the risk increases consequently if

severe atrophy is found either in both antrum and corpus (severe panCAG) (10, 11, 17, 18).

The prevalence of CAG and GC increases with increasing age and the risk for both diseases is highest among the subjects >45 years old (19, 20). The Correa's cascade represents a most accepted sequence of events from normal mucosa to gastric cancer, a multi-step gastric carcinogenesis including the progression from normal mucosa through chronic non-atrophic gastritis, atrophic gastritis and intestinal metaplasia to intraepithelial neoplasia (previously called dysplasia) and finally to carcinoma (1). On one hand, this long-lasting process of carcinogenesis represents an opportunity to elaborate a screening strategy to identify early stage, i.e. precancerous condition, on the other hand a difficulty is represented by the fact that the majority of

patients affected by CAG are asymptomatic or experiencing not specific manifestations like early satiety, epigastric fullness or bloating (13, 21-23).

The Updated Sydney System (USS) is the most widely accepted as sample protocol for diagnosis of CAG (24), being the OLGA (operative link for gastritis assessment) and/or OLGIM (operative link on gastric intestinal metaplasia) system, currently of reference for staging of gastritis (25, 26).

A proposed surrogate to investigate by non-invasive way both function and morphology of gastric mucosa is represented by serum pepsinogens (so called “serological biopsy”) (27).

Serum pepsinogens (PGI and PGII) were proposed in the diagnosis of gastric diseases in dyspeptic patients in general population (4, 28-30) and in selected patients (31). Moreover, Gastrin 17 (G17) has been proposed beside pepsinogens to verify the maintenance of negative feedback between acid and gastrin, critically modified in the picture of CAG, because of important modification in acid secretion (32-34). In particular, in body atrophic gastritis, in which atrophy leads to achlorhydria, G17 increases significantly, being the serological spectrum represented by low levels of PGI and high levels of gastrin (23).

Several meta-analyses (35-37) pointed out the properties of serum pepsinogens in the diagnosis of CAG, due to very high negative predictive value.

Based on the literature, pepsinogens were recommended by international experts for screening and diagnosis of CAG (10, 11).

In the literature the majority of studies documented an improvement of CAG and in some case of IM following H.p. eradication (38-41).

Di Mario by using serology, also demonstrated an improvement of the picture of CAG after H.p. eradication, lasting a four-year follow-up (42).

In 2009 IARC declared acetaldehyde as group 1 carcinogen equivalent to many others similar carcinogens (43). Acetaldehyde derives from the metabolism of ethanol through oxidation and it is an important downstream of tobacco smoke (44-46). The evidence for the carcinogenesis of acetaldehyde derives from several epidemiological studies on selected populations of alcohol drinkers and alcohol dehydrogenase (ADH) deficiency, as well as low aldehyde dehydrogenase ac-

tivity (47). The accumulation of acetaldehyde. In the saliva, in turn represents an increased risk for different upper gastrointestinal tract neoplasms (48, 49). In the mouth, it has been demonstrated the dissolution of acetaldehyde into the saliva during smoking (50). The carcinogenic effect of tobacco is mediated by noxious aldehydes in saliva at level of oral cavity and subsequently in the larynx, esophagus and stomach (51). Also stomach cancer is claimed to be dependent from the formation of acetaldehyde in gastric lumen through the role of local microbiota. In fact, some microbe strains of the saprophytic oral or intestinal microflora can exert ADH activity to induce acetaldehyde formation from ethanol by oxidation (52, 53). **Patients affected by CAG show some changes in** microbiome composition resulting in a presence of bacteria in gastric lumen producing both local ethanol and acetaldehyde from glucose metabolism (54). After assumption of ethanol, in this scenario, the production of both compounds has been described as a 5-fold increase (55).

Sprince et al. (56), demonstrated about forty years ago the properties of L-cysteine- a non-essential amino-acid – in order to avoid the toxicity of acetaldehyde by creating a covalent binding leading to the formation of a stable 2-methylthiazolidine-4- carboxylic acid (MTCA), which is a non-toxic compound cleared through feces and urine. (56, 57). Other studies focused on the protective role of ascorbic acid in prevention of precancerous condition such as atrophic gastritis (58). In the last decade, Salaspuro et al. (59) demonstrated that compounds slowly releasing L-cysteine locally in the oral cavity were able to successfully remove acetaldehyde from the saliva in alcohol and smokers’ consumers (59, 60). Several studies confirmed the properties of chewing gum containing L-cysteine of removing acetaldehyde from saliva in smokers (61, 62). Other studies demonstrated the capacity of L-cysteine to remove acetaldehyde also from gastric juice in achlorhydric stomach after alcohol administration (63, 64). A recent study, after oral administration of 15% alcohol solution in a total dose of 0.3 g/kg by using L-cysteine capsules shows that the compound was able to eliminate ethanol-derived acetaldehyde from gastric juice of patients with atrophic gastritis and the persistence of the effect at least for 40 min (65).

Aim

This study was designed to evaluate the assessment of gastric function by means of serology (Pepsinogen I and Gastrin 17) in patients with diagnosis of CAG treated for 1 year with 300 mg of L-cysteine daily compared with a control group of CAG untreated patients.

Patient and methods

Data collection

Data were recovered from the database and medical records from one single medical center located in Northeast of Italy. We collected patients affected by CAG, diagnosed by endoscopy and histology. The biopsies were collected according to USS (24) and evaluated according to OLGA grading (25).

Patients with CAG were characterized, according with different etiology: autoimmune gastritis and/or post-Hp infection. Autoimmune gastritis was diagnosed by means of positivity of anti-parietal cell antibodies and corresponding histological criteria. Post-infectious atrophic gastritis was diagnosed by previous confirmation of Hp infection (see Hp status methods) All patients at baseline were Hp negative for at least two years and off PPI therapy in the last six months.

Patients with CAG were subdivided in two groups, according with the intake of L-cysteine (Ace-tium®, Biohit Helsinki, Finland) at dosage of 100 mg capsules during meals, three times a day for 1 year (Group 1) and subjects without therapy, except anti-acid or prokinetics when needed (Group 2). Clinical data (age, sex, etiology of CAG - autoimmune or post HP infection) as well as smoking habit and alcohol intake were collected for each patient. We choose the cut-off of 10 cigarettes/day to define smoking habit and ethanol 40gr/day for positive alcohol intake.

Group 1 included 77 patients (24 males 53 females, mean age 52.7 ± 11.2 ys.) whose gastritis etiology was autoimmune (30 pts.) or Hp. related (47 pts.).

Group 2 included 50 patients (14 males, 36 females, mean age 50 ± 11.9 ys.) with autoimmune (30

patients) or Hp.-related (20 pts.) gastritis. Table 1 summarized the epidemiological data.

In all patients of both groups a serological sample was obtained to measure levels of PG I and G 17 using Elisa method (Biohit, Helsinki, Finland). Normal values are reported as follow: PG I: 30-160 µg/L, G 17: 1-10 pg/L. PGI and G17 levels were measured at baseline and after 3, 6, 12 months.

H.p. status was assessed by means of three methods in each patient: 13C-urea breath test, histology of gastric mucosa (using hematoxylin and eosin with modified Giemsa), serology by IgG antibody (Elisa method; Biohit, Helsinki, Finland)

Inclusion criteria

Only Hp negative patients, investigated by UBT, serology and histology of gastric mucosa, were enrolled in the study; age: >18 ys. < 75 ys. old. Only patients who have a biopsy confirmed CAG with corpus restricted disease or with pan-atrophy according to the OLGA staging or Sydney system were included in the study. Some subjects of both groups had an history of Hp infection previously successfully eradicated (at least two years before).

Exclusion criteria

Patients who underwent previous upper gastrointestinal tract surgery; previous history of neoplasms; PPI therapy in the last year.

Table 1. Demographics

| | Group 1 | Group 2 | p |
|---|--------------|--------------|--------|
| Patients N° | 77 | 50 | ns |
| Male | 24 (31.16%) | 14 (28.00%) | ns |
| Female | 53 (68.83%) | 36 (72.00%) | ns |
| Mean age | 52.7±11.2 ys | 50 ± 11.9 ys | ns |
| Age range | 37-74 ys | 43-79 ys | ns |
| Smoking habit | 14 (18,18%) | 9 (18,00%) | ns |
| Alcohol intake | 19 (24,68%) | 11 (22,00%) | ns |
| Previous H.p. | 47 (61.03%) | 20 (40.00%) | <0.001 |
| Autoimmune gastritis | 30 (38.96%) | 30 (60.00%) | <0.001 |
| Group 1 CAG treated by L-cysteine (study group) | | | |
| Group 2 untreated CAG (control group) | | | |

Study design

Single center, controlled, not randomized study that included two populations: the first (Group 1) included 77 pts. affected by chronic atrophic gastritis (CAG) diagnosed by histology treated with L-cysteine capsules (100mg three times daily with meals) for 12 months compared to 50 CAG pts. as control group (Group 2) off therapy.

Statistics

Statistical analysis was performed with the SPSS (version 20.1) statistical software program for windows. Chi-squared and Wilcoxon tests for paired data were used when appropriate. Data were expressed as mean for quantitative variables. All p were two-tailed with statistically significance indicated by a value of $p < 0.05$.

The study was performed in accordance with the declaration of Helsinki and approved by local Ethics Committee (Identifier: 92687).

Results

Table 2 summarized serological values of PGI and G17 at baseline, three, six and twelve months after entering the study, in both groups.

Serum levels of PGI and G17 in the two groups lasting 1 years of follow-up were reported in figures 1 and 2

In treated patients (group 1), PGI levels increase from baseline lasting the follow-up period from 7.88 $\mu\text{g/L}$ to 15.92 $\mu\text{g/L}$ at twelfth month, in statistically significant manner ($p < 0.001$). On the contrary, in control group (Group 2), PGI levels stay unchanged lasting the follow-up (baseline: 6.30 $\mu\text{g/L}$ vs twelve months: 7.10 $\mu\text{g/L}$; $p = \text{ns}$) (Fig. 1).

Conversely, G17 values show a decrease from baseline: 30.31 pg/L to 19.23 pg/L at twelve months; $p < 0.001$ in Group 1 opposite to Group 2, showing no significant modifications lasting the follow-up (baseline: 34.52 pg/L ; twelve month: 32.15; $p = \text{ns}$) (Fig. 2).

We stratified all patients, treated and untreated, according with OLGA Grading, in different severity of CAG. The distribution of OLGA scores in Group 1 was: OLGA 1: 10 pts, OLGA 2: 18 pts, OLGA 3: 26 pts, OLGA 4: 23 pts and in Group 2 was: OLGA 1: 5 pts, OLGA 2: 11 pts, OLGA 3: 19 pts, OLGA 4: 15 pts

We decide to group patients showing OLGA 1 and OLGA 2 in "OLGA Low-grade" and patients with OLGA 3 and OLGA 4 in "OLGA High-grade".

Fig. 3 and 4 summarizing values of PGI and G17 from baseline to twelve months, according with the OLGA severity score.

Data showed the lowest levels of PGI in the group of "OLGA High-grade" in comparison with patients showing "OLGA Low-grade", being this feature confirmed in both groups (treated and untreated).

Group 1 OLGA "Low-grade": 10.2 $\mu\text{g/L}$ vs 4.5 $\mu\text{g/L}$ in "OLGA High-grade";

Table 2: PGI and G17 values in Group 1 and Group 2, lasting one year of follow-up

| | Baseline | 3 months* | 6 months** | 12 months*** | p |
|----------------|----------------------|-----------------------|-----------------------|-----------------------|---|
| Group 1 PGI | 7.88 $\mu\text{g/L}$ | 11.51 $\mu\text{g/L}$ | 13.73 $\mu\text{g/L}$ | 15.92 $\mu\text{g/L}$ | * < 0.05 ** < 0.01 *** < 0.001 |
| Group 2 PGI | 6.90 $\mu\text{g/L}$ | 7.20 $\mu\text{g/L}$ | 6.80 $\mu\text{g/L}$ | 7.10 $\mu\text{g/L}$ | n* ns ** ns *** ns |
| Group 1 G17 | 30.31 pg/L | 25.46 pg/L | 21.84 pg/L | 19.23 pg/L | * < 0.01 ** < 0.001 *** < 0.001 |
| Group 2 G17 | 34.52 pg/L | 35.48 pg/L | 35.8 pg/L | 32.15 pg/L | * ns ** ns *** ns |

*3 months vs baseline values; **6 months vs baseline values; *** 12 months vs baseline values; Group 1 CAG treated by L-cysteine (study group); Group 2 untreated CAG (control group)




Figure 1. Comparison of PGI values between Group 1 and Group 2




Figure 2. Comparison of G17 values between Group 1 and Group 2

Group 2 OLGA "Low-grade" 11.3 µg/L vs 5.3 µg/L OLGA "High grade"; $p < 0.001$

Conversely, the highest values of G17 were found in OLGA "High-grade", both in Group 1 and Group 2: 39.7 pg/L and 35.6 pg/L, opposite to levels in OLGA "Low-grade": Group 1: 18.2 pg/L, Group 2 16.7 pg/L; $p < 0.001$.

Figures 3 and 4 summarize the modifications of both PGI and G17 lasting follow-up in the two groups, according with OLGA severity score. Ones more, treated patients show a statistically significant modification from baseline to twelfth month, of par-

ticular relevance as regards G17 in the group of OLGA "High-grade": baseline 39.7 pg/L vs 21.8 pg/L.

Discussion

Until now, Cag is lacking a reliable therapy, being the only suggestion to provide adequate follow up for such condition, considered at risk to develop GC (1-4).

From a functional point of view CAG is characterized by a low level of chlorohydric acid (hypochloridria) or – in the most severe cases – achlorhydria, in turn related with increased risk to develop GC (10).




Figure 3. PGI values according with OLGA grading




Figure 4. G17 values according with OLGA grading

The serological phenotype of body atrophic gastritis is represented by the picture of low levels of PGI (a surrogate of chlorohydric acid secretion) and high levels of G17 (due to negative feedback with acid) (23).

By using OLGA severity score, as expected, patients affected by OLGA 3 and 4 showed lowest levels of PGI and highest values of G17, in comparison with low severity scores (OLGA1 and 2).

The main result of the study is represented by a feature of a statistically increase of PGI levels in pa-

tients treated with L-cysteine, while the values of untreated CAG patients in control group remain unchanged, lasting the entire follow-up period. More important seems to be the critical decrease of G17 levels in treated patients from 30.31 pg/L at baseline to 19.23 at 12th month: in particular, in OLGA 3 and 4 groups, the decrease was from 39.7 pg/L at baseline to 21.8 at the end of follow-up. Interestingly, the major modification of both markers, PGI and G17, was observed after 3 months from the intake of L-cysteine

and remain stable, lasting the other 9 months of follow-up, suggesting possibly a hypertrophy of gastric glands, stimulated by the compound, more than an increase “de novo” of the number of glands (hyperplasia).

As we know, achlorhydria plays a critical role in carcinogenetic mechanisms in gastric mucosa, being acetaldehyde the most important factor involved in such transformation from CAG to adenocarcinoma (44,45, 46). In this view the recovery of gastric secretion represented by an increase of PGI levels could be of importance in limiting acetaldehyde production into the stomach after food ingestion, and in particular alcohol beverages (52, 53).

Similarly, the decrease in G17 levels lasting one year follow-up period after daily administration of 300 mg of L-cysteine, is important, considering the role of the hormone as a growth factor involved in pre-neoplastic conditions and in NEN (10).

In literature is generally accepted that CAG represents an irreversible stage of change for gastric mucosa, but recently some authors hypothesized a possible recovery of CAG picture (10), being the debate at the present open. The increase in PGI levels coupled with the decrease in G17 values seems support the possibility of a recovery of CAG at least as regards gastric function.

To prove this assumption, we need the support of histology, before and last the treatment, which is lacking in the present study, being, probably one year of follow-up, a too short interval to observe modifications in gastric morphology. The main purpose to administered L-cysteine in CAG patient was based on the ability of the compound in binding in a covalent way acetaldehyde in achlorhydric stomach, proved in both experimental and clinical studies (56,57,59). The modifications of acetaldehyde levels in gastric juice, was not investigated in the present study, but the feature of a recovery of gastric function in subjects treated with L-cysteine was totally unexpected, and to our knowledge, never documented, until now, in a full paper.

Strength points: all patients entered the study were consecutively enrolled and histologically diagnosed to confirm CAG, according to OLGA severity score, following a standardize biopsy sampling. The control group (untreated patients) was recruited in

same geographical area and diagnosed by same endoscopic and pathological team and was comparable as regards epidemiological factors.

The values of surrogate markers for gastric function (PGI and G17) were coherent at baseline with a picture of CAG, as well as the severity of OLGA scores and remain unchanged – as expected – in control group lasting all the follow-up period.

Weakness points: for sure the lack of histology to support the hypothesis of gastric function recovery following L-cysteine administration, represents the major weakness of the study and needs to confirm in next studies, hopefully based on a longest follow-up period (at least two years).

Additionally, we don't provide, in this study, any data on modifications in acetaldehyde values in gastric mucosa in both groups.

Conclusion

CAG is a precancerous condition in which – at present – the only suggestion is to propose a correct follow-up according to severity of picture. This study aimed to provide an indication of therapy for such patients by using L-cysteine - daily administered – based on the properties of this compound to bind in an irreversible way acetaldehyde in achlorhydric stomach and seems suggest a possible recovery in gastric function lasting therapy by using validated serologic markers.

Next studies are necessary to confirm these results by performing histology of gastric mucosa.

Conflict of Interest: Each author declares that he or she has no commercial associations (e.g. stock ownership, equity interest, patent/licensing arrangement etc.) that might pose a conflict of interest in connection with the submitted article

References

1. Correa P. Human gastric carcinogenesis: a multistep and multifactorial process—First American Cancer Society Award Lecture on Cancer Epidemiology and Prevention. *Cancer Res* 1992;52:6735-40.
2. Correa P. The biological model of gastric carcinogenesis. *IARC Sci Publ.* 2004;157:301-10.

3. de Vries AC, van Grieken NC, Looman CW, et al. Gastric cancer risk in patients with premalignant gastric lesions: a nationwide cohort study in the Netherlands. *Gastroenterology*. 2008 Apr; 134(4): 945–52.
4. Agreus L, Kuipers EJ, Kupcinskas L, et al. Rationale in diagnosis and screening of atrophic gastritis with stomach-specific plasma biomarkers. *Scand J Gastroenterol*. 2012;47:136–47.
5. Park YH, Kim N. Review of atrophic gastritis and intestinal metaplasia as a premalignant lesion of gastric cancer. *J Cancer Prev*. 2015;20:25–40.
6. Walter IR, Strickland RG, Ungar B, Mackay IR. Simple atrophic gastritis and gastric carcinoma. *Gut* 1971; 12:906–11.
7. Sipponen P, Kekki M, Haapakoski J et al. Gastric cancer risk in chronic atrophic gastritis: statistical calculations of cross-sectional data. *Int. J. Cancer* 1985; 35: 173–7.
8. Valle J, Kekki M, Sipponen P et al. Long-term course of a 32 year follow-up study. *Scand. J. Gastroenterol*. 1996; 31:546–50
9. Niemelä S, Karttunen T, Kerola T. Helicobacter pylori-associated gastritis. Evolution of histologic changes over 10 years. *Scand. J. Gastroenterol*. 1995; 30: 542–9.
10. Sugano K, Tack J, Kuipers EJ, et al. Faculty members of Kyoto Global Consensus Conference. Kyoto global consensus report on Helicobacter pylori gastritis. *Gut*. 2015 Sep;64(9):1353–67.
11. Malfertheiner P, Megraud F, O'Morain CA, et al. European Helicobacter and Microbiota Study Group and Consensus panel. Management of Helicobacter pylori infection—the Maastricht V/Florence Consensus Report. *Gut*. 2017 Jan;66(1):6–30.
12. Uemura N, Okamoto S, Yamamoto S, Matsumura N, Yamaguchi S and Yamakido M: Helicobacter pylori infection and the development of gastric cancer. *N Engl J Med* 2001; 345: 829–832.
13. Rugge M, Savarino E, Sbaraglia M, Bricca L, Malfertheiner P. Gastritis: The clinico-pathological spectrum. *Dig Liver Dis*. 2021 Mar 27:S1590–8658(21)00100–6.
14. Coati I, Fassan M, Farinati F, Graham DY, Genta RM, Rugge M. Autoimmune gastritis: Pathologist's viewpoint. *World J Gastroenterol*. 2015 Nov 14;21(42):12179–89.
15. Rugge M, Fassan M, Pizzi M, et al. Autoimmune gastritis: histology phenotype and OLGA staging. *Aliment Pharmacol Ther* 2012; 35: 1460–1466
16. Vannella L, Lahner E, Annibale B. Risk for gastric neoplasias in patients with chronic atrophic gastritis: a critical reappraisal. *World J Gastroenterol* 2012; 18: 1279–1285
17. Rugge M, Sugano K, Scarpignato C, Sacchi D, Oblitas W J, Naccarato AG. Gastric cancer prevention targeted on risk assessment: Gastritis OLGA staging. *Helicobacter*. 2019 Apr;24(2):e12571.
18. Rugge M, Genta RM, Fassan M, et al. OLGA Gastritis Staging for the Prediction of Gastric Cancer Risk: A Long-term Follow-up Study of 7436 Patients. *Am J Gastroenterol*. 2018 Nov;113(11):1621–1628.
19. Fock KM. Review article: the epidemiology and prevention of gastric cancer. *Aliment Pharmacol Ther*. 2014; Vol. 40, 3, pp 250–260.
20. Hyuna Sung PhD, Jacques Ferlay MSc, ME, Rebecca L. Siegel MPH, et al: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries CA: A Cancer Journal for Clinicians 2021; Vol. 71, 3, pp 209–249
21. Carabotti M, Lahner E, Esposito G, Sacchi MC, Severi C, Annibale B. Upper gastrointestinal symptoms in autoimmune gastritis: a cross-sectional study. *Medicine (Baltimore)*. 2017;96:e5784.
22. Tosetti C, Stanghellini V, Tucci A, et al. Gastric emptying and dyspeptic symptoms in patients with nonautoimmune fundic atrophic gastritis. *Dig Dis Sci*. 2000;45:252–7.
23. Lahner E, Conti L., Annibale B., Corleto V.D. Current Perspectives in Atrophic Gastritis. *Curr Gastroenterol Rep*. 2020 Jun 15;22(8):38.
24. Dixon MF, Genta RM, Yardley JH, et al. Classification and grading of gastritis: the updated Sydney system. *Am J Surg Pathol*. 1996; 20 (10):1161–1181.
25. Rugge M., Meggio A., Pennelli G., et al. Gastritis staging in clinical practice: the OLGA staging system. *Gut*. 2007 May;56(5):631–636
26. Capelle L.G., de Vries A.C., Haringsma J., et al. The staging of gastritis with the OLGA system by using intestinal metaplasia as an accurate alternative for atrophic gastritis. *Gastrointest Endosc*. 2010 Jun;71(7):1150–8.
27. Samloff IM, Varis K, Ihama K, Siurala M, Rotter JI. Relationships among serum pepsinogen I, serum pepsinogen II, and gastric mucosal histology. A study in relatives of patients with pernicious anemia. *Gastroenterology*, 1982; 83 (1 Pt 2):204–9.
28. Sjomina O, Pavlova J, Daugule I, et al. Pepsinogen test for the evaluation of precancerous changes in gastric mucosa: a population-based study. *J Gastrointest Liver Dis*. 2018 Mar;27(1):11–17.
29. Germaná, B., et al. Clinical usefulness of serum pepsinogens I and II, gastrin-17 and anti-Helicobacter pylori antibodies in the management of dyspeptic patients in primary care. *Dig. liv dis* 37.7 (2005): 501–508.
30. Ju Yup Lee, Nayoung Kim, Hye Seung Lee, et al. Correlations among endoscopic, histologic and serologic diagnoses for the assessment of atrophic gastritis *J Cancer Prev*. 2014;19(1):47–55
31. Venerito M, Radunz M, Reschke K, et al. Autoimmune gastritis in autoimmune thyroid disease. *Aliment Pharmacol Ther* 2015; 41: 686–93.
32. Sipponen, P., et al. Atrophic gastritis serum levels of amidated gastrin-17 and pepsinogen i in atrophic gastritis: an observational case-control study. *Scand. J. gastroenterol*. 37.7 (2002): 785–791.
33. Väänänen, H., et al. Non-endoscopic diagnosis of atrophic gastritis with a blood test. Correlation between gastric histology and serum levels of gastrin-17 and pepsinogen I: a multicentre study. *Eur j gastroenterol hepatol* 15.8 (2003):

- 885-891.
34. Barchi A, Miraglia C, Violi A, et al. A non-invasive method for the diagnosis of upper GI diseases. *Acta Biomed* 2018;89:40-43.
 35. Dinis-Ribeiro M, Yamaki G, Miki K, et al. Meta-analysis on the validity of pepsinogen test for gastric carcinoma, dysplasia or chronic atrophic gastritis screening. *J Med Screen* 2004; 11: 141-7.
 36. Huang YK, Yu JC, Kang WM, et al. Significance of Serum Pepsinogens as a Biomarker for Gastric Cancer and Atrophic Gastritis Screening: A Systematic Review and Meta-Analysis. *PLoS One* 2015; 10: e0142080.
 37. R M Zagari 1, S Rabitti 1, D C Greenwood 2, L H Eusebi 1, A Vestito 3, F Bazzoli 1 Systematic review with meta-analysis: diagnostic performance of the combination of pepsinogen, gastrin-17 and anti-*Helicobacter pylori* antibodies serum assays for the diagnosis of atrophic gastritis *Aliment Pharmacol Ther* . 2017 Oct;46(7):657-667.
 38. Yamada T, Miwa H, Fujino T, Hirai S, Yokoyama T, Sato N. Improvement of gastric atrophy after *Helicobacter pylori* eradication therapy. *J Clin Gastroenterol* 2003;36:405-410.
 39. Rokkas T, Pistiolas D, Sechopoulos P, Robotis I, Margantinis G. The long term impact of *Helicobacter pylori* eradication on gastric histology: a systematic review and meta analysis. *Helicobacter* 2007;12:32-38.
 40. Toyokawa T, Suwaki K, Miyake Y, Nakatsu M, Ando M. Eradication of *Helicobacter pylori* infection improved gastric mucosal atrophy and prevented progression of intestinal metaplasia, especially in the elderly population: a long term prospective cohort study. *J Gastroenterol Hepatol* 2010;25:544-547.
 41. Hwang YJ, Kim N, Lee HS, et al. Reversibility of atrophic gastritis and intestinal metaplasia after *Helicobacter pylori* eradication a prospective study for up to 10 years. *Aliment Pharmacol Ther* 2018;47:380-390.
 42. Di Mario F, Moussa AM, Dal Bò N, et al. Recovery of gastric function after *Helicobacter pylori* eradication in subjects with body atrophic gastritis: prospective 4-year study. *J Gastroenterol Hepatol*. 2005 Nov;20(11):1661-6
 43. Baan, R., Straif, K., Grosse, Y., et al. Carcinogenicity of alcoholic beverages. *The Lancet Oncology*, 2007, 8(4), 292-293.
 44. Driver, H.E. and Swann, P.F. (1987) Alcohol and human cancer. Anticancer implications in alcohol carcinogenesis. *Cancer Res.*, 39, 2844-2850.
 45. Blot, W.J. (1992) Alcohol and cancer. *Cancer Res.*, 52 (Suppl.), 2119-2123. *J. Natl Cancer Inst.*, 69, 947-951.
 46. Brugere, J., Guenel, P., Leclerc, A. and Rodriguez, J. (1986) Differential effects of tobacco and alcohol in cancer of the larynx, pharynx and mouth. *Cancer*, 57, 391-395.
 47. Breslow, N.E. and Enstrom, J.E. (1974) Geographic correlations between mortality rates, alcohol and tobacco consumption in the United States. *J. Natl Cancer Inst.*, 53, 631-639.
 48. Homann N., Jousimies-Somer H., Jokelainen K., Heine R. and Salaspuro M., High acetaldehyde levels in saliva after ethanol consumption: Methodological aspects and pathogenic implications *Carcinogenesis* vol.18 no.9 pp.1739-1743, 1997
 49. Dong, Y.-J., Peng, T.-K. and Yin, S.-J. (1996) Expression and activities of ethanol: a gastrointestinal barrier against the systemic toxicity of class IV alcohol dehydrogenase and class III aldehyde dehydrogenase in human mouth. *Alcohol*, 13, 257-262.
 50. Jauhonen, P., Baraona, E., Miyakawa, H. and Lieber, C.S. (1982) Origin of breath acetaldehyde during ethanol oxidation. Effect of long-term cigarette smoking. *J. Lab. Clin. Med.*, 100, 908-916. May 19, 1997
 51. Jokelainen, K., Heikkinen, E., Roine, R., Lehtonen, H. and Salaspuro, M. (1996) Increased acetaldehyde production by mouthwashings of patients with oral cavity, laryngeal or pharyngeal cancer. *Alcohol. Clin. Exp. Res.*, 20, 1206-1210.
 52. Jokelainen, K., Roine, R.P., Va'ana'nen, H., Fa'rkkila, M. and Salaspuro, M. (1994) In vitro acetaldehyde formation by human colonic bacteria. *Gut*, 35, 1271-1274.
 53. Jokelainen, K., Siitonen, A., Jousimies-Somer, H., Nosova, T., Heine, R. and (1981) In vitro alcohol dehydrogenase mediated acetaldehyde production by aerobic bacteria representing the normal colonic flora in man. *Alcohol. Clin. Exp. Res.*, 20, 967-972.
 54. Moreno, A. and Pares, X. (1991) Purification and characterization of a new alcohol dehydrogenase from human stomach. *J. Biol. Chem.*, 266, 1128-1233.
 55. Maier, H., Zöller, J., Herrmann, A., Kreiss, M. and Heller, W.D. (1993) Dental status and oral hygiene in patients with head and neck cancer. *Otolaryngol. Head Neck Surg.*, 108, 655-661
 56. Sprince H, Parker C M, Smith G G, Gonzales L J Protective action of ascorbic acid and sulfur compounds against acetaldehyde toxicity: implications in alcoholism and smoking *Agents Actions* 1975 May;5(2):164-73.
 57. Sprince H Protective action of sulfur compounds against aldehyde toxicants of cigarette smoke *Eur J Respir Dis Suppl*. 1985;139:102-12.
 58. Farinati, F., Della Libera, G., Cardin R, et al. Gastric Antioxidant, Nitrites, and Mucosal Lipoperoxidation in Chronic Gastritis and *Helicobacter pylori* Infection. *J Clin Gastroenterol* 1996 - Vol 22 - Issue 4 - p 275-281
 59. Salaspuro V, Hietala J, Kaihovaara P, et al. Removal of acetaldehyde from saliva by a slow-release buccal tablet of L-cysteine. *Int J Cancer*. 2002;97:361-364.
 60. Salaspuro VJ, Hietala JM, Marvola ML, et al. Eliminating carcinogenic acetaldehyde by cysteine from saliva during smoking. *Cancer Epid Biomark Prev*. 2006;15:146-149.
 61. Kartal A, Hietala J, Laakso I, et al. Formulation and in-vivo evaluation of L-cysteine chewing gums for binding carcinogenic acetaldehyde in the saliva during smoking. *J Pharm Pharmacol*. 2007 Oct;59(10):1353-8.
 62. Syrjänen K, Salminen J, Aresvuo U, et al. Elimination of Cigarette Smoke-derived Acetaldehyde in Saliva by Slow-release L-Cysteine Lozenge Is a Potential New Method to Assist Smoking Cessation. A Randomised, Double-blind,

- Placebo-controlled Intervention. *Anticancer Res.* 2016 May;36(5):2297-306
63. Linderborg K, Marvola T, Marvola M, Salaspuro M, Färkilä M, Väkeväinen S. Reducing carcinogenic acetaldehyde exposure in the achlorhydric stomach with cysteine. *Alcohol Clin Exp Res.* 2011 Mar;35(3):516-22.
64. Maejima R, Iijima K, Kaihovaara P, et al. Effects of ALDH2 genotype, PPI treatment and L-cysteine on carcinogenic acetaldehyde in gastric juice and saliva after intragastric alcohol administration. *PLoS One.* 2015 Apr 1;10(4):e0120397.
65. Hellström PM, Hendolin P, Kaihovaara P, et al. Slow-release L-cysteine capsule prevents gastric mucosa exposure to carcinogenic acetaldehyde: results of a randomised single-blinded, cross-over study of Helicobacter-associated atrophic gastritis. *Scand J Gastroenterol.* 2017 Feb;52(2):230-237

Correspondence:

Received: 1 January 2022

Accepted: 24 January 2022

Francesco Di Mario, MD,

Department of Medicine and Surgery,

University of Parma,

Via Gramsci, n. 14, 43126, Parma, Italy,

Tel.: +39 3318133870,

e-mail: francesco.dimario@unipr.it