

## R E V I E W

# Magnetic resonance enterography (MRE) and ultrasonography (US) in the study of the small bowel in Crohn's disease: state of the art and review of the literature

*Rosa Manetta<sup>1</sup>, Ilaria Capretti<sup>2</sup>, Noemi Belleggia<sup>2</sup>, Claudia Marsecano<sup>2</sup>, Angelo Viscido<sup>3</sup>, Federico Bruno<sup>2</sup>, Francesco Arrigoni<sup>2</sup>, Libeng Ma<sup>4</sup>, Giuseppe Guglielmi<sup>5</sup>, Alessandra Splendiani<sup>2</sup>, Ernesto Di Cesare<sup>2</sup>, Carlo Masciocchi<sup>2</sup>, Antonio Barile<sup>2</sup>*

<sup>1</sup>Division of Radiology, S. Salvatore Hospital, L'Aquila, Italy; <sup>2</sup>Department of Biotechnology and Applied Clinical Sciences, University of L'Aquila, L'Aquila, Italy; <sup>3</sup>Department of Life, Health and Environmental Sciences, University of L'Aquila, L'Aquila, Italy; <sup>4</sup>Department of Radiology, GuangDong Pharmaceutical University, Guangzhou, Cina; <sup>5</sup>Department of Clinical and Experimental Medicine, Foggia University School of Medicine, Foggia, Italy

**Summary.** Crohn's disease (CD) is a chronic idiopathic disease and its diagnosis is based on a combination of clinical symptoms, laboratory tests and imaging data. There isn't a diagnostic gold standard: the ileocolonoscopy with mucosal biopsies represents the standard for luminal disease, while cross-sectional imaging such as Ultrasound (US), Computed Tomography (CT) or Magnetic Resonance Imaging (MRI) can show transmural alterations and extraintestinal manifestations. CD is usually diagnosed in the young age and after baseline diagnosis, the patients have to undergo to variable follow-up depending on remission or active disease. The aim of our review is to compare Magnetic Resonance Enterography (MRE) to Ultrasonography (US) in the follow-up of CD. ([www.actabiomedica.it](http://www.actabiomedica.it))

**Key words:** MR enterography, Crohn's disease, IBD, CEUS

## Introduction

Crohn's disease (CD) is a chronic idiopathic disease that is commonly characterized by recurrent gastrointestinal tract inflammation. Patients affected by CD are mainly at reproductive ages and they need frequent follow-up (1)

The disease involves the whole gastrointestinal tract, in particular the small bowel (70%) in the tract of terminal ileum and colon (2); it is a pathology with multifactorial etiology and is more common in Europe and North America (3). It has an incidence in the U.K. of 83 per million people. Symptomatic manifestations may be unspecific. Danese et al. (4) have constructed the Red Flags index to individuate early symptoms and have established a value of 8 as highly predictive of CD diagnosis: chronic diarrhea (>3 bowel


movements and >4-week duration); chronic abdominal pain (>3 months); rectal bleeding; extra-intestinal manifestations. There is a poor correlation between symptomatology and disease severity (5). Frequent complications are intestinal strictures (40% of cases) but also abscesses, phlegmons and fistulas (6). Diagnosis and staging of CD require different diagnostic exams: serological testing (C-reactive protein and fecal calprotectin), clinical and endoscopical evaluation (7, 8), video-capsule endoscopy for proximal small bowel. Endoscopy consists in ileocolonoscopy with biopsies from the terminal ileum and colon in order to confirm the diagnosis (9); esophagogastroduodenoscopy is used for suspected upper tract disease (10). The ileocolonoscopy represents the standard for luminal disease (11). Cross-sectional imaging (MRI, CT and US) techniques, gained large application in

gastrointestinal radiology (12-48); in the setting of inflammatory bowel diseases, they are advised as first line techniques in the diagnosis, staging and follow-up (49, 50). An expert consensus committee from the European Society of Gastrointestinal and Abdominal Radiology (ESGAR) and European Society of Paediatric Radiology (ESPR) has established guidelines for performing these diagnostic techniques, including patient preparation, technical recommendations and scan protocol (51). Radiological examinations can evaluate accurately severity; they are non-invasive and not limited to the colon and terminal ileum (52) but can demonstrate complications of CD and extraintestinal manifestations. Aim of this review is to compare Magnetic Resonance Enterography (MRE) with Ultrasonography (US) as non-invasive, radiation-free and appropriate techniques for follow-up in patients affected by CD. In particular, dynamic imaging techniques can assess the degree of disease activity and the efficacy of treatment .


### **Magnetic resonance enterography**

Magnetic resonance imaging is one of the most largely used imaging tool in many diagnostic (53-57) and interventional settings (58, 59) thanks to its intrinsic excellent soft tissue contrast and the absence of ionizing radiation compared to CT (55, 60-87). Magnetic resonance enterography (MRE) plays an important role in supporting the diagnosis, and in establishing severity and presence of penetrating or extra-intestinal disease. To date, MRE is the most employed technique to assess the response to medical or surgical treatment . Common complications of Crohn's disease are intestinal strictures that may be of fibrotic, inflammatory and mixed types (88). The distinction is important in treatment planning; therefore, an appropriate diagnosis of inflammatory stenosis is required. The inflammatory stenosis responds to medical therapy while the fibrotic one requires endoscopic approach or intestinal resection. The exact distinction is sometimes difficult to be determined (89, 90). In a recent review of 2016, Westerland et al (90) analyze advantages and appropriate sequences that allow evaluation of the intestinal wall and distinction between inflammatory or fi-

brotic component. MRE is a panoramic technique that shows intestinal wall layers, presence of penetrating disease and extraluminal complications such as fistulas or abscesses. Important factors during MRE are: optimal distension of the small bowel, obtained with biphasic agent (mannitol or polyethylene glycol); antiperistaltic agent to minimize bowel peristalsis (91, 92). Conventional sequences, used to evaluate typical pathological alterations, are True-FISP or FIESTA (true fast imaging with steady-state free-precession), T2-weighted and T1-weighted fat-saturated sequences. FIESTA sequences are less sensible to motion artifact and show good contrast between the bowel wall, lumen and mesenteric fat. It is useful for a morphological evaluation of the bowel loops (93) (Fig. 1). T2-weighted sequences have a high sensitivity (83%-91%) and specificity (86%-100%) to detect inflamed bowel with a mural thickness greater than 3mm and T2-signal increased for the presence of mural and mesenteric edema. Other features of acute disease are: mesenteric vascular prominence (comb sign), hyperenhancing and enlarged lymph nodes (short axis up to 8 mm) (Fig. 2). In T1-weighted fat-saturated sequences acquired after contrast media, the active inflammatory appears with a stratified pattern of enhancement (hyper-intense mucosa and serosa and hypointense/edema submucosa) (Fig. 3). The chronic inflammation is characterized by T2 hypointense mural signal without adjacent mesenteric inflammation and homogeneous transmural enhancement on T1-weighted fat-saturated sequences (92) (Fig. 4). The distinction between inflammatory and fibrostenotic disease is facilitated by the peculiar tissue contrast obtained in MR images . Recently, Diffusion Weighted Imaging (DWI) has been added to conventional sequences and many studies (94) show that there is a restricted diffusion in active inflammation: the presence of a cellular inflammatory response in the bowel wall prevents the movement of water molecules, showing mucosal hyperintensity on DWI and corresponding low signal on the Apparent Diffusion Coefficient (ADC) map. A quantitative evaluation (for example through ADC) helps to determine the severity of the disease and to monitor treatment response: Rimola et al. (95, 96) have elaborated the MR index of activity (MaRIA), considering MR parameters such as wall thickness, relative contrast enhancement, presence



**Figure 1.** 14-year-old patient with abdominal pain, diarrhoea and elevated inflammatory biomarkers levels. Axial (a) and coronal (b) FIESTA sequences show active inflammation with increased wall thickness of the terminal ileum (arrows)




**Figure 2.** T2-weighted sequences show enlarged lymph nodes (a) and mural edema (b)

of edema, ulcers, pseudopolyps and enlarged lymph nodes. This index has a significant correlation with the endoscopic findings and, though being frequently quoted, it is not diffusely accepted in the clinical practice. In a study carried out in 2017, Kim et al (97) demonstrated that modified MaRIA scoring is quite accurate, because DWI, enteric and portal phase scans can improve reproducibility of the scoring system.

### Ultrasonography

Ultrasonography (US) is the least invasive imaging examination, well tolerated by patients. This technique does not employ ionizing radiations, is repeatable with high diagnostic accuracy and is widely used in the diagnostic setting and as guidance in many interventional radiology procedures (61, 63, 98-105).




**Figure 3.** T1-w fat saturated images in axial (a) and coronal (b) planes, after contrast injection, show hyperintense mucosa (arrows) during the arterial phase

US can identify morphological features of bowel wall drawing the layer pattern and wall thickness ( $>3$  mm in CD) (106) (Fig. 5). For this reasons, US is useful during active CD to show inflammatory processes of angiogenesis and hypervascularization into the intestinal wall (107); computed tomography (CT) and MR are unsuitable for repeated follow-up due to radiation exposure (CT) and high cost and complexity (MR) (108, 109). US can evaluate parietal vascularization using Color or Power Doppler: active disease is characterized by the increased number and caliber of vessels (110). Contrast-enhanced US is more accurate (111). The introduction of oral contrast solution (water and polyethylene glycol) to distend the bowel can improve the image quality and diagnostic accuracy in the evaluation of mild intestinal damage (112); the use of intravenous contrast media in US (CEUS) can emphasize the presence of an increased bowel wall vascularization that it is typical in the active form of CD. US with oral contrast agent (SICUS), as reported by Mocci et al. (113) in a review of 2017, has a sensitivity ranging from 96 to 100% if compared to that of conventional US (from 57 to 96%). Furthermore, it has been demonstrated that SICUS has high accuracy to identify complications and post-operative recurrence (114). An Italian group (115) has compiled a SICUS quantitative sonographic lesion index (SLIC) to study the evolution of the transmural bowel damage in CD

patients in medical therapy. The index includes wall thickness, length of damaged intestinal tract, dilation and strictures. It identifies five classes of severity ranging from the lower (class A) to the higher score (class E). CEUS uses an intravenous contrast (Sonovue, at the recommended dose of 2.4-4.8 ml) and has a good ability to distinguish hypo- or hypervascularized intestinal strictures, identifying inflammation from fibrostenotic lesions (116) (Fig. 6). Active inflammation characteristics include rapid wash-in, when the microbubbles appear  $<20$  second after infusion in bolus, and slow wash-out, when the microbubbles wash out in  $<80$ s (117). Some authors have shown that CEUS can be correlated with CD clinical activity (118). A quantitative method is proposed to evaluate vascularization of the intestinal wall in association with disease activity (the CD activity index, CDAI). Migaletto et al. (119) have demonstrated that CEUS has the highest performance: 93.5% sensitivity, 93.7% specificity and 93.6% accuracy. In post-surgical recurrences, the values are 97%, 91%, 96%, respectively (120). The distinction between inflammatory and fibrotic lesions, as already mentioned, is very important for the therapeutic approach but some studies are controversial about this distinction: increased echogenicity of the submucosal layer results in inflammation while a clear visibility of all intestinal layers suggests fibrosis with reduced blood volume and flow (121, 122). The introduction of







**Figure 4.** Follow-up at 5-years surgical treatment in 23-year-old patient. Images show, near ileum-ileal anastomosis, an inflamed loop, with increased thickness (\*), vascular prominence (arrowhead), fibro-fatty proliferation and progressive transmural hyperenhancement (arrows). Early enhancement is showed in axial plane (a) and transmural enhancement in late venous phase (b)

a dedicated US software, the QLAB (Philips, Koninklijke, Belgium) and the Qontrast (Bracco, Milan, Italy) has allowed a quantitative and semi-quantitative analysis of contrast enhancement of the inflamed area. These data are derived from a selected “region of interest” (ROI) in which median values of the image intensity and a perfusion analysis are calculated to generate a time-intensity curve and obtain a good correlation

between CD inflammatory activity and bowel wall vascularization (123, 124). In a recent study, Quiaia (125) shows a positive correlation between contrast enhancement in affected tracts of small bowel and laboratory data in CD, so that the quantitative CEUS can define responder from non-responder patients affected by active disease and treated with immunosuppressive and biological therapies. The strain elastography (SE)



**Figure 5.** B-mode US with linear probe shows loop affected by CD and with increased mural thickness



**Figure 6.** US after second-generation microbubble contrast shows slight enhancement, without signs of significant inflammation

is another diagnostic method useful but not routinely used for clinical management of inflammatory bowel diseases. It is a technique to evaluate the elasticity of the tissues as demonstrated in a case of ileum Crohn

stricture by Giannetti (126) and confirmed by macroscopic and microscopic examination: the terminal ileum wall appears with abnormal elasticity and higher percentage of “blue” areas, suggesting stiffness.


### Advantages and disadvantages of US

US has many advantages (127). It is widely used, low-cost, repeatable, noninvasive and radiation-free, all reasons that make of this technique one of the most tolerated by the patients. An experienced sonographer can find mural and extramural complications visualizing a good part of the small intestine and colon, in particular as the first approach in urgent assessment of complications. US is accurate in investigating stenosis, abscesses and fistulas; intraluminal oral contrast may improve image quality, but MR can show complicated fistulas better than US. Moreover, the use of intravenous contrast is necessary to distinguish CD strictures with predominantly inflammatory or fibrotic component especially when the stenosis is impassable with the endoscope. CEUS has an important role in revealing the presence and activity of CD in terminal ileum in particular in the follow-up of patients with known ileal localization of disease (128). Unfortunately, US

has some limitations: it is operator-dependent and the comparison or revision of images during the follow-up is not easy. Large body size, intestinal meteorism (in some cases it can be reduced by intraluminal solution) or depth of the region of interest can limit the exam. US cannot explore abdominal regions (retroperitoneum area) and intestinal tracts such as stomach, duodenum, jejunum, transverse colon, deep intrapelvic loops and rectum (129).

### Advantages and disadvantages of MRE

MRE is another radiation-free diagnostic technique and represents the first choice in the pelvic localization of CD (11). It has high diagnostic accuracy and ability to investigate the complete bowel tract; it has multiplanar reconstructions, good visualization of soft tissues and easy detection of complications (strictures, abscess and fistulas) (Fig. 7) thus playing an important



**Figure 7.** Example of enterocolic fistula formations (arrows) in a patient with CD for about 10 years. FIESTA images show «star sign» in axial (a) and coronal(b) planes and transmurial enhancement appears in axial (c) and coronal (d) T1-weighted fat-saturated images



role in surgical planning (127). It provides a panoramic view of the entire abdominal region, mesenteric tissue and retroperitoneum area, allowing a good visualization also in patients in whom US is limited (e.g., overweight patients). Unfortunately, MRE is less used in clinical practice, because it is expensive, needs specific radiological competences and is time consuming (130). Other limitations include presence of metal devices, claustrophobia or sensitivity to MR contrast agents (131, 132).

### CEUS vs MRE

A study by Malagò et al. (133) shows a good correlation between MRE and CEUS activity, with a Spearman's coefficient ( $\rho$ ) = 0.791 and a statistically significant p value ( $<0.0001$ ). In particular, a high correlation was found in the study of the small bowel to evaluate wall thickness, lymph nodes and vasa recta ( $\rho=0.926$ ;  $p<0.0001$ ); accurate correlation was obtained to assess layered wall appearance, disease extension and fibro adipose proliferation ( $\rho=0.716$ ;  $p<0.001$ ). An excellent correlation between imaging and clinical laboratory data was achieved. MRE and CEUS are non-invasive techniques used in the diagnosis of CD, and in monitoring its activity. The affected loop and the main signs of the disease can be determined by both imaging modalities, despite different accuracy rates. In active disease, CEUS offers the possibility of recognizing all the characteristic signs, using intravenous contrast agent (gas microbubbles) that remains inside the microcirculation throughout the procedure, breaks up in the vascular system and is not retained in the fibrous tissue. This technique better shows the bowel-wall enhancement and the increased vascularization of the affected bowel loop (133). Conversely, MRI contrast agent has an initial vascular phase, tends to migrate and to accumulate in the interstitium. Therefore, gadolinium can detect also chronic lesions. Quaiá et al. (134-137) demonstrated the accuracy of both techniques, with a quantitative analysis of enhancement patterns. Time intensity curves in CEUS and MRE are adequate for assessing intestinal-wall vascularity and for a more objective evaluation of the parietal enhancement compared with observer

experience. In chronic lesions (133), which are fibrous scar tissue, MRE shows moderate enhancement in the venous phase with a pattern restricted to the mucosa, layered or homogeneous, while CEUS does not reveal any vascular enhancement.

### Conclusions

An accurate diagnosis of CD is important to plan therapy and follow-up. The diagnosis is accomplished also by means of a bioptic extraction from the mucosal alteration performed during colonoscopy. In clinical practice, MRE is used as a first imaging exam to determine intestinal localization, extension and extra-intestinal complications. It can distinguish between inflamed and fibrotic component and helps in the therapeutic decision-making: medical therapy or surgical treatment. CEUS is useful in the explorable intestinal tracts, already visualized by previous MRI exams, and it is used in the follow-up after medical or surgical treatment or as fist-line exam in patients unsuitable to MR. Besides, CEUS can be helpful in emergency conditions as first approach examination in case of acute complications and recurrence. In conclusion, both techniques are useful and complementary in the study of CD evolution.

**Ethical approval:** This article does not contain any studies with human participants performed by any of the authors.

**Conflict of interest:** None to declare

### References

1. Panes J. Inflammatory bowel disease: pathogenesis and targets for therapeutic interventions. *Acta Physiol Scand* 2001; 173: 159-65.
2. Martin DR, Lauenstein T, Sitaraman SV. Utility of Magnetic Resonance Imaging in Small Bowel Crohn's Disease. *Gastroenterology* 2007; 133: 385-90.
3. Marampon F, Gravina GL, Ju X, et al. Cyclin D1 silencing suppresses tumorigenicity, impairs DNA double strand break repair and thus radiosensitizes androgen-independent prostate cancer cells to DNA damage. *Oncotarget* 2016;
4. Danese S, Fiorino G, Mary JY, et al. Development of red flags index for early referral of adults with symptoms and



- signs suggestive of Crohn's disease: An IOIBD initiative. *J Crohns Colitis* 2015; 9: 601-06.
5. Peyrin-Biroulet L, Reinisch W, Colombel JF, et al. Clinical disease activity, C-reactive protein normalisation and mucosal healing in Crohn's disease in the SONIC trial. *Gut* 2014; 63: 88-95.
  6. Cosnes J, Gower-Rousseau C, Seksik P, Cortot A. Epidemiology and natural history of inflammatory bowel diseases. *Gastroenterology* 2011; 140: 1785-94.
  7. Panes J, Bouhnik Y, Reinisch W, et al. Imaging techniques for assessment of inflammatory bowel disease: Joint ECCO and ESGAR evidence-based consensus guidelines. *J Crohns Colitis* 2013; 7: 556-85.
  8. Porcile C, Di Zazzo E, Monaco ML, et al. Adiponectin as novel regulator of cell proliferation in human glioblastoma. *J Cell Physiol* 2014; 229: 1444-54.
  9. Dalla Palma B, Guasco D, Pedrazzoni M, et al. Osteolytic lesions, cytogenetic features and bone marrow levels of cytokines and chemokines in multiple myeloma patients: Role of chemokine (C-C motif) ligand 20. *Leukemia* 2016; 30: 409-16.
  10. Deepak P, Park SH, Ehman EC, et al. Crohn's disease diagnosis, treatment approach, and management paradigm: what the radiologist needs to know. *Abdom Radiol (NY)* 2017; 42: 1068-86.
  11. Hommes DW, Van Deventer SJH. Endoscopy in inflammatory bowel diseases. *Gastroenterology* 2004; 126: 1561-73.
  12. Iacobellis F, Berritto D, Fleischmann D, et al. CT findings in acute, subacute, and chronic ischemic colitis: suggestions for diagnosis. *Biomed Res Int* 2014; 2014: 895248.
  13. Mazzei MA, Guerrini S, Squitieri NC, et al. The role of US examination in the management of acute abdomen. *Crit Ultrasound J* 2013; 5: 1-9.
  14. Reginelli A, Iacobellis F, Del Vecchio L, et al. VFMSS findings in elderly dysphagic patients: Our experience. *BMC Surg* 2013; 13:
  15. Somma F, Berritto D, Iacobellis F, et al. 7T  $\mu$ MRI of mesenteric venous ischemia in a rat model: Timing of the appearance of findings. *Magn Reson Imaging* 2013; 31: 408-13.
  16. Berritto D, Somma F, Landi N, et al. Seven-Tesla micro-MRI in early detection of acute arterial ischaemia: evolution of findings in an in vivo rat model. *Radiol Med* 2011; 116: 829-41.
  17. Grassi R, Lombardi G, Reginelli A, et al. Coccygeal movement: Assessment with dynamic MRI. *Eur J Radiol* 2007; 61: 473-79.
  18. Iacobellis F, Berritto D, Somma F, et al. Magnetic resonance imaging: A new tool for diagnosis of acute ischemic colitis? *World J Gastroenterol* 2012; 18: 1496-501.
  19. Mazzei MA, Mazzei FG, Marrelli D, et al. Computed tomographic evaluation of mesentery: Diagnostic value in acute mesenteric ischemia. *J Comput Assist Tomogr* 2012; 36: 1-7.
  20. Cappabianca S, Granata V, Di Grezia G, et al. The role of nasoenteric intubation in the MR study of patients with Crohn's disease: Our experience and literature review. *Radiol Med* 2011; 116: 389-406.
  21. Grassi R, Romano S, Micera O, Fioroni C, Boller B. Radiographic findings of post-operative double stapled trans anal rectal resection (STARR) in patient with obstructed defecation syndrome (ODS). *Eur J Radiol* 2005; 53: 410-16.
  22. Grassi R, Di Mizio R, Pinto A, Romano L, Rotondo A. Serial plain abdominal film findings in the assessment of acute abdomen: spastic ileus, hypotonic ileus, mechanical ileus and paralytic ileus. *Radiol Med* 2004; 108: 56-70.
  23. Buffa V, Solazzo A, D'Auria V, et al. Dual-source dual-energy CT: dose reduction after endovascular abdominal aortic aneurysm repair. *Radiol Med* 2014; 119: 934-41.
  24. Grassi R, Pinto A, Mannelli L, Marin D, Mazzei MA, New Imaging in Gastrointestinal Tract, *Gastroenterol Res Pract*, 2016.
  25. Di Cesare E, Gennarelli A, Di Sibio A, et al. Assessment of dose exposure and image quality in coronary angiography performed by 640-slice CT: a comparison between adaptive iterative and filtered back-projection algorithm by propensity analysis. *Radiol Med* 2014; 119: 642-49.
  26. Mandato Y, Reginelli A, Galasso R, Iacobellis F, Berritto D, Cappabianca S. Errors in the Radiological Evaluation of the Alimentary Tract: Part I. *Semin Ultrasound CT MR* 2012; 33: 300-07.
  27. Reginelli A, Russo A, Maresca D, Martiniello C, Cappabianca S, Brunese L. Imaging Assessment of Gunshot Wounds. *Semin Ultrasound CT MR* 2015; 36: 57-67.
  28. Pinto A, Pinto F, Faggian A, et al. Sources of error in emergency ultrasonography. *Crit Ultrasound J* 2013; 5 Suppl 1: S1-S1.
  29. Cappabianca S, Porto A, Petrillo M, et al. Preliminary study on the correlation between grading and histology of solitary pulmonary nodules and contrast enhancement and [18F] fluorodeoxyglucose standardised uptake value after evaluation by dynamic multiphase CT and PET/CT. *J Clin Pathol* 2011; 64: 114-19.
  30. Miele V, Piccolo CL, Trinci M, Galluzzo M, Ianniello S, Brunese L. Diagnostic imaging of blunt abdominal trauma in pediatric patients. *Radiol Med* 2016; 121: 409-30.
  31. Tamburrini S, Solazzo A, Sagnelli A, et al. Amyotrophic lateral sclerosis: Sonographic evaluation of dysphagia. *Radiol Med* 2010; 115: 784-93.
  32. Pinto A, Reginelli A, Cagini L, et al. Accuracy of ultrasonography in the diagnosis of acute calculous cholecystitis: review of the literature. *Crit Ultrasound J* 2013; 5 Suppl 1: S11.
  33. Pallotta N, Vincoli G, Montesani C, et al. Small intestine contrast ultrasonography (SICUS) for the detection of small bowel complications in crohn's disease: A prospective comparative study versus intraoperative findings. *Inflamm Bowel Dis* 2012; 18: 74-84.
  34. De Cecco CN, Buffa V, Fedeli S, et al. Preliminary experience with abdominal dual-energy CT (DECT): True versus virtual nonenhanced images of the liver. *Radiol Med* 2010; 115: 1258-66.

35. di Giacomo V, Trinci M, van der Byl G, Catania VD, Calisti A, Miele V. Ultrasound in newborns and children suffering from non-traumatic acute abdominal pain: imaging with clinical and surgical correlation. *J Ultrasound* 2015; 18: 385-93.
36. Pinto A, Miele V, Laura Schillirò M, et al. Spectrum of Signs of Pneumoperitoneum. *Semin Ultrasound CT MR* 2016; 37: 3-9.
37. Sessa B, Galluzzo M, Ianniello S, Pinto A, Trinci M, Miele V. Acute Perforated Diverticulitis: Assessment With Multi-detector Computed Tomography. *Semin Ultrasound CT MR* 2016; 37: 37-48.
38. Valentini V, Buquicchio GL, Galluzzo M, et al. Intussusception in Adults: The Role of MDCT in the Identification of the Site and Cause of Obstruction. *Gastroenterol Res Pract* 2016; 2016: 5623718-18.
39. Faggian A, Fracella MR, D'Alesio G, et al. Small-Bowel Neoplasms: Role of MRI Enteroclysis. *Gastroenterol Res Pract* 2016; 2016:
40. Valentini G, Marcoccia A, Cuomo G, et al. Early systemic sclerosis: marker autoantibodies and videocapillaroscopy patterns are each associated with distinct clinical, functional and cellular activation markers. *Arthritis Res Ther* 2013; 15: R63.
41. Reginelli A, Mandato Y, Solazzo A, Berritto D, Iacobellis F, Grassi R. Errors in the Radiological Evaluation of the Alimentary Tract: Part II. *Semin Ultrasound CT MR* 2012; 33: 308-17.
42. Dionigi G, Dionigi R, Rovera F, et al. Treatment of high output entero-cutaneous fistulae associated with large abdominal wall defects: single center experience. *Int J Surg* 2008; 6: 51-6.
43. Mocchegiani F, Vincenzi P, Coletta M, et al. Prevalence and clinical outcome of hepatic haemangioma with specific reference to the risk of rupture: A large retrospective cross-sectional study. *Dig Liver Dis* 2016; 48: 309-14.
44. Salvolini L, Urbinati C, Valeri G, Ferrara C, Giovagnoni A. Contrast-enhanced MR cholangiography (MRCP) with GD-EOB-DTPA in evaluating biliary complications after surgery. *Radiol Med* 2012; 117: 354-68.
45. Cappabianca S, Reginelli A, Monaco L, Del Vecchio L, Di Martino N, Grassi R. Combined videofluoroscopy and manometry in the diagnosis of oropharyngeal dysphagia: Examination technique and preliminary experience. *Radiol Med* 2008; 113: 923-40.
46. Maggioletti N, Capasso R, Pinto D, et al. Diagnostic value of computed tomography colonography (CTC) after incomplete optical colonoscopy. *Int J Surg* 2016; 33 Suppl 1: S36-44.
47. Sforza V, Martinelli E, Ciardiello F, et al. Mechanisms of resistance to anti-epidermal growth factor receptor inhibitors in metastatic colorectal cancer. *World J Gastroenterol* 2016; 22: 6345-61.
48. Valeri G, Mazza FA, Maggi S, et al. Open source software in a practical approach for post processing of radiologic images. *Radiol Med* 2015; 120: 309-23.
49. Grassi R, Rambaldi PF, Di Grezia G, et al. Inflammatory bowel disease: Value in diagnosis and management of MDCT-enteroclysis and 99mTc-HMPAO labeled leukocyte scintigraphy. *Abdom Imaging* 2011; 36: 372-81.
50. Lo Re G, Cappello M, Tudisca C, et al. CT enterography as a powerful tool for the evaluation of inflammatory activity in Crohn's disease: Relationship of CT findings with CDAI and acute-phase reactants. *Radiol Med* 2014; 119: 658-66.
51. Taylor SA, Avni F, Cronin CG, et al. The first joint ESGAR/ ESPR consensus statement on the technical performance of cross-sectional small bowel and colonic imaging. *Eur Radiol* 2017; 27: 2570-82.
52. Puylaert CAJ, Tielbeek JAW, Bipat S, Stoker J. Grading of Crohn's disease activity using CT, MRI, US and scintigraphy: a meta-analysis. *Eur Radiol* 2015; 25: 3295-313.
53. Di Cesare E, Cademartiri F, Carbone I, et al. Clinical indications for the use of cardiac MRI. by the SIRM Study Group on Cardiac Imaging. *Radiol Med* 2013; 118: 752-98.
54. Schicchi N, Valeri G, Moroncini G, et al. Myocardial perfusion defects in scleroderma detected by contrast-enhanced cardiovascular magnetic resonance. *Radiol Med* 2014; 119: 885-94.
55. Barile A, Arrigoni F, Bruno F, et al. Computed Tomography and MR Imaging in Rheumatoid Arthritis. *Radiol Clin North Am* 2017; 55: 997-1007.
56. Maurizi N, Passantino S, Spaziani G, et al. Long-term Outcomes of Pediatric-Onset Hypertrophic Cardiomyopathy and Age-Specific Risk Factors for Lethal Arrhythmic Events. *JAMA Cardiol* 2018; 3: 520-25.
57. Pradella S, Lucarini S, Colagrande S. Liver lesion characterization: The wrong choice of contrast agent can mislead the diagnosis of hemangioma. *Am J Roentgenol* 2012; 199:
58. Macchi M, Belfiore MP, Floridi C, et al. Radiofrequency versus microwave ablation for treatment of the lung tumours: LUMIRA (lung microwave radiofrequency) randomized trial. *Med Oncol* 2017; 34:
59. Tarantini G, Favaretto E, Napodano M, et al. Design and methodologies of the postconditioning during coronary angioplasty in acute myocardial infarction (POST-AMI) trial. *Cardiology* 2010; 116: 110-16.
60. Cazzato RL, Arrigoni F, Emanuele Boatta, et al. Percutaneous management of bone metastases: state of the art, interventional strategies and joint position statement of the Italian College of MSK Radiology (ICoMSKR) and the Italian College of Interventional Radiology (ICIR). *Radiol Med* 2018; 1: 3-3.
61. Barile A, Arrigoni F, Bruno F, et al. Present role and future perspectives of interventional radiology in the treatment of painful bone lesions. *Future Oncol* 2018; 14: 2945-55.
62. Bruno F, Barile A, Arrigoni F, et al. Weight-bearing MRI of the knee: A review of advantages and limits. *Acta Biomed* 2018; 89: 78-88.
63. Arrigoni F, Bruno F, Zugaro L, et al. Developments in the management of bone metastases with interventional radiology. *Acta Biomed* 2018; 89: 166-74.
64. Zoccali C, Arrigoni F, Mariani S, Bruno F, Barile A, Mas-

- ciocchi C. An unusual localization of chondroblastoma: The triradiate cartilage; from a case report a reconstructive technique proposal with imaging evolution. *J Clin Orthop Trauma* 2017; 8: S48-S52.
65. Bruno F, Smaldone F, Varrassi M, et al. MRI findings in lumbar spine following O2-O3 chemiodisclosyis: A long-term follow-up. *Interv Neuroradiol* 2017; 23: 444-50.
66. Barile A, Bruno F, Arrigoni F, et al. Emergency and Trauma of the Ankle. *Semi Musc Rad* 2017; 21: 282-89.
67. Splendiani A, Bruno F, Patriarca L, et al. Thoracic spine trauma: advanced imaging modality. *Radiol Med* 2016; 121: 780-92.
68. Giordano AV, Arrigoni F, Bruno F, et al. Interventional Radiology Management of a Ruptured Lumbar Artery Pseudoaneurysm after Cryoablation and Vertebroplasty of a Lumbar Metastasis. *Cardiovasc Intervent Radiol* 2017; 40: 776-79.
69. Barile A, Bruno F, Mariani S, et al. What can be seen after rotator cuff repair: a brief review of diagnostic imaging findings. *Musculoskelet Surg* 2017; 101: 3-14.
70. Barile A, Bruno F, Mariani S, et al. Follow-up of surgical and minimally invasive treatment of Achilles tendon pathology: a brief diagnostic imaging review. *Musculoskelet Surg* 2017; 101: 51-61.
71. Splendiani A, Perri M, Grattacaso G, et al. Magnetic resonance imaging (MRI) of the lumbar spine with dedicated G-scan machine in the upright position: a retrospective study and our experience in 10 years with 4305 patients. *Radiol Med* 2016; 121: 38-44.
72. Arrigoni F, Barile A, Zugaro L, et al. Intra-articular benign bone lesions treated with Magnetic Resonance-guided Focused Ultrasound (MRgFUS): imaging follow-up and clinical results. *Med Oncol* 2017; 34: 55.
73. Zappia M, Castagna A, Barile A, Chianca V, Brunese L, Pouliart N. Imaging of the coracoglenoid ligament: a third ligament in the rotator interval of the shoulder. *Skeletal Radiol* 2017; 46: 1101-11.
74. Barile A, La Marra A, Arrigoni F, et al. Anaesthetics, steroids and platelet-rich plasma (PRP) in ultrasound-guided musculoskeletal procedures. *Br J Radiol* 2016; 89: 20150355.
75. Reginelli A, Zappia M, Barile A, Brunese L. Strategies of imaging after orthopedic surgery. *Musculoskelet Surg* 2017; 101: 1.
76. Maccioni F, Al Ansari N, Mazzamurro F, et al. Detection of Crohn disease lesions of the small and large bowel in pediatric patients: Diagnostic value of MR enterography versus reference examinations. *Am J Roentgenol* 2014; 203: W533-W42.
77. Micheli G, Corridore A, Torlone S, et al. Dynamic MRI in the evaluation of the spine: State of the art. *Acta Biomed* 2018; 89: 89-101.
78. Masciocchi C, Arrigoni F, Ferrari F, et al. Uterine fibroid therapy using interventional radiology mini-invasive treatments: current perspective. *Med Oncol* 2017; 34: 52.
79. Scialpi M, Cappabianca S, Rotondo A, et al. Pulmonary congenital cystic disease in adults. Spiral computed tomography findings with pathologic correlation and management. *Radiol Med* 2010; 115: 539-50.
80. Muccio CF, Di Blasi A, Esposito G, Brunese L, D'Arco F, Caranci F. Perfusion and spectroscopy magnetic resonance imaging in a case of lymphocytic vasculitis mimicking brain tumor. *Pol J Radiol* 2013; 78: 66-69.
81. Battipaglia G, Avilia S, Morelli E, Caranci F, Perna F, Camera A. Posterior reversible encephalopathy syndrome (PRES) during induction chemotherapy for acute myeloblastic leukemia (AML). *Ann Hematol* 2012; 91: 1327-28.
82. Briganti F, Leone G, Marseglia M, Cicala D, Caranci F, Maiuri F. P64 Flow Modulation Device in the treatment of intracranial aneurysms: Initial experience and technical aspects. *J Neurointerv Surg* 2016; 8: 173-80.
83. Cirillo M, Caranci F, Tortora F, et al. Structural neuroimaging in dementia. *J Alzheimers Dis* 2012; 29: 16-19.
84. Sverzellati N, Calabrò E, Chetta A, et al. Visual score and quantitative CT indices in pulmonary fibrosis: Relationship with physiologic impairment. *Radiol Med* 2007; 112: 1160-72.
85. Bertolini L, Vaglio A, Bignardi L, et al. Subclinical interstitial lung abnormalities in stable renal allograft recipients in the era of modern immunosuppression. *Transplant Proc* 2011; 43: 2617-23.
86. Di Cesare E, Gennarelli A, Di Sibio A, et al. Image quality and radiation dose of single heartbeat 640-slice coronary CT angiography: A comparison between patients with chronic Atrial Fibrillation and subjects in normal sinus rhythm by propensity analysis. *Eur J Radiol* 2015; 84: 631-36.
87. Masciocchi C, Conti L, D'Orazio F, Conchiglia A, Lanni G, Barile A, Errors in Musculoskeletal MRI, in: Romano L., Pinto A. (Eds.), *Errors in Radiology*, Springer Milan, Milano, 2012, pp. 209-17.
88. Lenze F, Wessling J, Bremer J, et al. Detection and differentiation of inflammatory versus fibromatous Crohn's disease strictures: Prospective comparison of 18F-FDG-PET/CT, MR-enteroclysis, and transabdominal ultrasound versus endoscopic/histologic evaluation. *Inflamm Bowel Dis* 2012; 18: 2252-60.
89. Bettenworth D, Nowacki TM, Cordes F, Buerke B, Lenze F. Assessment of stricturing Crohn's disease: Current clinical practice and future avenues. *World J Gastroenterol* 2016; 22: 1008-16.
90. Westerland O, Griffin N. Magnetic Resonance Enterography in Crohn's Disease. *Semin Ultrasound CT MR* 2016; 37: 282-91.
91. Tolan DJM, Greenhalgh R, Zealley IA, Halligan S, Taylor SA. MR Enterographic Manifestations of Small Bowel Crohn Disease. *RadioGraphics* 2010; 30: 367-84.
92. Grand DJ, Guglielmo FF, Al-Hawary MM. MR enterography in Crohn's disease: current consensus on optimal imaging technique and future advances from the SAR Crohn's disease-focused panel. *Abdom Imaging* 2015; 40: 953-64.
93. Iannicelli E, Martini I, Fantini C, et al. Magnetic resonance enterography in Crohn's disease: New simple proposal to assess disease activity. *Clin Imaging* 2016; 40: 492-97.



94. Dohan A, Taylor S, Hoeffel C, et al. Diffusion-weighted MRI in Crohn's disease: Current status and recommendations. *Journal of magnetic resonance imaging : JMRI* 2016; 44: 1381-96.
95. Rimola J, Rodriguez S, García-Bosch O, et al. Magnetic resonance for assessment of disease activity and severity in ileocolonic Crohn's disease. *Gut* 2009; 58: 1113-20.
96. Rimola J, Ordás I, Rodriguez S, et al. Magnetic resonance imaging for evaluation of Crohn's disease: Validation of parameters of severity and quantitative index of activity. *Inflamm Bowel Dis* 2011; 17: 1759-68.
97. Kim JS, Jang HY, Park SH, et al. MR enterography assessment of bowel inflammation severity in Crohn disease using the MR index of activity score: Modifying roles of DWI and effects of contrast phases. *Am J Roentgenol* 2017; 208: 1022-29.
98. Arrigoni F, Bruno F, Zugaro L, et al. Role of interventional radiology in the management of musculoskeletal soft-tissue lesions. *Radiol Med* 2018; 1-6.
99. Barile A, Quarchioni S, Bruno F, et al. Interventional radiology of the thyroid gland: Critical review and state of the art. *Gland Surg* 2018; 7: 132-46.
100. Brunese L, Romeo A, Iorio S, et al. Thyroid B-flow twinkling sign: a new feature of papillary cancer. *Eur J Endocrinol* 2008; 159: 447-51.
101. Cantisani V, Grazhdani H, Drakonaki E, et al. Strain US elastography for the characterization of thyroid nodules: Advantages and limitation. *Int J Endocrinol* 2015; 2015: 1-6.
102. Gatta G, Parlato V, Di Grezia G, et al. Ultrasound-guided aspiration and ethanol sclerotherapy for treating endometrial cysts. *Radiol Med* 2010; 115: 1330-39.
103. Regine G, Atzori M, Miele V, et al. Second-generation sonographic contrast agents in the evaluation of renal trauma. *Radiol Med* 2007; 112: 581-87.
104. Iacobellis F, Segreto T, Berritto D, et al. A rat model of acute kidney injury through systemic hypoperfusion evaluated by micro-US, color and PW-Doppler. *Radiol Med* 2018; 1-6.
105. Perrotta FM, Astorri D, Zappia M, Reginelli A, Brunese L, Lubrano E. An ultrasonographic study of enthesitis in early psoriatic arthritis patients naive to traditional and biologic DMARDs treatment. *Rheumatol Int* 2016; 36: 1579-83.
106. Wale A, Pilcher J. Current Role of Ultrasound in Small Bowel Imaging. *Semin Ultrasound CT MR* 2016; 37: 301-12.
107. De Franco A, Di Veronica A, Armuzzi A, et al. Ileal Crohn disease: mural microvascularity quantified with contrast-enhanced US correlates with disease activity. *Radiology* 2012; 262: 680-8.
108. Allen BC, Baker ME, Einstein DM, et al. Effect of altering automatic exposure control settings and quality reference mAs on radiation dose, image quality, and diagnostic efficacy in MDCT enterography of active inflammatory Crohn's disease. *AJR Am J Roentgenol* 2010; 195: 89-100.
109. Taylor SA, Punwani S, Rodriguez-Justo M, et al. Mural Crohn disease: correlation of dynamic contrast-enhanced MR imaging findings with angiogenesis and inflammation at histologic examination--pilot study. *Radiology* 2009; 251: 369-79.
110. Esteban JM, Maldonado L, Sanchiz V, Minguez M, Benages A. Activity of Crohn's disease assessed by colour Doppler ultrasound analysis of the affected loops. *Eur Radiol* 2001; 11: 1423-28.
111. Manetta R, Pistoia ML, Bultrini C, Stavroulis E, Di Cesare E, Masciocchi C. Ultrasound enhanced with sulphur-hexafluoride-filled microbubbles agent (SonoVue) in the follow-up of mild liver and spleen trauma. *Radiol Med* 2009; 114: 771-9.
112. Nylund K, Hausken T, Gilja OH. Ultrasound and inflammatory bowel disease. *Ultrasound Q* 2010; 26: 3-15.
113. Mocci G, Migaleddu V, Cabras F, et al. SICUS and CEUS imaging in Crohn's disease: an update. *J Ultrasound* 2017; 20: 1-9.
114. Onali S, Calabrese E, Petruzzello C, et al. Endoscopic vs ultrasonographic findings related to Crohn's disease recurrence: a prospective longitudinal study at 3 years. *J Crohns Colitis* 2010; 4: 319-28.
115. Calabrese E, Zorzi F, Zuzzi S, et al. Development of a numerical index quantitating small bowel damage as detected by ultrasonography in Crohn's disease. *J Crohns Colitis* 2012; 6: 852-60.
116. Kratzer W, von Tirpitz C, Mason R, et al. Contrast-enhanced power Doppler sonography of the intestinal wall in the differentiation of hypervascularized and hypovascularized intestinal obstructions in patients with Crohn's disease. *J Ultrasound Med* 2002; 21: 149-57.
117. Romanini L, Passamonti M, Navarria M, et al. Quantitative analysis of contrast-enhanced ultrasonography of the bowel wall can predict disease activity in inflammatory bowel disease. *Eur J Radiol* 2014; 83: 1317-23.
118. Robotti D, Cammarota T, Debani P, Sarno A, Astegiano M. Activity of Crohn disease: value of Color-Power-Doppler and contrast-enhanced ultrasonography. *Abdom Imaging* 2004; 29: 648-52.
119. Migaleddu V, Scanu AM, Quaia E, et al. Contrast-enhanced ultrasonographic evaluation of inflammatory activity in Crohn's disease. *Gastroenterology* 2009; 137: 43-52.
120. Paredes JM, Ripolles T, Cortes X, et al. Contrast-enhanced ultrasonography: usefulness in the assessment of postoperative recurrence of Crohn's disease. *J Crohns Colitis* 2013; 7: 192-201.
121. Parente F, Maconi G, Bollani S, et al. Bowel ultrasound in assessment of Crohn's disease and detection of related small bowel strictures: a prospective comparative study versus x ray and intraoperative findings. *Gut* 2002; 50: 490-5.
122. Nylund K, Jirik R, Mezl M, et al. Quantitative contrast-enhanced ultrasound comparison between inflammatory and fibrotic lesions in patients with Crohn's disease. *Ultrasound Med Biol* 2013; 39: 1197-206.
123. Girlich C, Jung EM, Iesalnieks I, et al. Quantitative as-

- assessment of bowel wall vascularisation in Crohn's disease with contrast-enhanced ultrasound and perfusion analysis. *Clin Hemorheol Microcirc* 2009; 43: 141-8.
124. Girlich C, Jung EM, Huber E, et al. Comparison between preoperative quantitative assessment of bowel wall vascularization by contrast-enhanced ultrasound and operative macroscopic findings and results of histopathological scoring in Crohn's disease. *Ultraschall Med* 2011; 32: 154-9.
125. Quaia E, Sozzi M, Angileri R, Gennari AG, Cova MA. Time-Intensity Curves Obtained after Microbubble Injection Can Be Used to Differentiate Responders from Nonresponders among Patients with Clinically Active Crohn Disease after 6 Weeks of Pharmacologic Treatment. *Radiology* 2016; 281: 606-16.
126. Giannetti A, Biscontri M, Matergi M, Stumpo M, Minacci C. Feasibility of CEUS and strain elastography in one case of ileum Crohn stricture and literature review. *J Ultrasound* 2016; 19: 231-7.
127. Spinelli A, Allocca M, Jovani M, Danese S. Review article: optimal preparation for surgery in Crohn's disease. *Aliment Pharmacol Ther* 2014; 40: 1009-22.
128. Horjus Talabur Horje CS, Bruijnen R, Roovers L, Groenen MJ, Joosten FB, Wahab PJ. Contrast Enhanced Abdominal Ultrasound in the Assessment of Ileal Inflammation in Crohn's Disease: A Comparison with MR Enterography. *PLoS One* 2015; 10: e0136105.
129. Wilson SR, Burns PN. Microbubble-enhanced US in body imaging: what role? *Radiology* 2010; 257: 24-39.
130. Spinelli A, Allocca M, Jovani M, Danese S, Review article: Optimal preparation for surgery in Crohn's disease, 2014, pp. 1009-22.
131. Splendiani A, Perri M, Marsecano C, et al. Effects of serial macrocyclic-based contrast materials gadoterate meglumine and gadobutrol administrations on gadolinium-related dentate nuclei signal increases in unenhanced T1-weighted brain: a retrospective study in 158 multiple sclerosis (MS) patients. *Radiol Med* 2018; 123: 125-34.
132. Tedeschi E, Caranci F, Giordano F, Angelini V, Cocozza S, Brunetti A. Gadolinium retention in the body: what we know and what we can do. *Radiol Med* 2017; 122: 589-600.
133. Malago R, D'Onofrio M, Mantovani W, et al. Contrast-enhanced ultrasonography (CEUS) vs. MRI of the small bowel in the evaluation of Crohn's disease activity. *Radiol Med* 2012; 117: 268-81.
134. Quaia E, Migaleddu V, Baratella E, et al. The diagnostic value of small bowel wall vascularity after sulfur hexafluoride-filled microbubble injection in patients with Crohn's disease. Correlation with the therapeutic effectiveness of specific anti-inflammatory treatment. *Eur J Radiol* 2009; 69: 438-44.
135. Bozzetti C, Nizzoli R, Tiseo M, et al. ALK and ROS1 rearrangements tested by fluorescence in situ hybridization in cytological smears from advanced non-small cell lung cancer patients. *Diagnostic Cytopathology*, vol. 43, p. 941-946, ISSN: 8755-1039, doi: 10.1002/dc.23318.
136. De Filippo M, Gira F, Corradi D, Sverzellati N, Zompatori M, Rossi C. (2011). Benefits of 3D technique in guiding percutaneous retroperitoneal biopsies. *RAD. MED*, vol. 116(3), p. 407-416, ISSN: 0033-8362, doi: 10.1007/s11547-010-0604-2
137. De Filippo M, Onniboni M, Rusca M, et al. (2008). Advantages of multidetector row CT with multiplanar reformation in guiding percutaneous lung biopsies. *RAD. MED*, vol. 113, p. 945-953, ISSN: 0033-8362, doi: 10.1007/s11547-008-0325-y

Received: 26 March 2019

Accepted: 4 April 2019

Correspondence:

Antonio Barile M.D.

Department of Biotechnology and Applied Clinical Sciences  
University of L'Aquila, Vetoio Street, 1 - 67100 L'Aquila, Italy  
Tel. +390862414258

E-mail: antonio.barile@cc.univaq.it