

Supplemental Table 1. Standardized Mortality Ratios for All Causes by calendar year. Observed (O) and Expected (E) Cases, SMR and 95% Confidence Interval (95% CI) Follow Up Period 1989 – 2011*

Year	O	FVG Standard Rates ¹			Italian Standard Rates ²		
		E	SMR	95% CI	E	SMR	95% CI
1989	-	0.97	-	-	0.84	-	-
1990	-	1.17	-	-	1.02	-	-
1991	-	1.40	-	-	1.21	-	-
1992	-	1.74	-	-	1.52	-	-
1993	-	1.98	-	-	1.78	-	-
1994	-	2.66	-	-	2.36	-	-
1995	2	3.75	0.53	0.06-1.93	3.37	0.59	0.07-2.14
1996	3	4.86	0.62	0.13-1.80	4.39	0.68	0.14-2.00
1997	6	5.93	1.01	0.33-2.21	5.52	1.09	0.36-2.37
1998	4	7.57	0.53	0.14-1.35	7.01	0.57	0.16-1.46
1999	5	9.13	0.55	0.18-1.28	8.40	0.60	0.19-1.39
2000	9	11.09	0.81	0.37-1.54	10.36	0.87	0.40-1.65
2001	16	14.99	1.07	0.61-1.73	13.97	1.14	0.65-1.85
2002	23	19.48	1.18	0.75-1.77	18.25	1.26	0.80-1.89
2003	20	26.53	0.75	0.46-1.16	24.51	0.82	0.50-1.26
2004	27	33.00	0.82	0.54-1.19	30.48	0.89	0.58-1.29
2005	30	32.56	0.92	0.62-1.32	30.92	0.97	0.65-1.39
2006	31	36.73	0.84	0.57-1.21	34.95	0.89	0.60-1.27
2007	44	38.65	1.14	0.83-1.53	37.96	1.16	0.85-1.55
2008	42	43.44	0.97	0.69-1.31	41.30	1.02	0.73-1.38
2009	41	45.20	0.91	0.65-1.23	43.54	0.94	0.67-1.28
2010	41	46.50	0.88	0.63-1.20	44.36	0.92	0.66-1.26
2011	44	47.97	0.92	0.67-1.23	46.36	0.95	0.69-1.27

*Standard rate: age-specific mortality rates of FVG for years 1989-2003 and 2006-2011, provided by the Istituto Superiore di Sanità (ISS); ²Standard rate: age-specific mortality rates of Italy for years 1989-2003 and 2006-2011, provided by the Istituto Superiore di Sanità (ISS). *Follow Up Period: 1989 – 2011. Cohort included 2,488 subjects, for 26,415.77 total person-years.

Supplemental Table 2. Standardized Mortality Ratios for All Cancers by calendar year. Observed (O) and Expected (E) Cases, SMR and 95% Confidence Interval (95% CI) Follow Up Period 1989 – 2011*

Year	O	FVG Standard Rates ¹			Italian Standard Rates ²		
		E	SMR	95% CI	E	SMR	95% CI
1989	-	0.39	-	-	0.32	-	-
1990	-	0.49	-	-	0.39	-	-
1991	-	0.57	-	-	0.46	-	-
1992	-	0.74	-	-	0.59	-	-
1993	-	0.86	-	-	0.70	-	-
1994	-	1.17	-	-	0.96	-	-
1995	0	1.58	0.00	-	1.34	0.00	-
1996	2	2.08	0.96	0.12-3.47	1.74	1.15	0.14-4.15
1997	1	2.45	0.41	0.01-2.27	2.20	0.46	0.01-2.53
1998	2	3.22	0.62	0.08-2.24	2.77	0.72	0.09-2.61
1999	2	3.93	0.51	0.06-1.84	3.33	0.60	0.07-2.17
2000	5	4.67	1.07	0.35-2.49	4.13	1.21	0.39-2.82
2001	10	6.52	1.53	0.74-2.82	5.69	1.76	0.84-3.23
2002	9	8.75	1.03	0.47-1.96	7.51	1.20	0.55-2.28
2003	12	11.27	1.06	0.55-1.86	10.12	1.19	0.61-2.08
2004	14	13.95	1.00	0.55-1.69	12.53	1.12	0.61-1.88
2005	11	14.69	0.75	0.37-1.34	13.27	0.83	0.41-1.48
2006	15	16.45	0.91	0.51-1.50	14.85	1.01	0.57-1.67
2007	22	17.08	1.29	0.81-1.95	15.95	1.38	0.86-2.08
2008	20	18.99	1.05	0.64-1.62	16.92	1.18	0.72-1.82
2009	18	18.93	0.95	0.56-1.50	17.53	1.03	0.61-1.62
2010	20	19.89	1.01	0.61-1.55	17.83	1.12	0.69-1.73
2011	18	20.31	0.89	0.53-1.40	18.21	0.99	0.59-1.56

¹Standard rate: age-specific mortality rates of FVG for years 1989-2003 and 2006-2011, provided by the Istituto Superiore di Sanità (ISS); ²Standard rate: age-specific mortality rates of Italy for years 1989-2003 and 2006-2011, provided by the Istituto Superiore di Sanità (ISS). *Follow Up Period: 1989 – 2011. Cohort included 2,488 subjects, for 26,415.77 total person-years.

Supplemental Table 3. Standardized Mortality Ratios for Malignant Pleural cancer by calendar year. Observed (O) and Expected (E) Cases, SMR and 95% Confidence Interval (95% CI) Follow Up Period 1989 – 2011*

Year	O	FVG Standard Rates ¹			Italian Standard Rates ²		
		E	SMR	95% CI	E	SMR	95% CI
1989	-	0.00	-	-	0.00	-	-
1990	-	0.01	-	-	0.00	-	-
1991	-	0.00	-	-	0.00	-	-
1992	-	0.01	-	-	0.00	-	-
1993	-	0.01	-	-	0.01	-	-
1994	-	0.01	-	-	0.01	-	-
1995	-	0.02	-	-	0.01	-	-
1996	1	0.04	22.32	0.56-124.33	0.01	78.41	1.98-436.74
1997	-	0.03	-	-	0.02	-	-
1998	-	0.04	-	-	0.02	-	-
1999	1	0.06	17.73	0.45-98.77	0.03	32.01	0.81-178.30
2000	-	0.09	-	-	0.04	-	-
2001	2	0.14	13.81	1.67-49.86	0.05	37.27	4.51-134.55
2002	3	0.21	14.33	2.95-41.83	0.08	39.57	8.15-115.55
2003	-	0.14	-	-	0.10	-	-
2004	2	0.17	11.92	1.44-43.03	0.12	17.12	2.07-61.79
2005	1	0.32	3.13	0.08-17.45	0.15	6.77	0.17-37.68
2006	2	0.35	5.69	0.69-20.55	0.16	12.29	1.49-44.36
2007	3	0.31	9.73	2.00-28.41	0.18	16.51	3.40-48.21
2008	4	0.53	7.48	2.04-19.16	0.19	21.33	5.80-54.62
2009	3	0.43	7.01	1.44-20.48	0.21	14.62	3.01-42.69
2010	2	0.49	4.08	0.49-14.71	0.20	9.90	1.20-35.75
2011	1	0.22	4.59	0.12-25.59	0.20	4.93	0.12-27.47

¹Standard rate: age-specific mortality rates of FVG for years 1989-2003 and 2006-2011 calculated from the data of mortality provided by the National Institute of Statistics (ISTAT); ²Standard rate: age-specific mortality rates of Italy for years 1989-2003 and 2006-2011 calculated from the data of mortality provided by the National Institute of Statistics (ISTAT); *Follow Up Period: 1989 – 2011. Cohort included 2,488 subjects, for 26,415.77 total person-years.

Supplemental Table 4. Standardized Mortality Ratios for Lung cancer by calendar year. Observed (O) and Expected (E) Cases, SMR and 95% Confidence Interval (95% CI) Follow Up Period 1989 – 2011*

Year	O	FVG Standard Rates ¹			Italian Standard Rates ²		
		E	SMR	95% CI	E	SMR	95% CI
1989	-	0.12	-	-	0.11	-	-
1990	-	0.14	-	-	0.13	-	-
1991	-	0.16	-	-	0.15	-	-
1992	-	0.22	-	-	0.19	-	-
1993	-	0.26	-	-	0.23	-	-
1994	-	0.33	-	-	0.31	-	-
1995	-	0.44	-	-	0.43	-	-
1996	-	0.58	-	-	0.55	-	-
1997	1	0.73	1.37	0.03-7.62	0.69	1.45	0.04-8.06
1998	-	0.89	-	-	0.85	-	-
1999	-	1.06	-	-	1.03	-	-
2000	2	1.20	1.67	0.20-6.03	1.26	1.59	0.19-5.75
2001	2	1.74	1.15	0.14-4.16	1.71	1.17	0.14-4.23
2002	3	2.12	1.41	0.29-4.12	2.28	1.32	0.27-3.84
2003	5	2.87	1.74	0.57-4.06	3.01	1.66	0.54-3.87
2004	6	3.55	1.69	0.55-3.69	3.73	1.61	0.53-3.51
2005	7	3.73	1.88	0.75-3.86	3.86	1.81	0.73-3.73
2006	4	4.16	0.96	0.26-2.46	4.31	0.93	0.25-2.38
2007	10	4.15	2.41	1.16-4.43	4.56	2.19	1.05-4.04
2008	6	4.35	1.38	0.45-3.00	4.77	1.26	0.41-2.74
2009	9	4.36	2.06	0.95-3.92	4.88	1.84	0.84-3.50
2010	9	5.12	1.76	0.81-3.34	4.95	1.82	0.83-3.45
2011	6	4.79	1.25	0.41-2.73	4.98	1.20	0.39-2.63

¹Standard rate: age-specific mortality rates of FVG for years 1989-2003 and 2006-2011, provided by the Istituto Superiore di Sanità (ISS); ²Standard rate: age-specific mortality rates of Italy for years 1989-2003 and 2006-2011, provided by the Istituto Superiore di Sanità (ISS). *Follow Up Period: 1989 – 2011. Cohort included 2,488 subjects, for 26,415.77 total person-years.

²Standard rate: age-specific mortality rates of Italy for years 1989-2003 and 2006-2011, provided by the Istituto Superiore di Sanità (ISS). *Follow Up Period: 1989 – 2011. Cohort included 2,488 subjects, for 26,415.77 total person-years.

Supplemental Table 5. Number of observed deaths since first surveillance enrollment, by months, for all causes of death, lung cancer and malignant neoplasm of the pleura

Months since surveillance enrollment (N) by cause of death	N. of Obs. deaths (%)	Cumulative freq. (%)
All cause of death		
≤12	22 (5.67)	22 (5.67)
13 - 24	24 (6.19)	46 (11.86)
25 - 36	25 (6.44)	71 (18.30)
37 - 48	43 (11.08)	114 (29.38)
49 - 60	30 (7.73)	144 (37.11)
60+	244 (62.89)	388 (100.00)
All cancers		
≤12	8 (4.42)	8 (1.43)
13 - 24	12 (6.63)	20 (8.57)
25 - 36	11 (6.08)	31 (17.13)
37 - 48	20 (11.05)	51 (28.18)
49 - 60	13 (7.18)	64 (35.36)
60+	117 (64.64)	181 (100.00)
Lung cancer		
≤12	1 (1.43)	1 (1.43)
13 - 24	5 (7.14)	6 (8.57)
25 - 36	-	-
37 - 48	11 (15.71)	17 (24.29)
49 - 60	6 (8.57)	23 (32.86)
60+	47 (67.14)	70 (100.00)
Malignant neoplasm of pleura		
≤12	2 (8.00)	2 (8.00)
13 - 24	2 (8.00)	4 (16.00)
25 - 36	2 (8.00)	6 (24.00)
37 - 48	4 (16.00)	10 (40.00)
49 - 60	2 (8.00)	12 (48.00)
60+	13 (52.00)	25 (100.00)