

Appendixes

Appendix Table 1. Search strategies employed in PubMed to identify scientific literature focused on the health effects of climate change

1. Search strategy to identify publications focused on heat-related illness, cardiovascular, respiratory and kidney diseases		
#1	<i>It identifies hazards/exposures</i>	("air temperature" OR "climate change" OR "climate variability" OR "global warming" OR heat OR "hot temperature" OR "heat wave*")
#2	<i>It identifies the occupational setting</i>	("occupational diseases"[MeSH Terms] OR "occupational health"[TW] OR "occupational exposure"[MeSH Terms] OR "occupational medicine"[MeSH Terms] OR "occupational risk"[TW] OR "occupational hazard"[TW] OR "Industry"[Mesh:noexp] OR "construction industry"[MeSH Terms] OR agriculture[MeSH Terms] OR "occupational group"[TW] OR "work-related"[All Fields] OR "air pollutants, occupational"[MeSH Terms] OR "working environment"[TW] OR worker*[TW])
#3	<i>It identifies the outcomes</i>	(cardiovascular OR renal OR respiratory OR kidney OR dehydration OR fatigue OR "heat stress" OR syncope OR faint* OR heatstroke OR exhaustion OR mortality OR morbidity)
	Overall search strategy	#1 AND #2 AND #3 NOT animal*
2. Search strategy used to identify publications focused on traumatic injuries and acute death		
#1	<i>It identifies hazards/exposures</i>	("air temperature" OR "climate change" OR "climate variability" OR "global warming" OR heat OR "hot temperature" OR "heat wave*" OR flood* OR landslide* OR storm* OR draught* OR wildfire* OR hurricane*)
#2	<i>It identifies the occupational setting</i>	("occupational diseases"[MeSH Terms] OR "occupational health"[TW] OR "occupational exposure"[MeSH Terms] OR "occupational medicine"[MeSH Terms] OR "occupational risk"[TW] OR "occupational hazard"[TW] OR "Industry"[Mesh:noexp] OR "construction industry"[MeSH Terms] OR agriculture[MeSH Terms] OR "occupational group*" OR "work-related"[All Fields] OR "air pollutants, occupational"[MeSH Terms] OR "working environment"[TW] OR worker*[TW])
#3	<i>It identifies the outcomes</i>	(injur* OR "acute death" OR fatalit* OR safety)
	Overall search strategy	#1 AND #2 AND #3 NOT animal* (Filters activated: English, From 2000/01/01)
3. Search strategy to identify publications focused on the expansion of vector habitats on the increased risk of infectious diseases due to climate change		

#1	<i>It identifies hazards/exposures</i>	("air temperature" OR "climate change" OR "climate variability" OR "global warming" OR heat OR "hot temperature" OR "heat wave*")
#2	<i>It identifies the occupational setting</i>	("occupational diseases"[MeSH Terms] OR "occupational health"[TW] OR "occupational exposure"[MeSH Terms] OR "occupational medicine"[MeSH Terms] OR "occupational risk"[TW] OR "occupational hazard"[TW] OR "construction industry"[MeSH Terms] OR agriculture[MeSH Terms] OR occupational group*[TW] OR "work-related"[All Fields] OR "air pollutants, occupational"[MeSH Terms] OR "working environment"[TW] OR worker*[TW] OR livestock OR aviculture)
#3	<i>It identifies the outcomes</i>	((infect* OR vector* OR zoonos* OR mosquito* OR tick*) AND (distribution OR geographic* OR region* OR emergen*))
	Overall search strategy	#1 AND #2 AND #3

Appendix Table 2 Search strategies employed to retrieve the scientific literature on the effects of climate change on work capacity and productivity

PubMed		
#1	It identifies hazards/exposures	("air temperature" OR "Climate change" OR "climate variability" OR "global warming" OR heat OR "hot temperature" OR "heat wave*")
#2	It identifies the occupational setting	("occupational diseases"[MeSH Terms] OR "occupational health"[TW] OR "occupational exposure"[MeSH Terms] OR "occupational medicine"[MeSH Terms] OR "occupational risk"[TW] OR "occupational hazard"[TW] OR "Industry"[Mesh:noexp] OR "construction industry"[MeSH Terms] OR agriculture[MeSH Terms] OR "occupational group*"[TW] OR "work-related"[All Fields] OR "air pollutants, occupational"[MeSH Terms] OR working environment[TW] OR worker*[TW])
#3	It identifies the outcomes	("work capacity" OR strike* OR productivity OR break* OR efficiency OR rest)
	Overall search strategy	#1 AND #2 AND #3 NOT animal*
SCOPUS Database		
		productivity AND "Climate change" AND worker* All fields; Filters: Economics, Sociological

Appendix Table 3 Synthesis of studies found using the first search strategy focusing on HRI

Studies making use of administrative databases								
Paper	Codes employed to identify HRI	Study period	Heat index	Administrative database	Association measure	Type of study	Lag time	Definition of heatwave employed
Adam-Poupart et al (2)	Can_Stand_Assoc_Standard Z795 07200-Effects of heat or light, 07210-Heat stroke, 07220-Heat syncope, 07280-Multiple effects of heat or light, 07290-Effects of heat or light (not elsewhere classified) including heat-related fatigue and edema	From May 1st to September 30th of each year between 1998 and 2010	Daily maximum temperature	Workers' Compensation Board (WCB) of Quebec.	IRR per 1°C increase	Incidence	0-1-2; moving average (0-1; 0-1-2)	N/A
Bonauto et al (13)	ICD-9 codes: 992.0—Heat stroke and sunstroke; 992.1—Heat syncope; 992.2—Heat cramps; 992.3—Heat exhaustion, anhydrotic; 992.4—Heat exhaustion due to salt depletion; 992.5—Heat exhaustion, unspecified; 992.6—Heat fatigue, transient; 992.7—Heat edema; 992.8—Other specified heat effects; or 992.9—Effects of heat and light, unspecified; and/or an ANSI Z16.2 type code 151 (Contact with general heat—atmosphere or environment)	January 1, 1995 and December 31, 2005	Maximum daily temperature	Workers' compensation insurance	N/A	Descriptive; Incidence rates	3 days	N/A
Fortune et al (33)	In ED records, a case of heat illness was defined on the basis of ICD-10-CA codes for conditions diagnosed as heatstroke, sunstroke, heat collapse, heat cramps, heat exhaustion, heat fatigue, heat edema	January 2004 to December 2010	N/A	Emergency Department visits and Workplace Compensation Lost Time Claims	Proportionate morbidity ratio (PMR)	Incidence rates	N/A	N/A

	<p>or other effects of heat and light.</p> <p>Among lost time claims, a case of heat illness was defined by</p> <p>information describing the nature, event or source of injury.</p>							
Fortune et al (32)	<p>(a) Heat-related <i>Main or Other Problem listed:</i> T67: Effects of heat and light X30: Exposure to excessive natural heat W92: Exposure to excessive heat of man-made origin</p>	January 1, 2004 and December 31, 2010	Maximum daily T°	Emergency department, work-related, non-scheduled visits	Generalized Linear Mixed Model, with Bayesian inference	Ecologic time series analysis	The sum of the differences between temperature on the current day and that on the previous three days (Tlag3)	N/A
Fuhrmann et al (34)	ICD-9-CM code 992.x	2007-2011	Heat emergency response plan by National Weather Service (NWS). Apparent temperature.	Emergency Department Visits	Observed versus expected ED visits	??	N/A	At least one heat product must have been issued and verified across four or more NWS county warning areas in North Carolina for five or more consecutive days. A heat event was terminated if there was a lapse in these conditions for

								at least 5 days.
Garzon-Villalba et al (37)	Incidents with a primary or secondary code that suggested a heat-related disorder (e.g., 0721–0724). EHI included acute health events related to heat such as heat stroke, heat syncope, and heat exhaustion.	May 2010-March 2011	WBGTmax	British Petroleum Workers' compensation database	IRR per 1°C above – WBGTmax 20	Incidence	1	N.A.
Gubernot et al (38)	A heat-related death is identified in CFOI as an exposure to environmental heat with the BLS Occupational Injury and Illness Classification System event/exposure code 321 and the nature code of 072	2000-2010	N/A	Census of Fatal Occupational Injuries (CFOI) database	Rate Ratios per state	Incidence	N/A	N/A
Harduar-Morano et al (42)	HRI (ICD-9-CM 992.0-992.9 and E900.0, E900.1, E900.9). Work relatedness was identified through workers' compensation as the expected payer or the presence of a work-related Ecode	2007-2011	N/A	Inpatient hospitalization (IH) and emergency department (ED) data from nine southeastern states USA	Relative Risk for different categories of independent variables	Survey, incidence	N/A	N/A

Harduar-Morano et al (44)	<p>HRI was defined as the presence of an International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM, morbidity) or Tenth Revision (ICD-10, mortality) code for the effects of heat and light (992–992.9/T67–T67.9) or an excessive heat external cause of injury (Ecode) (E900.0/X30, E900.1/W92, E900.9) [</p>	<p>May and October (2005–2012)</p>	<p>N/A</p>	<p>Emergency department (ED) and hospital discharge data from the Florida Agency for Health Care Administration. Death certificate data from the Florida Department of Health (FDOH), Bureau of Vital Statistics</p>	<p>Crude rate ratios (RR) and corresponding 95% confidence intervals (CI)</p>	<p>Survey, incidence</p>	<p>N/A</p>	<p>N/A</p>
Harduar-Morano et al (43)	<p>HRI as occurring when: presence of (1) an ICD injury diagnosis code for the effects of heat and light (ICD-9-CM codes 992 or 992.0-992.9 or ICD-10 codes T67 or T67.0-T67.9) and/or (2) the presence of an ICD-9-CM or ICD-10 code for external cause of injury for excessive heat due to weather conditions (E900.0 or X30, respectively), excessive heat due to man-made conditions (E900.1 or W92, respectively), or excessive heat of unspecified origins (E900.9) or ICD-9-CM E900</p>	<p>2005 through 2012</p>	<p>N/A</p>	<p>ED visits and hospital Discharges. data on deaths for the same years from the Florida Department of Health, Office of Vital Statistics</p>	<p>N/A</p>	<p>Survey, descriptive</p>	<p>N/A</p>	<p>N/A</p>
Mirabelli et al (70)	<p>Heat-related deaths (ICD9 codes 692.71, 992.0–992.9, E900.0,</p>	<p>1977-2001</p>	<p>Average annual summer temperatures</p>	<p>Medical examiner's records;</p>	<p>Rate ratio</p>	<p>Incidence</p>	<p>N.A.</p>	<p>N.A.</p>

	E900.1, and E900.9)			statewide temperature data from the National Climatic Data Center				
	According to the Maricopa County Department of Public Health surveillance system, using both the ICD-10 codes and key phrases in the text fields (a death was considered environmental heat-associated if it was related to heat generated by the climate (e.g., sun, humidity, etc.) and excludes heat from man-made sources such as ovens or manufacturing equipment.)	May-October 2002-2009	NOT specified	Death certificates	Assessing the risk of HR death by occupation	Case-control		
	HRI International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM) at least one among the following codes: 992.xx (Effects of heat and light), E900 (Excessive heat), E900.0 (Excessive heat due to weather conditions), or E900.9 (Excessive heat of unspecified origin)	From May through September 2008–2010	Daily maximum temperature	Emergency department (ED) data	Mean daily HRI ED visits for increase in daily maximum ambient temperature	Ecological	0	N.A.
Petitti et al (75)	Industry/Sector Codes Used to Identify Heat-Related Illness Claims OR ICD9 CM codes (705.1 Prickly heat	January 1, 1995 to December 31, 2009	Maximum daily temperature (Tmax) and Heat Index (HImax)	workers' compensation claims	Mean Tmax and mean HImax for outdoor claims (TABLE III)	Ecological	0	N.A.

	<p>276 Hyperosmolality and/or hyponatremia 276.5 Volume depletion 276.5 Volume depletion, unspecified 276.51 Dehydration 276.52 Hypovolemia 584 Acute renal failure 584.9 Acute renal failure, unspecified 992 Effects of heat and light 992 Heat stroke and sunstroke 992.1 Heat syncope 992.2 Heat cramps 992.3 Heat exhaustion 992.3 Heat exhaustion, anhidrotic 992.4 Heat exhaustion due to salt depletion 992.5 Heat exhaustion, unspecified 992.6 Heat fatigue, transient 992.7 Heat edema 992.8 Other specified heat effects 992.9 Effects of heat and light)</p>							
	<p>The Type of Occurrence Classification System Other diseases and injuries; mental disorders; nervous systems and sense organ dis; skin and subcutaneous tissue dis; digestive system dis; resp. system diseases; infectious and parasitic dis; circulatory system dis; neoplasm; cancer</p>	<p>1 July 2001 to 30 June 2010</p>	<p>Daily maximum and minimum temperatures</p>	<p>Workers' compensation claim data</p>	<p>Incidence rate ratios (IRR) with 95% CIs, and interpreted as % in the N of daily work-related injury claims during heatwave periods compared with non-heat wave periods</p>	<p>Time-series</p>	<p>N.A.</p>	<p>≥ 3 consecutive days over 35°C); ≥ 5 days of max temp in excess of 35°C OR ≥ 3 days of max temp in excess of 40°C</p>

Rhea et al (82)	claims due to heat illnesses were identified using each of the following four codes: TOOCS nature classification G313 (heat stress/heat stroke), TOOCS mechanism classification G53 (exposure to environmental heat), TOOCS agency classification G7100 (weather and water) and TOOCS agency classification G7110 (sun)	2001-2010	Daily maximum temperature (Tmax)	Workers' compensation claim data	Incidence rate ratios (IRR) with 95% CIs,	Time series analysis/Case-crossover	N.A.	≥ 3 consecutive days with Tmax $\geq 35^{\circ}\text{C}$
Spector et al (89)	Definition of HRI	Study period	Heat index	Data source	Association measure	Type of study	Lag time	Definition of heatwave employed
	Reported sudden muscle cramps; nausea or vomiting; hot, dry skin; confusion; dizziness; or fainting	2012-2013	Reported "high temperature"	Data are from PACE4, a community-based participatory research project. Participants were recruited from three agricultural counties. A total of 235 farm workers completed the baseline interviews	Prevalence of characteristics among different subgroups. Chi square test of differences	Cross-sectional	N.A.	N/A

Xiang 2 et al (99)	Reported skin rash/skin bumps, painful muscle cramps/spasms, dizziness/light-headedness, fainting, headache, heavy sweating, extreme weakness/fatigue, nausea/vomiting, and confusion.	Interviews from Jul to Aug 2013	N/A	Eligible participants were: (1) 18 years or older; (2) farm workers engaging in outdoor crop production at the time of the interview; (3) able to speak English or Spanish; and (4) able to provide informed consent	Chi-square test and Fisher's exact test. Logistic regression models. Prevalence Rate.	Cross-sectional	N/A	N/A
	Heat-related symptoms were headache, tachycardia, muscle cramps in the arms or legs, fever, nausea, difficulty breathing, swelling of hands or feet, dizziness, vomiting and fainting. Dehydration symptoms were dry mouth and dysuria	February 2011	N/A	Harvesters were interviewed. A 64 items' symptom questionnaire with questions based on the US Military recommendations	Chi-square tests and Fisher's exact test	Cross-sectional survey with control group	N/A	N/A

Xiang et al (101)	Neurological symptoms Headache Loss of co-ordination Dizziness Tingling in hands/feet Blurred vision Fainting Electrolyte imbalances Abdominal cramps Spasms Systemic symptoms Fatigue Nausea Loss of appetite Early heat stress symptoms Heavy sweating Intense thirst Anxiety Dark coloured urine output Rashes Itching skin Dry mouth Elevated temperature Hot red or flushed dry skin	2013-2014 winter and summer	WBGT	Construction workers. Structured questionnaire	N/A	Cross sectional survey	N/A	N/A
	Symptoms of HRI (sudden muscle cramps; nausea or vomiting; hot, dry skin; confusion; dizziness; fainting; headache). The outcome of interest was experiencing three or more symptoms in the past week. Because HRI symptoms are common and general symptoms, using three or more symptoms was designed to improve the identification of HRI specifically	June 11–23, 2011	During data collection for the current study, temperatures ranged between 95°F and 104°F, and the heat index was 100°F or hotter every day, peaking at 108°F)	Structured questionnaire	Ors Migrant farm-workers versus “ideal”	Cross-sectional survey	N/A	N/A
Studies making use of ad hoc questionnaire surveys	Eight heat exhaustion symptoms (headache, nausea, vomiting, weakness, fatigue, dizziness, clammy/moist skin or irritability), and seven heat stroke symptoms (high body temperature, hot and dry skin, confusion, low muscle coordination, irrational behaviour, loss of consciousness or convulsions),	N/A	N/A	Data were collected from participants via a questionnaire. In completing the questionnaire, participants indicated whether they had	Independent samples t-test One-way analysis of variance	Cross-sectional with control	N/A	N/A

				experienced any symptoms of heat illness in the past 12 months.	Pearson's Chi-square analysis Rate ratios were calculated for significant Chi-square observations			
Paper	Symptom <hr/> Feeling tired Cramps Nausea Dizziness Thirst Vomiting Confusion Muscle weakness Heat sensations Chills Feeling light-headed	N/A	WBGT	Direct measurements on workers and symptoms' referral	Independent samples t-test One-way analysis of variance Pearson's Chi-square analysis Rate ratios were calculated for significant Chi-square observations	Cross-sectional	N/A	N/A
Arcury et al (8)	Overall 42 symptoms, of them 18 MSK and the rest "General"	The survey was carried out in the middle of August 2001	WBGT	self-administered questionnaire	Chi2 test or Student's t-test	Cross-sectional	N/A	N/A

Bethel et al (11)	A section of Critical Incident Report was included in the worker data collection sheet to obtain workers' personal heat illness experiences. Reported symptoms of heat disorders include heat rash (1 case), fatigue (10 cases), feeling thirsty (3 cases), feeling discomfort (5 cases), dizziness (18 cases), difficulty in breathing (4 cases), heat cramp (3 cases), dehydration (3 cases), over sweating (4 cases), dry and hot skin (1 case), fever (8 cases), headache (6 cases), vomiting (8 cases), loss of control (2 cases), fainting (7 cases), non-sweating (1 case), and heat stroke (2 cases).	N/A	Predicted Heat Strain (PHS) model WBGT and many others	Data collection sheet for the construction workers sample, a questionnaire and interview guide for the managers sample and a site observation checklist	N/A	Ethnographic study. Qualitative study	N/A	N/A
Crowe et al (20)	A range of health symptoms including minor conditions such as heat cramps, heat syncope, and heat exhaustion to life-threatening heat stroke	August to September 2013	Mean maximum temperatures; Maximum heat index	interviewer-administered questionnaire that included standard questions related to heat-related symptoms	Logistic and log-binominal regression models were used to assess associations between sun protection behaviors and heat-related symptoms and to calculate prevalence ratios.	Cross-sectional	N.A.	N.A.

Dutta et al (24)	Only heat stroke considered	November 2003	None	Questionnaire	Multiple linear regression analysis (BODY TEMP registered)	Cross-sectional : no association between heat index and heatstroke was evaluated	N.A.	N.A.
Fleischer et al (31)	A variety of symptoms (e.g. increased thirst, excessive perspiration, itchy skin, tiredness, dry nose, blister formation, sinus problems, teary or burning eyes, exhaustion, malaise, dehydration, headaches, backache, leg pains, nose bleeds, premature baldness, dizziness)	March 2009	Hourly humidity and hourly maximum temperature	Focus group discussions and interviews	NONE	Cross-sectional	N.A.	N.A.
Hunt et al (46)	Subjective symptoms; Physiological parameters: Systolic and diastolic blood pressure and heart rate, electrolyte and blood urea	August 18 th and 19 th in 1998 ("fine days")	WBGT	Field study	Systolic and diastolic blood pressure and heart rate measurement before and during work Did not study association with WBGT data	Physiological parameters monitoring	0	N.A.

Pradhan et al (77)	Faint, tension, irritation, laziness, lack of sleep, rash, diarrhoea, eye infection, etc.	2010	Heat index, humidity index, and WBGT, based on the HOTHAPS approach	Household survey	None	Cross-sectional	N.A.	N.A.
Sahu et al (84)	NO MENTION OF HRI Health complains: pain in different parts of the body, digestive problems, CV problems, other problems (eye, ears, respiratory); Heat concerns (Aware of heat symptoms, discomfort or exhaustion during hottest days)	April to June 2011	WBGT	Questionnaire	Student t-test used to test for any significant differences between the measured parameters at different times and heat exposures	Cross-sectional	0	N.A.
Singh et al (87)	perceptions of heat discomfort, occurrence of health symptoms, heat exhaustion	summer of 2010	Not specified	semi-structured telephone interviews	No association measure used	Cross-sectional	N.A.	N.A.
Venugopal et al (92)	self-reported heat related health illnesses One of the following heat related symptoms at work i.e., excessive sweating, excessive thirst, tiredness, cramps, headache, nausea/vomiting, fainting or prickly heat	2 Seasons, once during the "hotter season" (April– June 2013) and another during the	WBGT (The minimum, maximum and standard deviation of WBGT values from each workplace was computed from the	Questionnaire	Chi square test was used to compare seasonal differences in qualitative assessments of heat for workers assessed in both	Cross-sectional	N.A.	N.A.

		“cooler season” (November–January 2012)	total number of ambient heat stress measurements taken at various locations in the each workplace).		hotter and cooler seasons.			
--	--	---	---	--	----------------------------	--	--	--

Appendix Table 4 Synthesis of studies found using the second search strategy focusing on injuries.

Paper	Study period	Heat index	Data source	Association measures	Study design	Lag time	Heat wave definition
Adam-Poupart et al (2)	2003-2010	Maximum daily outdoor temperature	Workers' Compensation Board	IRR per 1°C increase	Incidence	0-1-2; moving average (0-1; 0-1-2)	N.A.
Bonauto et al (13)	May 2010-March 2011	WBGTmax	British Petroleum Workers' compensation database	IRR per 1°C above – WBGTmax 20	Incidence	1	N.A.
Fortune et al (33)	2002-2012	Maximum daily temperature and minimum daily temperature (minimum temperature during the night before the day of injury)	Workers' compensation claims database	OR per 1°C increase in temperature measure	Case-crossover	0	N.A.
Fortune et al (32)	1998-2003	Apparent temperature (AT) AT max, AT24 and AT day were assessed	Hospital discharge data	Proportions comparison	Ecological	0 1	N.A.

Fuhrmann et al (34)	2002-2013	Daily maximum and minimum temperatures	Workers' compensation claims database	Proportions comparison	Case-crossover	N.A.	≥3 days of max temp in excess of 35°C
Garzon-Villalba et al (37)	2000-2012	Maximum daily humidex exposures	Workers' compensation claims database	OR per category of humidex index	Case-crossover	0	N.A.
Gubernot et al (38)	2005	Experiences of heat stress derived from survey question	Questionnaire survey	OR	Cohort study	N.A.	N.A.
Harduar-Morano et al (42)	2001-2010	Daily maximum and minimum temperatures	Workers' compensation claims database	IRR	Time series	N.A.	≥3 days of max temp in excess of 35°C; ≥5 days of max temp in excess of 35°C OR ≥3 days of max temp in excess of 40°C
Harduar-Morano et al (44)	2001-2010	Daily maximum temperature	Workers' compensation claims database	IRR per 1°C between 14.2°C and 37.7°C	Ecological	0	N.A.

Appendix Table 5. Papers selected using the first search strategy

Paper	Year	Country/Region	Population	Heat exposure index	Study design	Outcome	Main results
Adam-Poupart et al (1)	2014	Canada	Working population	Maximum daily temperature	Ecological	Heat-related illness	The relationship between daily counts of compensations and maximum daily temperatures was log-linear; the pooled incidence rate ratio of daily heat-related compensations per 1 °C increase in daily maximum temperatures was 1.419 (95% CI 1.326 to 1.520).
Adam-Poupart et al (2)	2015	Canada	Working population	Daily mean temperature	Case-crossover	Respiratory diseases	It was unclear if the observed trends for outdoor workers are due to ozone levels, to high temperatures or to other unmeasured parameters that are associated with them.
Arbury et al (3)	2014	USA	Working population	Heat index	Case surveillance	Heat-related illness	Nine of the 13 deaths reported occurred in the first 3 days of working on the job, 4 of them occurring on the worker's 1st day. In 7 cases, 2 or more employees had a HRI. All heat illnesses and deaths occurred on days with a heat index 29.0°C–41.0°C.
Arcury et al (4)	2015	USA	Agriculture	Ambient temperature	Cross-sectional	Heat-related illness	The majority of days were in the heat index range of 90° to 103° F. Factors associated with HRI while working outside included working in wet clothes and shoes, harvesting and topping tobacco, and spending after-work time in an extremely hot house

Balbus et al (5)	2009	USA	General population, including working population	Not specified	Narrative review	Heat-related illness	Electricity and pipeline utility workers are at increased risk of heat stress, especially if work tasks involve heavy exertion.
Bates et al (6)	2008	United Arab Emirates	Construction	TWL (thermal work limit) and WBGT	Physiological parameters monitoring	Physiological parameters	Workers were not physiologically challenged despite fluctuating harsh environmental conditions
Bates et al (7)	2010	Middle East	Construction	Not specified	Cross-sectional	Physiological parameters	Although the expatriate workers were found in general to be better hydrated than their Australian counterparts, a high proportion were still found to be inadequately hydrated both on presentation for work and throughout the shift.
Bedno et al (8)	2014	USA	Armed force	Not specified	Cohort	Heat-related illness	There were 67 incidents of Exertional heat illness (EHI). Among weight qualified (WQ) subjects, step test failure was significantly associated with EHI (OR 2.00, 95% CI 1.13, 3.53). For those passing the step test, the risk of EHI was significantly higher in those who exceeded body fat than in WQ subjects.
Berry et al (9)	2011	Australia	Agriculture	Not specified	Narrative review	Mental health	Currently there is inconclusive evidence for the proposition that (all) farmers have elevated rates of mental health problems, even in the face of extreme adversities.
Bethel et al (10)	2014	USA (Oregon)	Agriculture	Not specified	Cross-sectional	Heat-related illness	64% of respondents reported experiencing a symptom consistent with HRI during a hot day at work in the past week. Nearly 30% of participants reported experiencing

							≥ 2 HRI symptoms during the previous work; 11% experienced ≥3 HRI symptoms.
Biswas et al (11)	2011	India	Other	WBGT	Cross-sectional	Physiological parameters	The WBGT index was above the recommended range in summer. In the summer, the relative cardiac cost above 30 was recorded for 35% of the work shift in the young workers and 47% of the work shift for older workers.
Bodin et al (12)	2016	El Salvador	Agriculture	WBGT	Physiological parameters monitoring	Heat-related illness	WBGT was >26°C from 9:00 (average max.: 29.3±1.7°C). There was a reduction of HR symptoms and dehydration before and after the intervention. Individual daily production increased from 5.1 to a high of 7.3 tons/person/day.
Bonauto et al (13)	2007	USA (Washington State)	Working population	Outdoor ambient temperature	Ecological	Heat-related illness	There were 480 Washington HRI claims during the study period. The significantly increased average daily Tmax on days with multiple HRI claims compared to those with a single HRI claim suggests a dose-response effect of environmental ambient temperature.
Boodosingh et al (14)	2014	Puerto Rico	Armed force	Not specified	Case report	Heat-related illness	A case of heat stroke with altered mental status was managed with mechanical ventilatory support, intravenous fluids and external cooling measures. The man was later discharged home without any neurological sequelae.

Brake et al (15)	2003	Australia	Mining underground	WGBT	Physiological parameters monitoring	Physiological parameters	Average environmental conditions were severe (WGBT 30.9°C, SD 2.0 °C), range 25.7-35.2 °C). A majority of workers were coming to work in a moderately hypohydrated state (average urinary specific gravity 1.024, SD 0.0059).
Bray (16)	2010	USA	Agriculture	Not specified	Case report	Heat-related illness	A case-report of a 19-year-old male landscaper in Delaware who died of heat stroke on his first day of work on a hot, humid day in late June.
Brearley et al (17)	2015	Australia (Northern territory)	Electrical	WGBT	Cross-sectional	Physiological parameters	The values of gastrointestinal temperature showed a peak of 38.4°C. The mean physiological strain index (PSI) was 2.6 (=overall low strain), with periods of moderate strain. They were dehydrated prior to and following the work shift.
Clark P.A. et al (18)	2016	Central America (Nicaragua)	Agriculture	Not specified	Narrative review	Kidney diseases	Sugarcane workers are working under direct sun exposure, high temperatures, for very long hours with minimal breaks. Those working conditions alone can cause severe rhabdomyolysis and acute kidney injury. Prolonged glyphosate use and exposure further damages renal tissue and worsens kidney function. Inappropriate use of antibiotics, NSAID, diuretics can also greatly aggravate kidney damage. Unremitting acute kidney injury eventually transitions to chronic kidney disease.

Crowe et al (19)	2015	Central America (Costa Rica)	Agriculture	Not specified	Cross-sectional	Heat-related illness	Heat and dehydration symptoms were experienced at least once per week more frequently among harvesters (P<0.05). Percentages of workers reporting such symptoms increased in accordance with increasing heat exposure categories.
Delgado Cortez (20)	2009	Nicaragua	Agriculture	WBGT	Physiological parameters monitoring and productivity	Physiological parameters	Although temperature increased to maximum values early in the morning, many workers did not follow the rehydration measures and drank less than 6 L, a potentially dangerously low volume.
D'Ovidio et al (21)	2016	Global	Working population	Not specified	Narrative review	Allergies	Workplaces may be an important source of exposure to air pollution not only for dust and fumes, but also for vegetable and animal allergens.
Dutta et al (22)	2015	India	Construction	WBGT	Cross-sectional	Heat-related illness	59% of all reports in summer were positive for symptoms (from mild to severe) as compared to 41% in winter.
Edwards et al (23)	2015	Australia	General population, including working population	Drought	Cross-sectional	Mental health	Overall, 15.7 % of farmers had a mental health problem compared to 9.1 % of farm workers, 8.9 % of those in non-agricultural employment and 17.4 % of those who were not employed.
Ellis et al (24)	2017	Australia	Agriculture	Not specified	Cross-sectional	Mental health	The recently observed patterns of climate change have exacerbated farmers' worries about the weather, undermined notions of self-identity, and contributed to

							cumulative and chronic forms of place-based distress, culminating in heightened perceived risk of depression and suicide.
Farshad et al (25)	2014	Iran	Construction	WBGT and TWL	Physiological parameters monitoring	Physiological parameters	There was a significant difference in Thermal Work Limit (TWL), WBGT and urinary specific gravity (USG) between exposed and non-exposed group (P<0.01). Mean USG was 1.0213 ± 0.0054 in control group and 1.026 ± 0.005 in exposure group, which does not show a clinically dehydration status
Fleischer et al (26)	2013	USA (Georgia)	Agriculture	Ambient temperature, heat index	Cross-sectional	Heat-related illness	Temperatures ranged between 95°F and 104°F, and the heat index was 100°F or hotter every day, peaking at 108°F. Most farmworkers (71%) reported experiencing at least 1 symptom, and one third reported experiencing ≥ 3 symptoms in the past week.
Fortune et al (27)	2013	Canada (Ontario)	Working population	Not specified	Case series	Heat-related illness	The peak incidence was observed in the summer. The risk of HRI was elevated for men, young workers, manual workers and those with shorter employment tenure.
Fortune et al (28)	2014	Canada (Ontario)	Working population	Ambient temperature	Ecological	Heat-related illness	The median increase in the daily rate of emergency department (ED) visits for HRI was 75% for each degree above 22°C (posterior 95% credible interval (CI) relative rate=1.56-1.99) in the daily maximum temperature.

Fuhrmann et al (29)	2016	USA	General population, including working population	Ambient temperature	Case series	Heat-related illness - cardiorespiratory diseases	Heat exhaustion was the most common HRI, while heat stroke was relatively low (<10 % of visits). About 1/3 of all HRI visits were work-related.
Galvao Barbosa et al (30)	2012	Brazil	Agriculture	WBGT	Physiological parameters monitoring	Physiological parameters	Work during the harvest period was associated with changes in blood markers and higher blood pressure, which may be related to autonomic imbalance.
Garcia-Trabanino et al (31)	2015	El Salvador	Agriculture	Temperature, WBGT, and HI	Cross-sectional	Physiological parameters	The mean urine specific gravity, urine osmolality and creatinine, serum creatinine, uric acid and urea nitrogen increased; urinary pH, chloride and potassium decreased; pre-shift eGFR was reduced (<60 mL/min) in 14% of male workers.
Garzon-Villalba et al (32)	2016	USA	Working population	WBGTmax	Incidence	Heat-related illness	Exertional Heat Illness (EHI) were higher in workers exposed above a WBGTmax of 20°C (RR 1.40). Exposures above 28°C-WBGTmax on the day of the EHI and/or the day before were associated with higher risk of EHI (RRs from 1.0-10.4).
Gubernot et al (33)	2014	USA	Working population	Not specified	Scoping review	Heat-related illness	Around 40% of HRI cases occur among workers. The length of service appears inversely related to the occurrence of HRI. Working in direct sunshine can add up to 15°F to the perceived temperature. Increased heat decreases workers' productivity.
Gubernot et al (34)	2015	USA	Working population	Not specified	Case surveillance	Heat-related illness	A higher rate of HR deaths for the years 2005-2010 than for the years

							2000-2004 was found (RR=1.4) The largest N of deaths occurred in July (34%) (86% occurred in June-August); 65% of workers fell ill between 12 and 6:00 pm.
Hanigan et al (35)	2012	Australia	Agriculture	Hutchinson Drought Index	Time-series	Mental health	Authors found an increased relative risk of suicide of 15% (95% CI 8%-22%) for rural males aged 30-49 y when the drought index rose from the 1st to the 3rd quartile. An increased risk of suicide was observed in spring and early summer.
Hanna et al (36)	2011	Australia	Working population	Ambient temperature	Narrative review	Heat-related illness	Workplaces with risks of extreme heat exposure include outdoor and maintenance work, mining, shearing, farmwork, firefighting, and other emergency and essential services. Indoor work near heat-generating equipment with poor ventilation is also potentially hazardous on hot days. Those with high aerobic fitness can tolerate core temperatures 0.9°C higher than those who are less fit before succumbing to heat exhaustion. The impacts of extreme heat are not restricted to physiological effects: cognitive impairment and psychological and behavioural effects have been demonstrated and climate change has been linked to increased suicide rates and other mental health concerns.

Harari-Arjona et al (37)	2016	Ecuador	Agriculture	Not specified	Narrative review	Heat-related illness	Farmers suffer dehydration and frequent episodes of sudden onset of cramps in limbs, which make workers unable to work again. Also, a higher prevalence of kidney diseases has been reported. In particular, there is great concern about an epidemic CKD.
Harduar-Morano et al (38)	2015	USA	Working population	Not specified	Case series	Heat-related illness	There were 8,315 occupational HRI ED visits (6.5/100,000 workers) and 1,051 IHs (0.61/100,000). Younger workers had elevated rates for ED visits, while older workers had higher IH rates. The majority of IHs and ED visits occurred in the summer months.
Harduar-Morano et al (39)	2016	USA	General population, including working population	Not specified	Case series	Heat-related illness	The largest proportion of work-related (WR) HRI morbidity and mortality occurred in August. The WR HRI cases accounted for 0.66%, 0.98%, and 2.3% of all-cause WR ED visits, hospitalizations, and deaths during the warm season.
Harduar-Morano et al (40)	2017	USA (Florida)	General population, including working population	Not specified	Case series	Heat-related illness	There 2979 ED visits, 415 hospitalizations, and 23 deaths in patients with work-related HRI. The proportion of patients with a severe HRI diagnosis increased with data source severity.
Holmes et al (41)	2011	Dubai	Other	Not specified	Cross-sectional	Physiological parameters	55% of workers were found to be clinically hyponatraemic during the summer period compared with only 8% during the winter period.
Hunt et al (42)	2013	Australia	Mining surface	Not specified	Cross-sectional	Heat-related illness	Heat illness symptoms were experienced by 87 and 79 % of

							surface and underground mine workers, respectively ($p = 0.189$), with 81-82 % of the symptoms reported being experienced by miners on more than one occasion.
Hunt et al (43)	2014	Australia	Mining surface	Not specified	Cross-sectional	Heat-related illness	Core body temperature averaged $37.46 \pm 0.13^{\circ}\text{C}$, with the group maximum $37.98 \pm 0.19^{\circ}\text{C}$. Mean urine-specific gravity was 1.024 ± 0.007 , with 78.6% of samples 1.020 or more. 73% of workers reported at least one HR symptom during the shift.
Inaba et al (44)	2007	Japan	Construction - Traffic control	WBGT	Cross-sectional	Heat-related illness	Work difficulty due to hot weather was significantly less prevalent in traffic control workers than among construction workers ($p < 0.01$).
Jackson et al (45)	2010	USA	Working population	Not specified	Case surveillance	Heat-related illness	From 1992 through 2006, 68 workers employed in crop production and related services died from heat-related illness. The Ag/For/Fis/ Hun sector had the highest average heat fatality rate (~ 0.3 deaths/100,000 full-time workers).
Jia et al (46)	2016	China (Hong Kong)	Construction	WBGT	Case-report	Heat-related illness	Among the 207 workers, 36 valid heat illness cases (17.4%) were reported. There was a peak in HRI right before lunch, and another at 2 pm, lasting until 3 pm. No incident was reported after 4 pm.
Joubert et al (47)	2011	United Arab Emirates	Working population	Not specified	Case surveillance	Heat-related illness	One company reported a combined 79.5% decrease in cases (15.3 vs 1.16 cases per 1000 workers) while the other experienced a 50%

							reduction in serious cases (0.08-0.04 cases per 100,000 work hours).
Kearney et al (48)	2015	USA	Agriculture	Mean maximum temperatures and maximum heat index	Cross-sectional	Heat-related illness	Mean max. temp. ranged between 86°F and 95°F; max. heat index was 117°F. The prevalence of having 1 HRI symptom was 72% and 27% among workers had ≥3 HRI symptoms.
Kenefick et al (49)	2007	USA	Working population	Ambient temperature	Narrative review	Physiological parameters	Bishop et al. observed that, in simulated industrial work conditions, encapsulated protective clothing increased sweat rates up to 2.25 L/h. Likewise, wearing protective equipment such as full or half face masks can make fluid consumption more difficult and can further contribute to dehydration in the work place. Firefighters, who wear heavy protective clothing and are exposed to intense heat., were reported to have sweat rates up to 2.1 L/h. Brake et al. observed fluid losses and hydration status of mine workers under thermal stress working extended shifts (12 hours): 60% of the miners reported to work dehydrated and that their hydration status did not improve throughout the 10 to 12 hour shift.
Kjellstrom 1 et al (50)	2009	Global, with a special focus on LMIC	Working population	WBGT	Narrative review	Heat-related illness	423 heat-related deaths were reported among agricultural workers in the USA, 1992-2006.

							There are few studies in low and middle-income countries of deaths or serious clinical heat stroke among heat-exposed workers, except for India. Beyond the acute heat stress, more chronic effects on the heart and kidneys may develop after repeated excessive body heating or dehydration.
Kjellstrom et al (51)	2011	Global, with a special focus on Central America	Working population	WBGT	Narrative review	Heat-related illness	28°C is the WBGT threshold limit value for risks to health for heat-acclimatized workers carrying out moderate labor, and 27°C for heavy labour. 4% of HR deaths in the USA were in the Ag/For/Fis/ Hun sector and of these 67% were in workers employed in crop production. Above 26°C, the increasing need for rest periods can be approximated by straight lines that differ depending on the work intensity
Kjellstrom et al (52)	2016	Global	Working population	WBGT	Narrative review	Heat-related illness	Death from heat stroke have been reported among South African mine workers, US agricultural workers, as well as in China, India, Qatar, and other countries. Specific serious heat stroke symptoms and heat exhaustion were notified in many hot workplaces, including China and India. Clinical damage of organs, heart overload and kidney damage were reported among US military, Central American sugar workers and migrant construction workers in Qatar. Heat exhaustion,

							and cognitive and psychological performance effects among South African mine workers; Australian farmers. Reports on HR deaths show that symptoms of heat strain are often ignored.
Krake et al (53)	2003	USA (Grand Canyon)	Other	WBGT	Cross-sectional	Physiological parameters	All participant hiking into the canyon exceeded the heart rate and core body temperature (CBT) criteria. All but one participant developed mild dehydration; 1 case of hypernatremia occurred. The majority had a history of HR symptoms or illness
Kravchenko et al (54)	2013	USA and Europe	General population, including working population	Not specified	Narrative review	Heat-related illness	A total of 423 workers died from exposure to heat (an average annual heat-related death rate of 0.39 deaths per 100,000 workers for crop workers, compared with 0.02 deaths per 100,000 for all U.S. civilian workers) between 1992 and 2006 in the U.S.
Lao et al (55)	2016	Australia	Working population	Not specified	Cross-sectional	Heat-related illness	Common heat-related symptoms including headaches, sweating, dizziness, and tiredness. Most workers highlighted that heat impedes their work efficiency and causes a substantial slowing of work rate.
Laws et al (56)	2016	Nicaragua	Agriculture	Not specified	Physiological parameters monitoring	Kidney diseases	Mean estimated glomerular filtration rate (eGFR) was 113 mL/min/1.73 m ² and <5% of workers had albuminuria. Field workers had increases in neutrophil gelatinase-associated lipocalin and

							IL-18 levels that were 1.49 (95% CI, 1.06 to 2.09) and 1.61 (95% CI, 1.12 to 2.31) times as high, respectively, as in non-field workers.
Lundgren et al (57)	2013	Global	Working population	Not specified	Narrative review	Heat-related illness	The years 1995-2006 have rank among the warmest. Effects of increasing temperatures include a high prevalence of CKD, in Central America, especially among sugarcane workers, and a global reduction of work productivity.
Lundgren et al (58)	2014	India	Working population	WBGT	Within-group comparison	Physiological parameters	Avg WBGT was 29.7. Significant impacts on productivity in all workplaces, apart from the laundry facility, were shown, e.g. in the canteen, the core temperature limit of 38°C predicted by the model was reached in only 64 min for women.
Lunyera et al (59)	2016	Global	General population, including working population	Not specified	Systematic review	Kidney diseases	Assessment of the associations between dehydration and CKDu across studies was limited by inconsistent reporting of the measurements for assessing dehydration.
Luo et al (60)	2014	China	Other	Not specified	Case-control	Urinary system diseases	Outdoor workers were more likely to present with urolithiasis compared with indoor ones; workers with longer cumulative exposure time (OR=1.5; 95% CI: 1.2, 1.8) and abnormal blood pressure (OR=1.6; 95% CI: 1.0, 2.5) had higher risk for urolithiasis.
Maeda et al (61)	2006	Japan	Forestry	Not specified	Cross-sectional	Heat-related illness	1/3 of forestry workers developed some symptoms of early heatstroke. A short duration of

							forestry service was one of the risk factors contributing to the onset of heatstroke, in addition to heat stress, loss of body water and electrolytes, and obesity.
Marchetti et al (62)	2016	World	Working population	Not specified	Narrative review	Human health	A recent CDC report identified, during 1992-2006, a total of 423 worker deaths from exposure to environmental heat were reported in the United States. Of these 423 deaths, 102 (24%) occurred in workers employed in the agriculture, forestry, fishing, and hunting industries (rate: 0.16 per 100 000 workers)
Mathee et al (63)	2010	South Africa	Outdoor workers	Ambient temperature	Cross-sectional	Heat-related illness	Hourly max. temp. ranged from 23-36°C in Upington and 21-27°C in Johannesburg. Heat-related effects reported included sunburn, sleeplessness, irritability, and exhaustion leading to difficulty in maintaining work levels and output during very hot weather.
Mbanu et al (64)	2007	USA	Firefighters	Apparent temperature	Case-crossover	CHD	Overall, temperature was not associated with increased risk of on-duty death.
Meade et al (65)	2015	USA	Electrical	Not specified	Physiological parameters monitoring	Physiological parameters	Prior to the start of the work shift, 38% of workers were euhydrated, whereas 75% were dehydrated (USG > 1.020) after. The high levels of thermal strain were paralleled by high mean and peak HR and PSI responses, which were greater in Manual Pole Workers.

Mirabelli et al (66)	2005	USA (North Carolina)	General population, including working population	Average annual summer temperatures	Incidence	Heat-related illness	Of the 161 HR deaths, 40 occurred on the job: of these 45% were among farm labourers. All decedents whose HR injuries occurred in occupational settings were men, and most were young adults. All HR deaths among Hispanic men occurred on the job. For each 1°F increase in average summer temperature, the rate of heat-related death in the working population increased 37% (Rate Ratio=1.37; 95% CI=0.99, 1.90).
Mirabelli et al (67)	2010	USA (North Carolina)	Agriculture	Not specified	Cross-sectional	Heat-related illness	Working in extreme heat was reported by 281 respondents (94%), among whom 112 (40%) reported symptoms of heat illness. 37% reported changes in their work hours and 34% in their work activities.
MMWR -	2008	USA	Working population	Not specified	Case surveillance	Heat-related illness	423 HR deaths were reported among workers. Of these, 24% occurred in the agriculture, forestry, fishing, and hunting industries and 2/3 of whom occurred in the crop production. A not statistically significant increase in fatality rates over time was observed.
CDC (68)	2013	Iran	Construction	Thermal Work Limit	Physiological parameters monitoring	Physiological parameters	In the exposed group, 38% of workers had a USG level between 1.026-1.030 and 12.72% had a USG level above 1.030 (in the control group 15.2% and 0.58%, respectively). The Pearson

							correlation measure showed a significant correlation between USG and TWL.
Montazer et al (69)	2006	Japan	Construction	WBGT	Physiological parameters monitoring	Heat-related illness	The WBGT outdoors varied from 23 to 34 °C. Few items of subjective symptoms increased after work compared with before work. Electrolytes and blood urea nitrogen did not change. Blood sugar before work was higher than before lunch and after work.
Morioka et al (70)	2017	USA	Agriculture	Not specified	Physiological parameters monitoring	Kidney diseases	Among men, 10 experienced heat strain, as estimated by physiological strain index (PSI) ≥ 7.5 , and 3 of those met criteria for Acute kidney injury (AKI); 35 participant were incident AKI (12.3%). Adjusted OR of AKI was 1.34 (95% CI 1.04 to 1.74) among those experiencing heat strain.
Moyce et al (71)	2011	USA	Armed force	Not specified	Case surveillance	Heat-related illness	Overall, in 2010 crude incidence rates of "heat stroke" and "other heat injury" were 0.21 and 1.77 per 1,000 p-yrs, respectively. The overall unadjusted incidence rate of "other heat injury" was higher in 2010 than in any prior year of the period.
MSMR - Armed Forces Health Surveillance Branch (72)	2016	USA	Armed force	Not specified	Case surveillance	Heat-related illness	The incidence rate of heat stroke in 2015 was higher than rates in the previous 4 years. More service members were treated for "other heat injuries" in 2015 (n=1,933) than in either of the previous 2 years.

MSMR - Armed Forces Health Surveillance Branch (73)	2016	Finland	General population, including working population	Mean temperature, highest daily temperature	Cross-sectional	Heat-related cardiorespiratory symptoms	In the summer preceding the survey, the temperatures exceeded the long-term average by 1-2 °C in the areas studied. Farmers had a higher risk of presenting HR cardiorespiratory symptoms (OR 2.27; 1.14-4.46) compared with those working in industry.
Näyhä et al (74)	2013	USA (Northern Illinois)	Agriculture	Temperature and humidity	Case report	Heat-related illness	Case-report of multiple organ failure due to heatstroke.
Nevarez (75)	2014	Finland	Electrical	Ambient temperature	Physiological parameters monitoring	Physiological parameters	AT varied between -30°C and +28°C. Average cardiorespiratory strain was $48 \pm 3\%VO_{2max}$. 40% exceeded the recommended 50% VO_{2max} level. Both muscular and cardiorespiratory strain may reduce work efficiency. Thermal strain remained at a tolerable level.
Oksa et al (76)	2014	Global	General population, including working population	Not specified	Systematic review	Heat-related illness	The adverse health aspects related to climate change may include heat-related disorders, such as heat stress and respiratory disorders, including those exacerbated by air pollution and aeroallergens, such as asthma.
Patz et al (77)	2013	Australia	Mining surface	WBGT	Physiological parameters monitoring	Physiological parameters	A significant effect for time was observed for core temperature with greater mean core temperatures measured mid-shift ($37.5 \pm 0.4^\circ C$) and post-shift ($37.6 \pm 0.3^\circ C$) compared with pre-shift values ($37.0 \pm 0.5^\circ C$).
Peiffer et al (78)	2012	El Salvador	Agriculture	Not specified	Cross-sectional	Physiological parameters	The adjusted ORs of decreased kidney function for 10-year increments of coastal sugarcane or

							cotton plantation work were 3.1 (95% CI, 2.0-5.0) in men and 2.3 (95% CI, 1.4-3.7) in women.
Peraza et al (79)	2013	USA (Arizona)	Working population	Not specified	Case-control	Heat-related illness	75% of HR deaths occurred in men: the age-adjusted ORs for HR death were 2.32 (95% CI 1.55, 3.48) for construction/extraction and 3.50 (95% CI 1.94, 6.32) for agriculture occupations. Men aged ≥ 65 y.o. in agriculture were at especially high risk.
Petitti et al (80)	2011	Australia	Agriculture	Not specified	Cross-sectional	Mental health	Older farmers faced the same drought-related pressures as their younger peers but these pressures were compounded by the discomforts of ageing and by fighting a losing battle to cope with rapid social and agricultural change.
Polain et al (81)	2013	Nepal	Manufacturing industry	Heat index, humidity index, and WBGT, based on the HOTHAPS approach	Cross-sectional	Heat-related illness	The average temperature in the summer reached to over 39°C: the environmental conditions were inadequate for workers to work continuously during the day. HRI included fainting, mental irritation, laziness, sleepless nights, dehydration and giddiness.
Pradhan et al (82)	2014	Nicaragua	Agriculture	Not specified	Cross-sectional with nested case-control analysis	Kidney diseases	Prevalence of GFR $<60\text{mL}/\text{min}/1.73\text{m}^2$ was 10% among women and 42% among men. Risk factors included increased lifetime days cutting sugarcane in the dry season, non-deliberate pesticide inhalation and sugarcane chewing.

Raines et al (83)	2014	India	Working population	Not specified	Case series	Kidney diseases	Most of the CKD (48.4%) cases were registered relatively high between the months of March and May; most were agricultural workers (24%), construction workers or laborers (23%) and industrial labor workers (19%) who belong to low income group.
Raju et al (84)	2012	North Carolina	General population, including working population	Daily maximum temperature	Ecological	Heat-related illness	7.6% of HRI ED visits were work-related. These were more common than other causes in 19-45 year-olds. A threshold ambient maximum daily temperature of 98°F is apparent. The mean daily number of HRI ED visits increased by 1.4 for each 1°F increase from 90°F to 98°F and by 15.8 for each 1°F increase from 98°F to 100°F. Beyond 100°F, the mean number of HRI ED visits decreased with an increasing daily maximum ambient temperature: between 100°F and 102°F, the mean daily number of HRI ED visits decreased by 12.5 for each 1°F increase.
Rhea et al (85)	2016	Central America	Agriculture	Not specified	Physiological parameters monitoring	Kidney diseases	The presence of urate crystals in the urine and high end-of-workday urinary uric acid concentrations were common. Hyperuricemia may induce glomerular hypertension, whereas the increased urinary uric acid may directly injure renal tubules.

Roncal-Jimenez et al (86)	2013	India	Agriculture	WBGT	Cross-sectional	Heat-related illness: health complains: pain in different parts of the body, digestive problems, CV problems, other problems (eye, ears, respiratory); Heat concerns (Aware of heat symptoms, discomfort or exhaustion during hottest days)	Most workers reported exhaustion and pain in hot days. Heart rate recovered more slowly at high heat, indicating cardiovascular strain. Peak HR was significantly higher in the air temperature (Ta) ranges of 31-33.5°C (p<0.05) and 35-36°C (p<0.001) than at 28-30°C. The % of resting heart rate also increases with temperature, as do the sum of recovery heartbeats and the cardiac cost variables.
Sahu et al (87)	2014	India	Manufacturing industry	WBGT	Cross-sectional	Cardiovascular parameters	The cardiac parameters were significantly higher on hotter days for the brick molders.
Sett et al (88)	2015	Australia	Working population	Not specified	Cross-sectional	Heat-related illness	All interviewees reported that exposure to high temperatures was routine for workers during the summer months, and many reported varying degrees of symptomatology of heat-related illnesses. More than half of respondents (15/20) reported known cases of heat exhaustion in their industry, and several offered that there were frequent episodes each summer amongst the workforce.

Singh et al (89)	2014	USA (Washington State)	Agriculture - Forestry	Maximum daily temperature (Tmax) and Heat Index (HI _{max})	Ecological	Heat-related illness	There were 84 HRI claims and 60 citations. HRI claims and citations were most common in crop production and support subsectors; 95% of outdoor claims occurred between May and September, and 84% occurred between June and August. The mean (IQR) Tmax for outdoor HRI claims was 95 (89, 100) °F, and the mean (IQR) HI _{max} was 99 (90, 106) °F.
Spector et al (90)	2010	Arabian Gulf	Armed force	Ambient Air Temperature	Cohort	Heat-related illness	No cases of heat illness were reported due to employing simple strategies, such as maintaining hydration, rest periods outside of hot workspaces and using cooling techniques.
Stevenson et al (91)	2010	Thailand	Working population	Not specified	Cross-sectional	Human health	Working under heat stress conditions is associated with worse overall health and psychological distress (aOR ranging from 1.49 to 1.84). The heat stress effects are increased for workers >45 years, the better educated, and for work located in rural areas.
Tawatsupa et al (92)	2012	Thailand	Working population	Experiences of heat stress derived from survey question	Cross-sectional	Kidney diseases	A significant association between heat stress and incident kidney disease was observed in men (aOR = 1.48, 95% CI: 1.01-2.16). The incidence of kidney disease was even higher among men aged ≥35 years in a physical job.
Tawatsupa et al (93)	2015	India	Working population	WBGT	Cross-sectional	Heat-related illness	Workplace WBGT > 27 °C was associated with significantly

							more heat-related symptoms (74%; chi square = 5.152, p = 0.023), as compared to workplaces where WBGT ≤ 27 °C. Workers with heavy workloads reported more heat-related health issues (chi square = 23.67, p \leq 0.001) and reduced productivity (chi square = 15.82, p \leq 0.001), especially outdoor workers. Of the workers' assessed in both hotter and cooler seasons, 65% vs. 32% were exposed to higher than recommended WBGT levels in hotter and cooler seasons respectively. This corresponded to a significant increase in self-reported heat-related health impacts (p = 0.011) as well as an increase of 1.8 °C–4.3 °C WBGT in the hotter season, as compared to the cooler.
Venugopal et al (94)	2016	India	Agriculture - Manufacturing industry	WBGT	Cross-sectional	Physiological parameters	Avg. WBGT was 30°C. Above normal CBT, Sweat rate (SwR), and USG in about 10% women indicated heat strain and moderate dehydration; those who had no access to proper toilet facilities reported more HR symptoms (n=194) ($\chi^2=4.03$, p=0.0444), including dehydration.
Venugopal et al (95)	2003	Austria	Construction	Not specified	Case report	Heat-related illness	A case of heatstroke-induced acute hepatic failure with complete recovery under conservative management in a 24 year-old male is reported.

Wagner et al (96)	2016	Nicaragua	Agriculture	Not specified	Physiological parameters monitoring	Kidney diseases	The pre-shift renal function decreased significantly during 9 weeks of work in the cane cutters. Mean serum creatinine increased (20%), mean eGFR decreased (9%, 10mL/min), blood urea nitrogen (BUN) increased (41%), and mean urinary Neutrophil gelatinase-associated lipocalin (NGAL) increased four times.
Wesseling 1 et al (97)	2016	Nicaragua	Agriculture - Construction	Not specified	Cross-sectional	Kidney diseases	The study found evidence for more frequent heat stress, dehydration and kidney dysfunction among sugarcane cutters, and to a lesser degree also reduced kidney function among construction workers but not among small-scale farmers.
Wesseling et al (98)	2015	North America	Firefighters	Not specified	Narrative review	Heat-related illness	The trend in fatalities follows that of climate change, with a general upward trend in deaths from the 1980s to the present.
Withen (99)	2014	Australia	Working population	Daily maximum and minimum temperatures	Time-series	Heat-related illness	The mean Tmax during heat waves was 38.8 °C. Daily claims increased significantly by 6.2% during heatwaves for outdoor industries. Burns, wounds, lacerations, amputations and HRI were significantly associated with heatwaves.

Xiang et al (100)	2014	Global	Working population	25 (60%) studies used WGBT, 7 (16%) used subjective heat stress and 6 (14%) used air temperature	Systematic review	Heat-related illness	Manual workers be at risk of heat stress, especially those in LMIC. 79% of identified studies indicated that participants were suffering from heat strain, with outdoor workplaces (90%) being much higher than indoor workplaces (65%).
Xiang et al (101)	2015	Australia	Working population	Daily maximum temperature (Tmax)	Time series analysis/Case-crossover	Heat-related illness	The overall risk of HRI was positively associated with Tmax, especially when Tmax was over 35.5 °C. 1°C increase of Tmax was associated with a 12.7% increase of occupational heat illness claims. The risk was 4-7 times higher during heatwaves periods.
Xiang et al (102)	2017	China (Beijing)	General population, including working population	Daily maximum temperature, daily mean relative humidity, and daily mean atmospheric pressure	Ecological	CVD diseases	When the daily maximum temperature was > 33 °C from the 10th consecutive day onward, the excess mortality risk (ER) of CVD death among outdoor workers was 149%, considerably higher than the ER for the overall population (87%; p < 0.05).
Yin et al (103)	2016	USA (Colorado)	Agriculture	Daily maximum, mean and minimum temperatures, and mean apparent temperature (meAT)	Time-series	CVD diseases	Estimates of heat effects on average daily clinic visits among migrant farm workers were positive (88.0%, 95% CI: 26.2% to 180.0%). Authors did not observe statistically significant associations between heat and clinic visits among other groups.

References

1. Adam-Poupart A, Smargiassi A, Busque MA, Duguay P, Fournier M, Zayed J, et al. Summer outdoor temperature and occupational heat-related illnesses in Quebec (Canada). *Environ Res* [Internet]. Elsevier; 2014;134:339–44. Available from: <http://dx.doi.org/10.1016/j.envres.2014.07.018>
2. Adam-Poupart A, Labrèche F, Busque MA, Brand A, Duguay P, Fournier M, et al. Association between outdoor ozone and compensated acute respiratory diseases among workers in Quebec (Canada). *Ind Health* [Internet]. 2015;53:171–5. Available from: <http://dx.doi.org/10.2486/indhealth.2014-0136>
3. Arbury S, Jacklitsch B, Farquah O, Hodgson M, Lamson G, Martin H, et al. Heat illness and death among workers - United States, 2012–2013. *MMWR Morb Mortal Wkly Rep* [Internet]. 2014 Aug 8;63(31):661–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25102413>
4. Arcury TA, Summers P, Talton JW, Chen H, Sandberg JC, Spears Johnson CR, et al. Heat Illness Among North Carolina Latino Farmworkers. *J Occup Environ Med* [Internet]. 2015 Dec;57(12):1299–304. Available from: <http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00043764-201512000-00007>
5. Balbus John M, Malina Catherine M. Identifying Vulnerable Subpopulations for Climate Change Health Effects in the United States. *J Occup Environ Med*. 2009;51(1):33–7.
6. Bates GP, Schneider J. Hydration status and physiological workload of UAE construction workers: A prospective longitudinal observational study. *J Occup Med Toxicol* [Internet]. 2008;3:21. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2561022&tool=pmcentrez&rendertype=abstract>
7. Bates GP, Miller VS, Joubert DM. Hydration status of expatriate manual workers during summer in the Middle East. *Ann Occup Hyg*. 2010;54(2):137–43.
8. Bedno SA, Urban N, Boivin MR, Cowan DN. Fitness, obesity and risk of heat illness among army trainees. *Occup Med (Chic Ill)*. 2014;64(6):461–7.
9. Berry HL, Hogan A, Owen J, Rickwood D, Fragar L. Climate Change and Farmers' Mental Health: Risks and Responses. *Asia-Pacific J Public Heal* [Internet]. 2011;23(2 Suppl):119S–132S. Available from: <http://aph.sagepub.com/cgi/doi/10.1177/1010539510392556>
10. Bethel JW, Harger R. Heat-related illness among Oregon farmworkers. *Int J Environ Res Public Health*. 2014;11(9):9273–85.
11. Biswas R, Saha P, Samanta A. Cardiac strain of confectionery worker in relation to heat exposure during regular work shift. *Indian J Occup Environ Med* [Internet]. 2011;15(3):120. Available from: <http://www.ijoem.com/text.asp?2011/15/3/120/93202>
12. Bodin T, García-Trabanino R, Weiss I, Jarquín E, Glaser J, Jakobsson K, et al. Intervention to reduce heat stress and improve efficiency among sugarcane workers in El Salvador: Phase 1. *Occup Environ Med* [Internet]. 2016;oemed-2016-103555. Available from: <http://oem.bmj.com/lookup/doi/10.1136/oemed-2016-103555>
13. Bonauto D, Anderson R, Rauser E, Burke B. Occupational Heat Illness in Washington State , 1995 – 2005. *Am J Ind Med*. 2007;950(50):1995–2005.
14. Boodosingh DR, Robles-Arias C, Alemán-Ortiz JR, Rodríguez-Cintrón W. A rare cause of altered mental status and fever in a young military recruit in Puerto Rico. *P R Health Sci J*. 2014;33(4):200–2.
15. Brake DJ, Bates GP. Fluid losses and hydration status of industrial workers under thermal stress working extended shifts. *Occup Environ Med* [Internet]. 2003;60(2):90–6. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1740457&tool=pmcentrez&rendertype=abstract>
16. Bray P. Heat-Related Illnesses: Opportunities for Prevention. *J Occup Environ Med*. 2010;52(8):844–5.
17. Brearley M, Harrington P, Lee D, Taylor R. Working in Hot Conditions—A Study of Electrical Utility Workers in the Northern Territory of Australia. *J Occup Environ Hyg* [Internet]. 2015 Mar 4;12(3):156–62. Available from: <http://www.tandfonline.com/doi/full/10.1080/15459624.2014.957831>
18. Clark PA, Chowdhury J, Chan B, Radigan N. Chronic kidney disease in Nicaraguan sugarcane workers: A historical, medical, environmental analysis and ethical analysis. *Internet J Third World Med* [Internet]. 2016;12(1):1–16. Available from:

- <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=emed18b&AN=613372262%5Cnhttps://www.mobilelibrary.com/vlib/order/OpenURLReceive.aspx?sid=OVID:embase&id=pmid:&id=doi:10.5580%2FIJWH.39545&issn=1539-4646&isbn=&volume=12&issue=1&spage=no+>
19. Crowe J, Nilsson M, Kjellstrom T, Wesseling C. Heat-Related symptoms in sugarcane harvesters. *Am J Ind Med*. 2015;58(5):541–8.
 20. Cortez OD. Heat stress assessment among workers in a Nicaraguan sugarcane farm. *Glob Health Action*. 2009;2(1):1–6.
 21. D’Ovidio MC, Annesi-Maesano I, D’Amato G, Cecchi L. Climate change and occupational allergies: an overview on biological pollution, exposure and prevention. *Ann Ist Super Sanita* [Internet]. 52(3):406–14. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27698299>
 22. Dutta P, Rajiva A, Andhare D, Azhar G, Tiwari A, Sheffield P, et al. Perceived heat stress and health effects on construction workers. *Indian J Occup Environ Med* [Internet]. 2015;19(3):151. Available from: <http://www.ijoem.com/text.asp?2015/19/3/151/174002>
 23. Edwards B, Gray M, Hunter B. The Impact of Drought on Mental Health in Rural and Regional Australia. *Soc Indic Res*. 2015;121(1):177–94.
 24. Ellis NR, Albrecht GA. Climate change threats to family farmers’ sense of place and mental wellbeing: A case study from the Western Australian Wheatbelt. *Soc Sci Med* [Internet]. Elsevier Ltd; 2017;175:161–8. Available from: <http://dx.doi.org/10.1016/j.socscimed.2017.01.009>
 25. Farshad A, Montazer S, Monazzam MR, Eyvazlou M, Mirkazemi R. Heat stress level among construction workers. *Iran J Public Health*. 2014;43(4):492–8.
 26. Fleischer NL, Tiesman HM, Sumitani J, Mize T, Amarnath KK, Bayakly AR, et al. Public health impact of heat-related illness among migrant farmworkers. *Am J Prev Med* [Internet]. Elsevier Inc.; 2013;44(3):199–206. Available from: <http://dx.doi.org/10.1016/j.amepre.2012.10.020>
 27. Fortune MK, Mustard CA, Etches JJC, Chambers AG. Work-attributed illness arising from excess heat exposure in Ontario, 2004–2010. *Can J Public Heal*. 2013;104(5):2004–10.
 28. Fortune M, Mustard C, Brown P. The use of Bayesian inference to inform the surveillance of temperature-related occupational morbidity in Ontario, Canada, 2004–2010. *Environ Res* [Internet]. Elsevier; 2014;132:449–56. Available from: <http://dx.doi.org/10.1016/j.envres.2014.04.022>
 29. Fuhrmann CM, Sugg MM, Konrad CE, Waller A. Impact of Extreme Heat Events on Emergency Department Visits in North Carolina (2007–2011). *J Community Health*. Springer US; 2016;41(1):146–56.
 30. Barbosa CMG, Terra-Filho M, de Albuquerque ALP, Di Giorgi D, Grupi C, Negrão CE, et al. Burnt Sugarcane Harvesting - Cardiovascular Effects on a Group of Healthy Workers, Brazil. *PLoS One*. 2012;7(9):1–10.
 31. García-Trabanino R, Jarquín E, Wesseling C, Johnson RJ, González-Quiroz M, Weiss I, et al. Heat stress, dehydration, and kidney function in sugarcane cutters in El Salvador - A cross-shift study of workers at risk of Mesoamerican nephropathy. *Environ Res* [Internet]. Elsevier; 2015;142:746–55. Available from: <http://dx.doi.org/10.1016/j.envres.2015.07.007>
 32. Garzon-Villalba XP, Mbah A, Wu Y, Hiles M, Moore H, Schwartz SW, et al. Exertional heat illness and acute injury related to ambient wet bulb globe temperature. *Am J Ind Med*. 2016;59(12):1169–76.
 33. Gubernot DM, Anderson GB, Hunting KL. The epidemiology of occupational heat exposure in the United States: a review of the literature and assessment of research needs in a changing climate. *Int J Biometeorol* [Internet]. 2014 Oct 11;58(8):1779–88. Available from: <http://link.springer.com/10.1007/s00484-013-0752-x>
 34. Gubernot DM, Anderson GB, Hunting KL. Characterizing occupational heat-related mortality in the United States, 2000–2010: An analysis using the census of fatal occupational injuries database. *Am J Ind Med*. 2015;58(2):203–11.
 35. Hanigan IC, Butler CD, Kocic PN, Hutchinson MF. Suicide and drought in New South Wales, Australia, 1970–2007. *Proc Natl Acad Sci*. 2012;109(35):13950–5.

36. Hanna EG, Kjellstrom T, Bennett C, Dear K. Climate Change and Rising Heat: Population Health Implications for Working People in Australia. *Asia-Pacific J Public Heal* [Internet]. 2011 Mar 1;23(2 Suppl):14S–26S. Available from: <http://aph.sagepub.com/cgi/doi/10.1177/1010539510391457>
37. Harari Arjona R, Piñeiros J, Ayabaca M, Harari Freire F. Climate change and agricultural workers' health in Ecuador: occupational exposure to UV radiation and hot environments. *Ann Ist Super Sanita* [Internet]. 52(3):368–73. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27698295>
38. Harduar Morano L, Bunn TL, Lackovic M, Lavender A, Dang GTT, Chalmers JJ, et al. Occupational heat-related illness emergency department visits and inpatient hospitalizations in the southeast region, 2007-2011. *Am J Ind Med*. 2015;58(10):1114–25.
39. Harduar Morano L, Watkins S, Kintziger K. A Comprehensive Evaluation of the Burden of Heat-Related Illness and Death within the Florida Population. *Int J Environ Res Public Health* [Internet]. 2016 May 31;13(6):551. Available from: <http://www.mdpi.com/1660-4601/13/6/551>
40. Harduar Morano L, Watkins S. Evaluation of Diagnostic Codes in Morbidity and Mortality Data Sources for Heat-Related Illness Surveillance. *Public Health Rep* [Internet]. 2017;132(3):326–35. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28379784><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC5415256><http://journals.sagepub.com/doi/10.1177/0033354917699826>
41. Holmes NA, Miller VS, Schneider J, Hasan O, Bates GP. Plasma sodium levels and dietary sodium intake in manual workers in the middle east. *Ann Occup Hyg*. 2011;55(4):397–402.
42. Hunt AP, Parker AW, Stewart IB. Symptoms of heat illness in surface mine workers. *Int Arch Occup Environ Health*. 2013;86(5):519–27.
43. Hunt AP, Parker AW, Stewart IB. Heat strain and hydration status of surface mine blast crew workers. *J Occup Environ Med* [Internet]. 2014;56(4):409–14. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24504322>
44. Inaba R, Kurokawa J, Mirbod SM. Comparison of subjective symptoms and cold prevention measures in winter between traffic control workers and construction workers in Japan. *Ind Health*. 2009;47(3):283–91.
45. Jackson LL, Rosenberg HR. Preventing heat-related illness among agricultural workers. *J Agromedicine*. 2010;15(April):200–15.
46. Jia YA, Rowlinson S, Ciccarelli M. Climatic and psychosocial risks of heat illness incidents on construction site. *Appl Ergon* [Internet]. Elsevier Ltd; 2016;53:25–35. Available from: <http://dx.doi.org/10.1016/j.apergo.2015.08.008>
47. Joubert D, Thomsen J, Harrison O. Safety in the heat: A comprehensive program for prevention of heat illness among workers in Abu Dhabi, United Arab Emirates. *Am J Public Health*. 2011;101(3):395–8.
48. Kearney GD, Hu H, Xu X, Hall M, Balanay JA. Estimating the Prevalence of Heat-Related Symptoms and Sun Safety Related Behavior among Latino Farmworkers in Eastern North Carolina. *J Agromedicine* [Internet]. 2015;813(April):15–23. Available from: <http://dx.doi.org/10.1080/1059924X.2015.1106377>
49. Kenefick RW, Sawka MN. Hydration at the work site. *J Am Coll Nutr*. 2007;26(5 Suppl):597S–603S.
50. Kjellstrom T, Holmer I, Lemke B. Workplace heat stress, health and productivity-an increasing challenge for low and middle-income countries during climate change. *Glob Health Action*. 2009;2(1):1–6.
51. Kjellstrom T, Crowe J. Climate Change, Workplace Heat Exposure, and Occupational Health and Productivity in Central America. *Int J Occup Environ Health* [Internet]. 2011 Jul;17(3):270–81. Available from: <http://www.tandfonline.com/doi/full/10.1179/107735211799041931>
52. Kjellstrom T, Briggs D, Freyberg C, Lemke B, Otto M, Hyatt O. Heat, Human Performance, and Occupational Health: A Key Issue for the Assessment of Global Climate Change Impacts. *Annu Rev Public Health* [Internet]. 2016 Mar 18;37(1):97–112. Available from: <http://www.annualreviews.org/doi/10.1146/annurev-publhealth-032315-021740>

53. Krake A, McCullough J, King B. Health Hazards to Park Rangers from Excessive Heat at Grand Canyon National Park. *Appl Occup Environ Hyg* [Internet]. 2003 May;18(5):295–317. Available from: <http://www.tandfonline.com/doi/abs/10.1080/10473220301364>
54. Kravchenko J, Abernethy AP, Fawzy M, Lyerly HK. Minimization of heatwave morbidity and mortality. *Am J Prev Med* [Internet]. Elsevier Inc.; 2013;44(3):274–82. Available from: <http://dx.doi.org/10.1016/j.amepre.2012.11.015>
55. Lao J, Hansen A, Nitschke M, Hanson-Easey S, Pisaniello D. Working smart: An exploration of council workers' experiences and perceptions of heat in Adelaide, South Australia. *Saf Sci* [Internet]. Elsevier Ltd; 2016;82:228–35. Available from: <http://dx.doi.org/10.1016/j.ssci.2015.09.026>
56. Laws RL, Brooks DR, Amador JJ, Weiner DE, Kaufman JS, Ramírez-Rubio O, et al. Biomarkers of Kidney Injury among Nicaraguan Sugarcane Workers. *Am J Kidney Dis*. 2016;67(2):209–17.
57. Lundgren K, Kuklane K, Gao C, Holmér I. Effects of heat stress on working populations when facing climate change. *Ind Health* [Internet]. 2013;51(1):3–15. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23411752>
58. Lundgren K, Kuklane K, Venugopal V. Occupational heat stress and associated productivity loss estimation using the PHS model (ISO 7933): A case study from workplaces in Chennai, India. *Glob Health Action*. 2014;7(1):1–9.
59. Lunyera J, Mohottige D, von Isenburg M, Jeuland M, Patel UD, Stanifer JW. CKD of uncertain etiology: A systematic review. *Clin J Am Soc Nephrol*. 2016;11(3):379–85.
60. Luo H, Turner LR, Hurst C, Mai H, Zhang Y, Tong S. Exposure to ambient heat and urolithiasis among outdoor workers in Guangzhou, China. *Sci Total Environ* [Internet]. Elsevier B.V.; 2014;472:1130–6. Available from: <http://dx.doi.org/10.1016/j.scitotenv.2013.11.042>
61. Maeda T, Kaneko SY, Ohta M, Tanaka K, Sasaki A, Fukushima T. Risk factors for heatstroke among Japanese forestry workers. *J Occup Health*. 2006;48(4):223–9.
62. Marchetti E, Capone P, Freda D. Climate change impact on microclimate of work environment related to occupational health and productivity. *Ann Ist Super Sanita* [Internet]. 2016;52(3):338–42. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27698292>
63. Mathee A, Oba J, Rose A. Climate change impacts on working people (the HOTHAPS initiative): findings of the South African pilot study Angela. *Glob Health Action*. 2010;3(5612):1–9.
64. Mbanu I, Wellenius GA, Mittleman MA, Peeples L, Stallings LA, Kales SN. NIH Public Access. 2013;24(4):715–26.
65. Meade RD, Lauzon M, Poirier MP, Flouris AD, Kenny GP. An Evaluation of the Physiological Strain Experienced by Electrical Utility Workers in North America. *J Occup Environ Hyg* [Internet]. 2015;12(10):708–20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26011148>
66. Mirabelli MC, Richardson DB. Heat-related fatalities in North Carolina. *Am J Public Health*. 2005;95(4):635–7.
67. Mirabelli MC, Quandt SA, Crain R, Grzywacz JG, Robinson EN, Vallejos QM, et al. Symptoms of heat illness among Latino farm workers in North Carolina. *Am J Prev Med* [Internet]. Elsevier Inc.; 2010;39(5):468–71. Available from: <http://dx.doi.org/10.1016/j.amepre.2010.07.008>
68. Centers for Disease Control and Prevention (CDC). Heat-related deaths among crop workers--United States, 1992--2006. *MMWR Morb Mortal Wkly Rep* [Internet]. 2008 Jun 20;57(24):649–53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18566563>
69. Montazer S, Farshad AA, Monazzam MR, Eyvazlou M, Yaraghi AAS, Mirkazemi R. Assessment of construction workers' hydration status using urine specific gravity. *Int J Occup Med Environ Health* [Internet]. 2013;26(5):762–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24464540>
70. Morioka I, Miyai N, Miyashita K. Hot environment and health problems of outdoor workers at a construction site. *Ind Health*. 2006;44(3):474–80.
71. Moyce S, Mitchell D, Armitage T, Tancredi D, Joseph J, Schenker M. Heat strain, volume depletion and kidney function in California agricultural workers. *Occup Environ Med* [Internet]. 2017;d:oemed-2016-103848. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28093502>

72. Armed Forces Health Surveillance Center (AFHSC). Update: heat injuries, active component, U.S. Armed Forces, 2010. *MSMR* [Internet]. 2011 Mar;18(3):6–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21793607>
73. Armed Forces Health Surveillance Branch. Update: Heat injuries, active component, U.S. Army, Navy, Air Force, and Marine Corps, 2015. *MSMR* [Internet]. 2016 Mar;23(3):16–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27030928>
74. Näyhä S, Rintamäki H, Donaldson G, Hassi J, Jousilahti P, Laatikainen T, et al. The prevalence of heat-related cardiorespiratory symptoms: the vulnerable groups identified from the National FINRISK 2007 Study. *Int J Biometeorol* [Internet]. *International Journal of Biometeorology*; 2017;61(4):657–68. Available from: <http://link.springer.com/10.1007/s00484-016-1243-7>
75. Nevarez J. OSHA Compliance Issues. *J Occup Environ Hyg* [Internet]. 2013 Jun;10(6):D67–70. Available from: <http://www.tandfonline.com/doi/abs/10.1080/15459624.2013.784177>
76. Oksa J, Hosio S, Mäkinen T, Lindholm H, Rintamäki H, Rissanen S, et al. Muscular, cardiorespiratory and thermal strain of mast and pole workers. *Ergonomics* [Internet]. 2014 May 4;57(5):669–78. Available from: <http://www.tandfonline.com/doi/abs/10.1080/00140139.2014.895854>
77. Patz JA, Frumkin H, Holloway T, Vimont DJ, Haines A. Climate Change. *Jama* [Internet]. 2014;312(15):1565. Available from: <http://jama.jamanetwork.com/article.aspx?doi=10.1001/jama.2014.13186>
78. Peiffer JJ, Abbiss CR. Thermal stress in North Western Australian iron ore mining staff. *Ann Occup Hyg*. 2013;57(4):519–27.
79. Peraza S, Wesseling C, Aragon A, Leiva R, García-Trabanino RA, Torres C, et al. Decreased kidney function among agricultural workers in El Salvador. *Am J Kidney Dis* [Internet]. Elsevier Inc.; 2012;59(4):531–40. Available from: <http://dx.doi.org/10.1053/j.ajkd.2011.11.039>
80. Petitti DB, Harlan SL, Chowell-Puente G, Ruddell D. Occupation and Environmental Heat-Associated Deaths in Maricopa County, Arizona: A Case-Control Study. *PLoS One*. 2013;8(5).
81. Polain JD, Berry HL, Hoskin JO. Rapid change, climate adversity and the next “big dry”: Older farmers’ mental health. *Aust J Rural Health*. 2011;19(5):239–43.
82. Pradhan B, Shrestha S, Shrestha R, Pradhanang S, Kayastha B, Pradhan P. Assessing climate change and heat stress responses in the Tarai region of Nepal. *Ind Health* [Internet]. 2013;51(1):101–12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23411760>
83. Raines N, González Quiroz M, Wyatt C, Kurzrok M, Pool C, Lemma T, et al. Risk factors for reduced glomerular filtration rate in a nicaraguan community affected by mesoamerican nephropathy. *MEDICC Rev* [Internet]. 2014;16(2):16–22. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24878645>
84. Raju DSSK, Kiranmayi P, Vijaya Rachel K. Climate change and chronic kidney disease. *Asian J Pharm Clin Res* [Internet]. 2014;7(2):53–7. Available from: <http://innovareacademics.in/journals/index.php/ajpcr/article/download/953/615%5Cnhttp://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed12&NEWS=N&AN=2014240557>
85. Rhea S, Ising A, Fleischauer AT, Deyneka L, Vaughan-Batten H, Waller A. Using near real-time morbidity data to identify heat-related illness prevention strategies in North Carolina. *J Community Health*. 2012;37(2):495–500.
86. Roncal-Jimenez C, García-Trabanino R, Barregard L, Lanaspá MA, Wesseling C, Harra T, et al. Heat Stress Nephropathy from Exercise-Induced Uric Acid Crystalluria: A Perspective on Mesoamerican Nephropathy. *Am J Kidney Dis*. 2016;67(1):20–30.
87. Sahu S, Sett M, Kjellstrom T. Heat exposure, cardiovascular stress and work productivity in rice harvesters in India: implications for a climate change future. *Ind Health* [Internet]. 2013;51(4):424–31. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23685851>
88. Sett M, Sahu S. Effects of occupational heat exposure on female brick workers in West Bengal, India. *Glob Health Action*. 2014;7(1):1–11.

89. Singh S, Hanna EG, Kjellstrom T. Working in Australia's heat: Health promotion concerns for health and productivity. *Health Promot Int*. 2015;30(2):239–50.
90. Spector JT, Krenz J, Rauser E, Bonauto DK. Heat-related illness in Washington State agriculture and forestry sectors. *Am J Ind Med*. 2014;57(8):881–95.
91. Stevenson T, Roiz de Sa D. Managing the risk of heat illness on board type 23 frigates deploying to the Arabian Gulf. *J R Nav Med Serv*. 2010;96(3):150–7.
92. Tawatsupa B, Lim LL-Y, Kjellstrom T, Seubsman S-A, Sleight A, The Thai Cohort Study Team. The association between overall health, psychological distress, and occupational heat stress among a large national cohort of 40,913 Thai workers. *Glob Health Action*. 2010;3(9):1–10.
93. Tawatsupa B, Lim LL-Y, Kjellstrom T, Seubsman S, Sleight A, Study Team the TC. Association Between Occupational Heat Stress and Kidney Disease Among 37 816 Workers in the Thai Cohort Study (TCS). *J Epidemiol* [Internet]. 2012;22(3):251–60. Available from: <http://japanlinkcenter.org/JST.JSTAGE/jea/JE20110082?from=CrossRef&type=abstract>
94. Venugopal V, Chinnadurai JS, Lucas RAI, Kjellstrom T. Occupational heat stress profiles in selected workplaces in India. *Int J Environ Res Public Health*. 2015;13(1):1–13.
95. Venugopal V, Rekha S, Manikandan K, Latha PK, Vennila V, Ganesan N, et al. Heat stress and inadequate sanitary facilities at workplaces an occupational health concern for women? *Glob Health Action*. 2016;1(March 2017):1–9.
96. Wagner M, Kaufmann P, Fickert P, Trauner M, Lackner C, Stauber RE. Successful conservative management of acute hepatic failure following exertional heatstroke. *Eur J Gastroenterol Hepatol* [Internet]. 2003;15(10):1135–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/14501625>
97. Wesseling C, Aragón A, González M, Weiss I, Glaser J, Bobadilla NA, et al. Kidney function in sugarcane cutters in Nicaragua - A longitudinal study of workers at risk of Mesoamerican nephropathy. *Environ Res* [Internet]. Elsevier; 2016;147:125–32. Available from: <http://dx.doi.org/10.1016/j.envres.2016.02.002>
98. Wesseling C, Aragón A, González M, Weiss I, Glaser J, Rivard CJ, et al. Heat stress, hydration and uric acid: a cross-sectional study in workers of three occupations in a hotspot of Mesoamerican nephropathy in Nicaragua. *BMJ Open* [Internet]. 2016;6(12):e011034. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27932336%0Ahttp://bmjopen.bmj.com/lookup/doi/10.1136/bmjopen-2016-011034>
99. Withen P. Climate change and wildland firefighter health and safety. *New Solut a J Environ Occup Heal policy*. 2015;24(4):577–84.
100. Xiang J, Bi P, Pisaniello D, Hansen A. The impact of heatwaves on workers' health and safety in Adelaide, South Australia. *Environ Res* [Internet]. Elsevier; 2014;133:90–5. Available from: <http://europepmc.org/abstract/med/24906072%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/24906072>
101. Xiang J, Bi P, Pisaniello D, Hansen A. Health impacts of workplace heat exposure: An epidemiological review. *Ind Health* [Internet]. 2014;52(2):91–101. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24366537>
102. Xiang J, Hansen A, Pisaniello D, Bi P. Extreme heat and occupational heat illnesses in South Australia, 2001-2010. *Occup Environ Med* [Internet]. 2015;72(8):580–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26081622>
103. Yin Q, Wang J. The association between consecutive days' heat wave and cardiovascular disease mortality in Beijing, China. *BMC Public Health* [Internet]. BMC Public Health; 2017;17(1):223. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28228117%5Cnhttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC5322604>
104. Zhang K, Arauz RF, Chen T-H, Cooper SP. Heat effects among migrant and seasonal farmworkers: a case study in Colorado. *Occup Environ Med*. 2016;73(5):324–8.

Appendix Table 6. Papers selected using the second search strategy

Papers selected using search strategy N.2						
Paper	Year	Country/Region	Population	Heat exposure index	Study design	Main results
Adam-Poupart 1 et al (1)	2015	Canada	Working population	Maximum daily outdoor temperature	Ecological	The relationship between daily counts of compensations and maximum daily temperatures reached statistical significance for three health regions. The incidence rate ratio of daily compensations per 1°C increase was 1.002 (95%CI 1.002 to 1.003).
Bonafede et al (2)	2016	Global	Working population	Daily maximum temperature daily mean temperature or apparent temperature	Systematic review	Only 2 studies analysed the dose-response relationship between temperature and the health outcomes finding a reversed U-shaped exposure-response relationship, or linear relationship or linear above/below a threshold.
Dutta et al (3)	2015	India	Construction	WBGT	Cross-sectional	Injuries reported in summer were among workers with < 36 months of experience.
Garzon-Villalba et al (4)	2016	USA	Working population	WBGTmax	Incidence	Acute injuries (AI) were higher in workers exposed above a WBGTmax of 20°C (RR 1.06/°C). The unadjusted model showed a 13% increment in risk to experience an AI when the ambient thermal conditions increase 1°C-WBGT.
McInnes et al (5)	2017	Australia	Working population	Maximum daily temperature and minimum daily temperature (minimum temperature during the night before the day of injury)	Case-crossover	Significant positive associations between temperature and acute injury were seen for younger workers, with the odds of injury increasing by 1% for each 1°C increase in daily minimum temperature, and by 0.8% for each 1°C increase in daily maximum temperature.
Morabito et al (6)	2006	Italy (Tuscany)	Working population	Apparent temperature (AT) AT max, AT24 and	Ecological	The peak of work-related accidents occurred on days characterized by high, but not extreme, thermal conditions. The maximum peak of work accidents in each month occurred on days

				AT day were assessed		characterized by a mean ATday ranging between 24.8°C and 27.5 °C.
Nag et al (7)	2001	India	Manufacturing industry	Ambient temperature	Case surveillance	Accident prevalence was significantly high in the summer months (May-June) when the ambient temperature ranged between 42 and 48°C (hot-dry). Permanent night workers were more vulnerable and less tolerant to heat compared to the rotating day workers.
Otte-im-Kampe et al (8)	2016	High-income countries	General population, including working population	Not specified	Systematic review	Two out of 3 studies found an increase in work-related accidents during increased temperatures; the other reported no association.
Rameezdeen et al (9)	2017	Australia	Construction	Daily maximum and minimum temperatures	Case-crossover	The number of accidents reported during heat wave periods was less compared to control periods, suggesting some control measures are in operation in construction sites to prevent accidents during heat waves
Spector et al (10)	2016	USA	Agriculture	Maximum daily humidex exposures	Case-crossover	The traumatic injury OR was 1.14 (95% CI 1.06, 1.22), 1.15 (95% CI 1.06, 1.25), and 1.10 (95% CI 1.01, 1.20) for daily maximum humidex of 25±29, 30±33, and ≥34, respectively, compared to < 25, adjusted for self-reported duration of employment.
Tawatsupa et al (11)	2013	Thailand	Working population	Experiences of heat stress derived from survey question	Cohort	Nearly 20% of workers experienced occupational heat stress, which strongly and significantly associated with occupational injury (adjusted OR 2.12, 95%CI 1.87-2.42 for males and 1.89, 95%CI 1.64-2.18 for females).
Withen (12)	2015	North America	Firefighters	Not specified	Narrative review	The trend in fatalities follows that of climate change, with a general upward trend in deaths from the 1980s to the present.

Xiang et al (13)	2014	Australia	Working population	Daily maximum temperature	Ecological	A reversed U-shaped exposure-response relationship between Tmax and overall daily injury claims was observed. Overall, a 1°C increase in maximum temperature between 14.2°C and 37.7°C was associated with a 0.2% increase in daily injury claims.
Xiang et al (14)	2014	Australia	Working population	Daily maximum and minimum temperatures	Time-series	The mean Tmax during heat waves was 38.8 °C. Daily claims increased significantly by 6.2% during heatwaves for outdoor industries. Burns, wounds, lacerations, amputations and HRI were significantly associated with heatwaves.
Xiang et al (15)	2014	Global	Working population	25 (60%) studies used WGBT, 7 (16%) used subjective heat stress and 6 (14%) used air temperature	Systematic review	Manual workers are at risk of heat stress, especially those in LMIC; 79% of identified studies indicated that participants were suffering from heat strain, with outdoor workplaces (90%) being much higher than indoor workplaces (65%).

References

1. Adam-Poupart A, Smargiassi A, Busque M-A, Duguay P, Fournier M, Zayed J, et al. Effect of summer outdoor temperatures on work-related injuries in Quebec (Canada). *Occup Environ Med* [Internet]. 2015;72(5):338–45. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25618108>
2. Bonafede M, Marinaccio A, Asta F, Schifano P, Michelozzi P, Vecchi S. The association between extreme weather conditions and work-related injuries and diseases. A systematic review of epidemiological studies. *Ann Ist Super Sanita* [Internet]. 52(3):357–67. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27698294>
3. Dutta P, Rajiva A, Andhare D, Azhar G, Tiwari A, Sheffield P, et al. Perceived heat stress and health effects on construction workers. *Indian J Occup Environ Med* [Internet]. 2015;19(3):151. Available from: <http://www.ijoem.com/text.asp?2015/19/3/151/174002>
4. Garzon-Villalba XP, Mbah A, Wu Y, Hiles M, Moore H, Schwartz SW, et al. Exertional heat illness and acute injury related to ambient wet bulb globe temperature. *Am J Ind Med*. 2016;59(12):1169–76.
5. McInnes JA, Akram M, Macfarlane EM, Keegel T, Sim MR, Smith P. Association between high ambient temperature and acute work-related injury: A case-crossover analysis using workers' compensation claims data. *Scand J Work Environ Heal*. 2017;43(1):86–94.
6. Morabito M, Cecchi L, Crisci A, Modesti PA, Orlandini S. Relationship between work-related accidents and hot weather conditions in Tuscany (central Italy). *Ind Health* [Internet]. 2006;44(3):458–64. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16922190>
7. Nag PK, Nag A. Shiftwork in the hot environment. *J Hum Ergol (Tokyo)* [Internet]. 2001;30(1–2):161–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/14564876>
8. Otte im Kampe E, Kovats S, Hajat S. Impact of high ambient temperature on unintentional injuries in high-income countries: a narrative systematic literature review. *BMJ Open* [Internet]. 2016;6(2):e010399. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26868947>
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4762150>
<http://bmjopen.bmj.com/lookup/doi/10.1136/bmjopen-2015-010399>
9. Rameezdeen R, Elmualim A. The impact of heat waves on occurrence and severity of construction accidents. *Int J Environ Res Public Health*. 2017;14(1):1–13.
10. Spector JT, Bonauto DK, Sheppard L, Busch-Isaksen T, Calkins M, Adams D, et al. A case-crossover study of heat exposure and injury risk in outdoor agricultural workers. *PLoS One* [Internet]. 2016;11(10):30–3. Available from: <http://dx.doi.org/10.1371/journal.pone.0164498>
11. Tawatsupa B, Yiengprugsawan V, Kjellstrom T, Berecki-Gisolf J, Seubsman S-A, Sleigh A. Association between heat stress and occupational injury among Thai workers: findings of the Thai Cohort Study. *Ind Health* [Internet]. 2013;51(1):34–46. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23411755>
12. Withen P. Climate change and wildland firefighter health and safety. *New Solut a J Environ Occup Heal policy*. 2015;24(4):577–84.
13. Xiang J, Bi P, Pisaniello D, Hansen A, Sullivan T. Association between high temperature and work-related injuries in Adelaide, South Australia, 2001–2010. *Occup Environ Med* [Internet]. 2014;71(4):246–52. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24334260>

14. Xiang J, Bi P, Pisaniello D, Hansen A. The impact of heatwaves on workers' health and safety in Adelaide, South Australia. *Environ Res* [Internet]. Elsevier; 2014;133:90–5. Available from: <http://europepmc.org/abstract/med/24906072><http://www.ncbi.nlm.nih.gov/pubmed/24906072>
15. Xiang J, Bi P, Pisaniello D, Hansen A. Health impacts of workplace heat exposure: An epidemiological review. *Ind Health* [Internet]. 2014;52(2):91–101. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24366537>

Appendix Table 7. Papers selected using the fourth search strategy

Paper	Year	Country/Region	Population	Heat exposure index	Study design	Main results
Altinsoy et al (1)	2015	Turkey	Working population	WBGT	Simulation based on climate driven model	Between 2071 and 2100, deficiency in labour productivity may reach up to 52 % during the summer across so Central Anatolia, Cyprus, and parts of the Aegean and eastern Mediterranean coastal areas, particularly in agriculture and construction.
Bodin et al (2)	2016	El Salvador	Agriculture	WBGT	Physiological parameters monitoring	WBGT was >26°C from 9:00 (average max.: 29.3±1.7°C). There was a reduction of HR symptoms and dehydration before and after the intervention. Individual daily production increased from 5.1 to a high of 7.3 tons/person/day.
Delgado_Cortez (3)	2009	Nicaragua	Agriculture	WBGT	Physiological parameters monitoring and productivity	Output production increased significantly (p=0.005) among those best hydrated, from 5.5 to 8 tons of cut sugarcane per worker per day.
Dell et al (4)	2014	Global	Working population	Not specified	Narrative review	Many of the papers discussed seek to address the impact of "harvesting". Temperatures can also affect income through labour productivity, which can in turn affect health.
Esteban et al (5)	2009	Taiwan	Working population	Not specified	Simulation based on climate driven model	Annual downtime from tropical cyclones could increase from 1.5% nowadays to up to 2.2% by 2085, an increase of almost 50%. This decrease in productivity could result in a loss of up to 0.7% of the annual Taiwanese GDP by 2085.
Gubernot et al (6)	2014	USA	Working population	Not specified	Scoping review	Around 40% of HRI cases occur among workers. The length of service appears inversely related to the occurrence of HRI. Working in direct sunshine can add up to 15°F to the perceived temperature. Increased heat decreases workers' productivity.
Hanna et al (7)	2011	Australia	Working population	Ambient temperature	Narrative review	Workers carried out on average 1 hour less work per day when temperatures exceeded 37°C (compared with days <30°C), as workers self-paced to maintain thermal

						comfort. Heat-related health risks increase when work is "externally paced".
Heal et al (8)	2016	Global	Working population	Not specified	Systematic review	A day with temperatures exceeding 32°C can increase local monthly mortality rates by more than 1% and reduces daily labour supply in exposed sectors by up to 14%.
Kenefick et al (9)	2007	USA	Working population	Ambient temperature	Narrative review	A study showed that when subjects were dehydrated, productivity of stacking and debarking pulpwood was reduced by 12%. Decision-making and cognitive performance are also adversely influenced by dehydration
Kershaw et al (10)	2013	United Kingdom	Other	Different levels of climate change	Simulation based on climate driven model	As we move further into the future there is an increase in summertime overheating and hence a decrease in productivity but also a decrease in cold related winter discomfort and an associated increase in productivity.
Kjellstrom (11)	2016	Global	Working population	WBGT	Simulation based on climate driven model	In South-East Asia as much as 15% to 20% of annual work hours may already be lost in heat-exposed jobs, and this may double by 2050. The annual cost of reduced labour productivity at country level already in 2030 can be several percent of GDP.
Kjellstrom et al (12)	2009	Global, with a special focus on LMIC	Working population	WBGT	Narrative review	The resulting work capacity during different hours for a person who works at a heavy work intensity of 500 W is very low: on average only 20% of work capacity remains at 12 noon.
Kjellstrom et al (13)	2009	Global	Working population	WBGT	Simulation based on climate driven model	In terms of absolute change in labour productivity by the 2080s, the greatest losses (11.4% to 26.9%) are seen under A2 in Southeast Asia, Andean and Central America, Eastern Sub-Saharan Africa, and the Caribbean.
Kjellstrom et al (14)	2011	Global, with a special focus on Central America	Working population	WBGT	Narrative review	24% of HR deaths in the USA were in the Ag/For/Fis/Hun sector and of these 67% were in workers employed in crop production. Above 26°C, the increasing need for rest periods can be approximated by straight lines that differ depending on the work intensity

Kjellstrom et al (15)	2013	South-East Asia	Working population	WBGT	Simulation based on climate driven model	Heavy work in the shade is now affected in the hottest areas so that 50-60% of afternoon work time is lost due to heat. Heavy work in the sun has losses up to 80% and above, and even moderate work is affected: >50% of afternoon work time is lost.
Kjellstrom et al (16)	2016	Global	Working population	WBGT	Narrative review	Reduced work capacity, labour productivity, and economic loss, as well as heat impact on gross domestic product were reported in India, USA and South Africa. When hourly WBGT exceeds 26°C, work capacity is reduced in heavy-labour jobs, and above 32°C (90°F) any work activity is made difficult. GDP losses will be greater than 20% by 2100.
Langkulsen et al (17)	2010	Thailand	Agriculture - Construction	WBGT	Cross-sectional	Productivity as perceived by the workers revealed that for more than half (60%) the workers, productivity loss varied from 10 to 66.7%, whereas vegetable field workers displayed no loss of productivity.
Lao et al (18)	2016	Australia	Working population	Not specified	Cross-sectional	Common heat-related symptoms including headaches, sweating, dizziness, and tiredness. Most workers highlighted that heat impedes their work efficiency and causes a substantial slowing of work rate.
Li et al (19)	2016	China	Construction	WBGT	Within-group comparison	Direct work time decreased by 0.57% and idle time increased by 0.74% when the WBGT increased by 1 °C; direct work time increased by 0.33% when the workers' experience increased by 1 year and decreased by 0.72% when the workers' age increased by 1 year.
Lundgren et al (20)	2013	Global	Working population	Not specified	Narrative review	The years 1995-2006 have rank among the warmest. Effects of increasing temperatures include a high prevalence of CKD, in Central America, especially among sugarcane workers, and a global reduction of work productivity.
Lundgren et al (21)	2014	India	Working population	WBGT	Within-group comparison	Avg WBGT was 29.7. Significant impacts on productivity in all workplaces, apart from the laundry facility, were shown, e.g. in the canteen, the core temperature limit of 38°C predicted by the model was reached in only 64 min for women.

Marchetti et al (22)	2016	World	Working population	Not specified	Narrative review	Productivity is affected after 1 hour of moderate physical work above 32 °C. In SE Asia up to 20% of annual work hours may already be lost in heat-exposed jobs. By 2080, the greatest losses are foreseen in SE Asia, Central America, Sub-Saharan Africa.
Mathee et al (23)	2010	South Africa	Outdoor workers	Ambient temperature	Cross-sectional	Hourly max. temp. ranged from 23-36°C in Upington and 21-27°C in Johannesburg. Heat-related effects reported included sunburn, sleeplessness, irritability, and exhaustion leading to difficulty in maintaining work levels and output during very hot weather.
Mirabelli et al (24)	2010	USA (North Carolina)	Agriculture	Not specified	Cross-sectional	Working in extreme heat was reported by 281 respondents (94%), among whom 112 (40%) reported symptoms of heat illness. 37% reported changes in their work hours and 34% in their work activities.
Oyekale (25)	2015	Nigeria	Agriculture	Not specified	Cross-sectional	Missing regular times scheduled for spraying cocoa pods (45.7% in Ondo state) was one of the forms of reported climate change induced occupational stresses
Patz et al (26)	2014	Global	General population, including working population	Not specified	Systematic review	Heat stress has reduced labour capacity by 10% in summer's peak over the past few decades. Projected reduction may double by 2050.
Pradhan et al (27)	2013	Nepal	Manufacturing industry	Heat index, humidity index, and WBGT, based on the HOTHAPS approach	Cross-sectional	The average temperature in the summer reached to over 39°C: the environmental conditions were inadequate for workers to work continuously during the day. HRI included fainting, mental irritation, laziness, sleepless nights, dehydration and giddiness.
Sahu et al (28)	2013	India	Agriculture	WBGT	Cross-sectional	At WBGT>26°C the hourly N of rice bundles collected was reduced approximately 5% per °C of increased WBGT.
Sett et al (29)	2014	India	Manufacturing industry	WBGT	Cross-sectional	There is a linear decline in productivity with an increase in maximum air temperature above 34.9°C, and the lost productivity for every degree rise in temperature is about 2%.

Singh et al (30)	2015	Australia	Working population	Not specified	Cross-sectional	Productivity is significantly reduced at 35°C, and approximately one-third of baseline work productivity can be lost in certain physically demanding jobs when working at 40°C.
Suzuki-Parker et al (31)	2016	Japan	Working population	WBGT	Simulation based on climate driven model	"Light labour safe" hours are projected to decrease by 30-40 % by the end of the 21st century. The number of "heavy labour restricted days" is projected to increase from ~5 days in the 2000s to nearly 2/3 of the days in August in the 2090s.
Venugopal et al (32)	2015	India	Working population	WBGT	Cross-sectional	Workers with heavy workloads reported more heat-related health issues (chi square = 23.67, $p \leq 0.001$) and reduced productivity (chi square = 15.82, $p \leq 0.001$), especially outdoor workers. Of the workers' assessed in both hotter and cooler seasons, 65% vs. 32% were exposed to higher than recommended WBGT levels in hotter and cooler seasons respectively. This corresponded to a significant increase in self-reported productivity losses ($p \leq 0.016$).
Xiang et al (33)	2014	Global	Working population	WBGT, heat stress and air temperature	Systematic review	Manual workers be at risk of heat stress, especially those in LMIC. 79% of identified studies indicated that participants were suffering from heat strain, with outdoor workplaces (90%) being much higher than indoor workplaces (65%).
Zander et al (34)	2015	Australia	Working population	Not specified	Cross-sectional	70% said heat made them less productive on at least 1 day in the previous year: on average, they were 35% less productive on days on which they had suffered from heat, were less productive on 10 days, and worked for 27.1 h less.
Zhao et al (35)	2016	China	Outdoor workers	Daily Tmax	Simulation based on climate driven model	The total HTS is estimated at 38.6 billion yuan/y over the 1979-2005 period (0.2% of GDP), the share of GDP devoted to HTS could become as high as 3% at the end of 21st century.

Zivin et al (36)	2014	USA	Working population	Historical and forecasted temperature distribution - daily maximum and minimum temperature, precipitation, snowfall, and relative humidity.	Econometric model	At daily max. temp. >85°F, workers in industries with high exposure to climate reduce daily time allocated to labour by as much as 1 h. Almost all of the decrease in time allocated to labour happens at the end of the day.
------------------	------	-----	--------------------	---	-------------------	---

References

1. Altinsoy H, Yildirim HA. Labor productivity losses over western Turkey in the twenty-first century as a result of alteration in WBGT. *Int J Biometeorol.* 2015;59(4):463–71.
2. Bodin T, García-Trabanino R, Weiss I, Jarquín E, Glaser J, Jakobsson K, et al. Intervention to reduce heat stress and improve efficiency among sugarcane workers in El Salvador: Phase 1. *Occup Environ Med* [Internet]. 2016;oemed-2016-103555. Available from: <http://oem.bmj.com/lookup/doi/10.1136/oemed-2016-103555>
3. Cortez OD. Heat stress assessment among workers in a Nicaraguan sugarcane farm. *Glob Health Action.* 2009;2(1):1–6.
4. Dell M, Jones BF, Olken BA. What Do We Learn from the Weather? The New Climate-Economy Literature †. *J Econ Lit* [Internet]. 2014 Sep;52(3):740–98. Available from: <http://pubs.aeaweb.org/doi/abs/10.1257/jel.52.3.740>
5. Esteban M, Webersik C, Shibayama T. Effect of a global warming-induced increase in typhoon intensity on urban productivity in Taiwan. *Sustain Sci.* 2009;4(2):151–63.
6. Gubernot DM, Anderson GB, Hunting KL. The epidemiology of occupational heat exposure in the United States: a review of the literature and assessment of research needs in a changing climate. *Int J Biometeorol* [Internet]. 2014 Oct 11;58(8):1779–88. Available from: <http://link.springer.com/10.1007/s00484-013-0752-x>
7. Hanna EG, Kjellstrom T, Bennett C, Dear K. Climate Change and Rising Heat: Population Health Implications for Working People in Australia. *Asia-Pacific J Public Heal* [Internet]. 2011 Mar 1;23(2 Suppl):14S–26S. Available from: <http://aph.sagepub.com/cgi/doi/10.1177/1010539510391457>
8. Heal G, Park J. Reflections—Temperature Stress and the Direct Impact of Climate Change: A Review of an Emerging Literature. *Rev Environ Econ Policy* [Internet]. 2016 Jul;10(2):347–62. Available from: <https://academic.oup.com/reep/article-lookup/doi/10.1093/reep/rew007>
9. Kenefick RW, Sawka MN. Hydration at the work site. *J Am Coll Nutr.* 2007;26(5 Suppl):597S–603S.

10. Kershaw T, Lash D. Investigating the productivity of office workers to quantify the effectiveness of climate change adaptation measures. *Build Environ*. Elsevier Ltd; 2013;69:35–43.
11. Kjellstrom T. Impact of Climate Conditions on Occupational Health and Related Economic Losses: A New Feature of Global and Urban Health in the Context of Climate Change. *Asia-Pacific J Public Heal*. 2015;1–10.
12. Kjellstrom T, Holmer I, Lemke B. Workplace heat stress, health and productivity-an increasing challenge for low and middle-income countries during climate change. *Glob Health Action*. 2009;2(1):1–6.
13. Kjellström T, Kovats RS, Lloyd SJ, Holt T, Tol RSJ. The direct impact of climate change on regional labor productivity. *Arch Environ Occup Health* [Internet]. 2009;64(4):217–27. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20007118>
14. Kjellstrom T, Crowe J. Climate Change, Workplace Heat Exposure, and Occupational Health and Productivity in Central America. *Int J Occup Environ Health* [Internet]. 2011 Jul;17(3):270–81. Available from: <http://www.tandfonline.com/doi/full/10.1179/107735211799041931>
15. Kjellstrom T, Lemke B, Otto M. Mapping occupational heat exposure and effects in South-East Asia: ongoing time trends 1980-2011 and future estimates to 2050. *Ind Health* [Internet]. 2013;51(1):56–67. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23411757>
16. Kjellstrom T, Briggs D, Freyberg C, Lemke B, Otto M, Hyatt O. Heat, Human Performance, and Occupational Health: A Key Issue for the Assessment of Global Climate Change Impacts. *Annu Rev Public Health* [Internet]. 2016 Mar 18;37(1):97–112. Available from: <http://www.annualreviews.org/doi/10.1146/annurev-publhealth-032315-021740>
17. Langkulsen U, Vichit-Vadakan N, Taptagaporn S. Health impact of climate change on occupational health and productivity in Thailand. *Glob Health Action* [Internet]. 2010 Dec 9;3. Available from: <http://www.globalhealthaction.net/index.php/gha/article/view/5607>
18. Lao J, Hansen A, Nitschke M, Hanson-Easey S, Pisaniello D. Working smart: An exploration of council workers' experiences and perceptions of heat in Adelaide, South Australia. *Saf Sci* [Internet]. Elsevier Ltd; 2016;82:228–35. Available from: <http://dx.doi.org/10.1016/j.ssci.2015.09.026>
19. Li X, Chow KH, Zhu Y, Lin Y. Evaluating the impacts of high-temperature outdoor working environments on construction labor productivity in China: A case study of rebar workers. *Build Environ* [Internet]. Elsevier Ltd; 2016;95:42–52. Available from: <http://dx.doi.org/10.1016/j.buildenv.2015.09.005>
20. Lundgren K, Kuklane K, Gao C, Holmér I. Effects of heat stress on working populations when facing climate change. *Ind Health* [Internet]. 2013;51(1):3–15. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23411752>
21. Lundgren K, Kuklane K, Venugopal V. Occupational heat stress and associated productivity loss estimation using the PHS model (ISO 7933): A case study from workplaces in Chennai, India. *Glob Health Action*. 2014;7(1):1–9.
22. Marchetti E, Capone P, Freda D. Climate change impact on microclimate of work environment related to occupational health and productivity. *Ann Ist Super Sanita* [Internet]. 2016;52(3):338–42. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27698292>
23. Mathee A, Oba J, Rose A. Climate change impacts on working people (the HOTHAPS initiative): findings of the South African pilot study Angela. *Glob Health Action*. 2010;3(5612):1–9.
24. Mirabelli MC, Quandt SA, Crain R, Grzywacz JG, Robinson EN, Vallejos QM, et al. Symptoms of heat illness among Latino farm workers in North Carolina. *Am J Prev Med* [Internet]. Elsevier Inc.; 2010;39(5):468–71. Available from: <http://dx.doi.org/10.1016/j.amepre.2010.07.008>
25. Oyekale AS. Climate change induced occupational stress and reported morbidity among cocoa farmers in South-Western Nigeria. *Ann Agric Environ Med*. 2015;22(2):357–61.

26. Patz JA, Frumkin H, Holloway T, Vimont DJ, Haines A. Climate Change. *Jama* [Internet]. 2014;312(15):1565. Available from: <http://jama.jamanetwork.com/article.aspx?doi=10.1001/jama.2014.13186>
27. Pradhan B, Shrestha S, Shrestha R, Pradhanang S, Kayastha B, Pradhan P. Assessing climate change and heat stress responses in the Tarai region of Nepal. *Ind Health* [Internet]. 2013;51(1):101–12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23411760>
28. Sahu S, Sett M, Kjellstrom T. Heat exposure, cardiovascular stress and work productivity in rice harvesters in India: implications for a climate change future. *Ind Health* [Internet]. 2013;51(4):424–31. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23685851>
29. Sett M, Sahu S. Effects of occupational heat exposure on female brick workers in West Bengal, India. *Glob Health Action*. 2014;7(1):1–11.
30. Singh S, Hanna EG, Kjellstrom T. Working in Australia's heat: Health promotion concerns for health and productivity. *Health Promot Int*. 2015;30(2):239–50.
31. Suzuki-Parker A, Kusaka H. Future projections of labor hours based on WBGT for Tokyo and Osaka, Japan, using multi-period ensemble dynamical downscale simulations. *Int J Biometeorol*. 2016;60(2):307–10.
32. Venugopal V, Chinnadurai JS, Lucas RAI, Kjellstrom T. Occupational heat stress profiles in selected workplaces in India. *Int J Environ Res Public Health*. 2015;13(1):1–13.
33. Xiang J, Bi P, Pisaniello D, Hansen A. Health impacts of workplace heat exposure: An epidemiological review. *Ind Health* [Internet]. 2014;52(2):91–101. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24366537>
34. Zander KK, Botzen WJW, Oppermann E, Kjellstrom T, Garnett ST. Heat stress causes substantial labour productivity loss in Australia. *Nat Clim Chang* [Internet]. 2015;5(May):1–6. Available from: <http://www.nature.com/doi/10.1038/nclimate2623>
35. Zhao Y, Sultan B, Vautard R, Braconnot P, Wang HJ, Ducharne A. Potential escalation of heat-related working costs with climate and socioeconomic changes in China. *Proc Natl Acad Sci U S A* [Internet]. 2016;113(17):4640–5. Available from: <http://www.pnas.org/content/113/17/4640.full>
36. Zivin JG, Neidell MJ. Temperature and the Allocation of Time: Implications for Climate Change. 2010;32(1). Available from: <http://www.nber.org/papers/w15717>

Appendix Table 8. Papers selected using the third search strategy

Paper	Year	Country/Region	Population	Heat exposure index	Study design	Outcome	Main results
Afshan et al (1)	2014	Pakistan	General population	Mean environmental temp; Mean, min and max temp; Relative humidity; Mean evaporation; Mean sunshine hours; Mean solar radiation; Wet Day Index (Mt)	Ecological	Fascioliasis	Increased fascioliasis risk throughout the Punjab plain and its decrease in the northern highlands of the province became evident during the study period. The results point in the direction of evident trend modifications for the summer months
Alimi et al (2)	2015	northern South America (nSA)	Vectors	Temperature and precipitation	Simulation based on climate driven model	Malaria	While areas with malaria are expected to decrease, both vectors are predicted to experience range expansions in the future. Elevation, annual precipitation and temperature were influential in all models both current and future.
Anyamba et al (3)	2012	Eastern and Southern Africa	General population	Surface air temperature; precipitation anomalies	Ecological	Rift Valley fever	RVF outbreaks occurred after mo. of abnormal rainfall. Chikungunya outbreaks in conditions of anomalous heat and drought in Eastern Africa; in Southeast Asia, they were negatively correlated with drought, but positively correlated with higher temperatures and rainfall.

Atkinson et al (4)	2013	Global	General population	Not specified	Systematic review	Echinococcus granulosus and E. multilocularis	Causal links between specific environmental variables and echinococcosis in humans are yet to be established. Authors recommend for incorporation of environmental, animal and human data collection within a sentinel site surveillance network
Balbus et al (5)	2009	USA	General population, including working population	Not specified	Narrative review	Vector-borne diseases	Electricity and pipeline utility workers are at increased risk of infection with Lyme Disease, although evidence is lacking for greater risk of clinical illness.
Bennet et al (6)	2010	Global	Working population	Not specified	Narrative review	Infectious diseases	Incidence schistosomiasis, distribution of Aedes albopictus, prevalence of leishmaniasis, have increased. Outbreaks and endemic diarrhoeal disease are already a major source of lost work days, especially in developing countries
Bett et al (7)	2017	Global	General population	Not specified	Narrative review	Vector-borne diseases	Some of the studies are used to show a positive association between temperature and expansion of the geographical ranges of arthropod vectors, while others are used to illustrate an opposite trend.

Bostan et al (8)	2016	Pakistan	General population	Average temperature and rainfall trends	Ecological	Dengue fever	An immense potential for the sustenance of DF in major metropolitan cities in Pakistan for more than 9 months, with temperature at $\geq 18^{\circ}\text{C}$ (Punjab and Sindh Province, where maximum incidences have been reported over the years).
Caminade et al (9)	2015	Europe	Livestock	Maximum and minimum temperatures	Simulation based on climate driven model	Fascioliasis	The estimated pattern of climate change for northern Europe will possibly extend the season suitable for development of the parasite in the environment by up to four months. For southern Europe, the risk will increase during the winter months.
Chaves et al (10)	2010	East African highlands	Vectors	Not specified	Narrative review	Malaria	Authors reviewed the evidence for the role of climate as a driving force for highland invasion by malaria and concluded that it is robust. However, other aspects are involved (e.g. deforestation, predators, competitors) and should be taken into account.
Daniel et al (11)	2009	Czech Republic	Vectors	Ground-level temperature	Cross-sectional	Tick-borne Encephalitis Virus and <i>Borrelia burgdorferi</i> s.l.	A trend of increasing mean air temperatures during the last three decades was found. The trend was most pronounced in the spring and summer months with

							the highest activity of <i>I. ricinus</i> ticks.
Desai et al (12)	2009	Romania, Slovakia, Poland	Agriculture	Not specified	Retrospective cohort study	Leptospirosis	The unusually warm winter of 2006-2007 supported vole population growth and contributed to this resurgence of leptospirosis in Germany
Elbers et al (13)	2015	Global	Vectors	Not specified	Narrative review	Mosquitoes' geographic distribution	Over the last few decades, multiple Asian species have established themselves in Europe, spread and are unlikely to ever be eradicated. Ecological interactions with newly encountered species can also be modified by factors, such as photoperiodism.
Escobedo et al (14)	2015	Cuba	Other	the Oceanic Niño Index (ONI)	Ecological	Giardiasis	The cumulated rates for children were significantly higher, 244.73 cases/10,000 pop. (95%CI 238.5-251.0), compared to the workers, 101.82 cases/10,000 pop. (95%CI 93.5-110.2)
Feria-Arroyo et al (15)	2014	Texas and a portion of northeast Mexico	Vectors	Annual mean temperature, Max temp (warmest month); min temp (coldest month); temp annual range; mean temp driest, warmest and coldest	Simulation based on climate driven model	Lyme disease	The habitat suitable for the distribution of <i>I. scapularis</i> in the Texas-Mexico transboundary region will remain relatively stable until 2050.

				quarter + precipitation indexes			
Gale et al (16)	2009	Global	Livestock	Not specified	Narrative review	Vector-borne diseases	Warmer night-time temperatures influence many insect vectors that fly and feed at night. Rainfall affects mosquito abundance. Climate change could promote biological invasion of nonindigenous species with their associated pathogens and parasites.
Gale et al (17)	2010	Europe	Livestock	Not specified	Simulation based on climate driven model	Crimean- Congo haemorrhagic fever	Southern European bird species may expand their breeding ranges into more northern parts. The chances of moulting and survival of immature ticks may increase. An expansion of <i>H. marginatum</i> distribution northwards may be expected.
Gikungu et al (18)	2016	Kenya	Livestock	Mean maximum and minimum monthly temperature, relative humidity, sea surface temperatures	Simulation based on climate driven model	Rift valley fever	Besides rainfall, minimum temperature was found to be the most significant predictor in each of the models.
Holy et al (19)	2011	Germany	General population	Mean monthly air temperatures	Simulation based on climate driven model	Malaria	Both modelling approaches employed resulted in prolonged seasonal transmission gates in the

							future, enabling malaria transmissions up to 6 months in the climate reference period 2051-2080.
Ivanescu et al (20)	2016	Romania	General population	Mean monthly temperature	Simulation based on climate driven model	Malaria	Doing an extrapolation of the evolution of temperatures in 2030, a slight increase of temperatures by an average of 24°C in 2030 can be estimated, which may ensure a favourable climate for the development of Culicidae.
Jiménez-Clavero (21)	2012	Global	General population	Not specified	Narrative review	West Nile fever	West Nile virus (WNV) has expanded in the last 12 years, particularly in the Americas, where it first occurred in 1999, extending relentlessly since then. In Europe, is since the last years of the XX century that its incidence has risen substantially.
Joyner et al (22)	2010	Kazakhstan	Infectious agent	Emissions scenarios produced by the IPCC	Simulation based on climate driven model	Bacillus anthracis	Overall the predicted current distribution of B. anthracis stretches across the northern tier, eastern quarter, and south-eastern regions of Kazakhstan. It is predicted that these areas are potentially maintaining suitable environments for B. anthracis.

Konrad et al (23)	2012	USA	Infectious agent	Daily maximum and minimum temperatures	Simulation based on climate driven model	Rift Valley fever	Areas most at risk include the Baltimore and NYC metro areas, most of Texas, especially around Houston; Florida; Atlanta; southwest Nebraska; southern California and Arizona; and the central valley of California.
Lal et al (24)	2013	Global	General population	Several, depending on the selected studies	Narrative review	Cryptosporidiosis and giardiasis	Future environmental change is expected to have the greatest health impacts in countries with limited resources.
Lal et al (25)	2015	New Zealand	General population	Ambient temperature	Systematic review	Campylobacteriosis, salmonellosis, giardiasis and cryptosporidiosis.	Enteric disease risk from environmental reservoirs is pathogen specific. In some rural regions, environmental pathogen load is considerable, with multiple opportunities for zoonotic transmission.
Léger et al (26)	2013	Global	Vectors	Not specified	Narrative review	Ticks distribution	The northward expansion of <i>I. ricinus</i> has been well-documented in Scandinavia; <i>Ixodes uriae</i> , a tick associated with colonial seabirds, may be increasing in abundance and occurrence at high latitudes. Expansions have also been observed in African ticks.
Lindsay et al (27)	2001	Great Britain	Vectors	Temperature surfaces and future temperature change scenarios	Simulation based on	Vivax malaria	Under all climate-change scenarios, the risk of transmission increases in

					climate driven model		the south of England, spreading northwards to southern Scotland.
Moore et al (28)	2012	Africa	Humans, animals and vectors	Mean temperature	Simulation based on climate driven model	African trypanosomiasis	The model predicts that epidemics can occur when mean temperatures are between 20.7°C and 26.1°C. It predicts a large shift of up to 60% in the geographical extent of the range, and that 46-77 million additional people may be at risk of exposure by 2090.
Patz et al (29)	2008	Global	General population	Not specified	Narrative review	Infectious diseases	Although in isolated cases, disease resurgence has been attributed to recent warming trends, some of the long-term and complex problems posed by climate change may not be readily discernible from other causal factors.
Paz et al (30)	2007	Israel	Other	Mean, maximum and minimum temperatures	Ecological	Systemic V. vulnificus	Significant correlations between temperature values and hospital admission dates were revealed. Higher significant results were detected for the daily minimum temperatures in summer 1996 compatible with the disease eruption.

Sainz-Elipe et al (31)	2010	Spain	General population	Mean max. and min temperatures, mean environmental temperature, relative humidity, potential evapo transpiration, wind speed, vapour pressure, global radiation	Simulation based on climate driven model	Malaria	Climatological analyses and Gradient Model Risk index show that a transmission risk presently exists, lasting from May until September for <i>P. falciparum</i> , and from May until October for <i>P. vivax</i> .
Singh et al (32)	2011	India	Vectors	Not specified	Narrative review	Zoonoses	In recent years, vector-borne diseases have emerged as a serious public health issue in countries of the South- East Asia region. Many of these, particularly dengue fever and Japanese encephalitis, now occur in epidemic form, almost on an annual basis.
Tanga et al (33)	2010	Cameroon	Vectors	Monthly mean maximum and minimum temperatures	Cross-sectional	Vectors of malaria	There was a strong positive correlation between the abundance of the <i>Anopheles</i> species and the maximum, mean and minimum temperatures, whereas, for <i>An. hancocki</i> , the correlation was significant (positive) only for minimum temperature.
WHO (34)	2014	Global	General population	Mean temp (coldest and warmest month); mean annual temperature; mean precipitation (driest and wettest month); mean annual precipitation.	Simulation based on climate driven model	Malaria and dengue	Compared with a future without climate change, 60,000 additional deaths due to malaria are projected for the year 2030. There is very little projected increase

							in deaths due to dengue fever.
Xue et al (35)	2011	Ethiopia	General population	Monthly maximum air temperature, monthly average land surface temperature, monthly total precipitation and precipitation estimates	Ecological	Schistosoma mansoni	An apparent association between temperature and number of patients was observed: most peaks and valleys of the two time series appeared synchronized. This correlation was stronger for land surface temperature than for air temperature
Yang et al (36)	2017	China	General population, including working population	Annual average temperature (AAT), annual cumulative rainfall (ACR)	Case surveillance	Vectors of malaria	Annual average temperature, annual cumulative rainfall, rice yield per square kilometre and proportion of rural employees mainly showed a positive association with the malaria incidence rate.
Yusa et al (37)	2015	Canada	General population, including working population	Not specified	Systematic review	Infectious diseases	Exposure to vector-borne diseases, such as WNV, that may be linked to drought is increased in people spending greater amounts of time outdoors (e.g., agricultural workers, parks services)

References

1. Afshan K, Fortes-Lima CA, Artigas P, Valero AM, Qayyum M, Mas-Coma S. Impact of climate change and man-made irrigation systems on the transmission risk, long-term trend and seasonality of human and animal fascioliasis in Pakistan. *Geospat Health*. 2014;8(2):317–34.
2. Alimi TO, Fuller DO, Qualls WA, Herrera S V, Arevalo-Herrera M, Quinones ML, et al. Predicting potential ranges of primary malaria vectors and malaria in northern South America based on projected changes in climate, land cover and human population. *Parasit Vectors* [Internet]. *Parasites & Vectors*; 2015;8(1):431. Available from: <http://www.parasitesandvectors.com/content/8/1/431>
3. Anyamba A, Linthicum KJ, Small JL, Collins KM, Tucker CJ, Pak EW, et al. Climate teleconnections and recent patterns of human and animal disease outbreaks. *PLoS Negl Trop Dis*. 2012;6(1).
4. Atkinson JAM, Gray DJ, Clements ACA, Barnes TS, Mcmanus DP, Yang YR. Environmental changes impacting *Echinococcus* transmission: Research to support predictive surveillance and control. *Glob Chang Biol*. 2013;19(3):677–88.
5. Balbus John M, Malina Catherine M. Identifying Vulnerable Subpopulations for Climate Change Health Effects in the United States. *J Occup Environ Med*. 2009;51(1):33–7.
6. Bennett CM, McMichael AJ. Non-heat related impacts of climate change on working populations. *Glob Health Action* [Internet]. 2010 Dec 17;3. Available from: <http://journals.sfu.ca/coaction/index.php/gha/article/view/5640>
7. Bett B, Kiunga P, Gachohi J, Sindato C, Mbotha D, Robinson T, et al. Effects of climate change on the occurrence and distribution of livestock diseases. *Prev Vet Med* [Internet]. Elsevier B.V.; 2017;137(November 2015):119–29. Available from: <http://dx.doi.org/10.1016/j.prevetmed.2016.11.019>
8. Bostan N, Javed S, Nabgha-e-Amen, Eqani SAMAS, Tahir F, Bokhari H. Dengue fever virus in Pakistan: effects of seasonal pattern and temperature change on distribution of vector and virus. *Rev Med Virol*. 2017;27(1):1–17.
9. Caminade C, van Dijk J, Baylis M, Williams D. Modelling recent and future climatic suitability for fasciolosis in Europe. *Geospat Health*. 2015;9(2):301–8.
10. Chaves LF, Koenraadt CJM. Climate change and highland malaria: fresh air for a hot debate. *Q Rev Biol* [Internet]. 2010 Mar;85(1):27–55. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20337259>
11. Daniel M, Materna J, Honig V, Metelka L, Danielová V, Harcarik J, et al. Vertical distribution of the tick *Ixodes ricinus* and tick-borne pathogens in the northern Moravian mountains correlated with climate warming (Jeseníky Mts., Czech Republic). *Cent Eur J Public Health* [Internet]. 2009 Sep;17(3):139–45. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20020603>
12. Desai S, van Treeck U, Lierz M, Espelage W, Zota L, Sarbu A, et al. Resurgence of field fever in a temperate country: an epidemic of leptospirosis among seasonal strawberry harvesters in Germany in 2007. *Clin Infect Dis*. 2009;48(6):691–7.
13. ELBERS ARW, KOENRAADT C, MEISWINKEL R. Mosquitoes and *Culicoides* biting midges: vector range and the influence of climate change. *Rev Sci Tech l'OIE* [Internet]. 2015 Apr 1;34(1):123–37. Available from: <http://doc.oie.int:8080/dyn/portal/index.seam?page=alo&alold=32331>
14. Escobedo AA, Almirall P, Rumbaut R, Rodríguez-Morales AJ. Potential impact of macroclimatic variability on the epidemiology of giardiasis in three provinces of Cuba, 2010-2012. *J Infect Public Health* [Internet]. King Saud Bin Abdulaziz University for Health Sciences; 2015;8(1):80–9. Available from: <http://dx.doi.org/10.1016/j.jiph.2014.06.001>

15. Feria Arroyo TP, Castro-Arellano I, Gordillo-Peréz G, Cavazos AL, Vargas-Sandoval M, Grover A, et al. Implications of climate change on the distribution of the tick vector *Ixodes scapularis* and risk for Lyme disease in the Texas-Mexico transboundary region. *Parasit Vectors* [Internet]. 2014;7(1):199. Available from: <http://parasitesandvectors.biomedcentral.com/articles/10.1186/1756-3305-7-199>
16. Gale P, Drew T, Phipps LP, David G, Wooldridge M. The effect of climate change on the occurrence and prevalence of livestock diseases in Great Britain. *J Appl Microbiol.* 2009;106(5):1409–23.
17. Gale P, Estrada-Peña A, Martínez M, Ulrich RG, Wilson A, Capelli G, et al. The feasibility of developing a risk assessment for the impact of climate change on the emergence of Crimean-Congo haemorrhagic fever in livestock in Europe: A Review. *J Appl Microbiol.* 2010;108(6):1859–70.
18. Gikungu D, Wakhungu J, Siamba D, Neyole E, Muita R, Bett B. Dynamic risk model for Rift Valley fever outbreaks in Kenya based on climate and disease outbreak data. *Geospat Health.* 2016;11(2):95–103.
19. Holy M, Schmidt G, Schröder W. Potential malaria outbreak in Germany due to climate warming: Risk modelling based on temperature measurements and regional climate models. *Environ Sci Pollut Res.* 2011;18(3):428–35.
20. Ivanescu L, Bodale I, Florescu SA, Roman C, Acatrinei D, Miron L. Climate Change Is Increasing the Risk of the Reemergence of Malaria in Romania. *Biomed Res Int.* 2016;2016.
21. Jiménez-Clavero MA. Animal viral diseases and global change: Bluetongue and West Nile fever as paradigms. *Front Genet.* 2012;3(JUN):1–15.
22. Joyner TA, Lukhnova L, Pazilov Y, Temiralyeva G, Hugh-Jones ME, Aikimbayev A, et al. Modeling the potential distribution of *Bacillus anthracis* under multiple climate change scenarios for Kazakhstan. *PLoS One.* 2010;5(3).
23. Konrad SK, Miller SN. A temperature-limited assessment of the risk of Rift Valley fever transmission and establishment in the continental United States of America. *Geospat Health.* 2012;6(2):161–70.
24. Lal A, Baker MG, Hales S, French NP. Potential effects of global environmental changes on cryptosporidiosis and giardiasis transmission. *Trends Parasitol* [Internet]. Elsevier Ltd; 2013;29(2):83–90. Available from: <http://dx.doi.org/10.1016/j.pt.2012.10.005>
25. Lal A, Lill AWT, McIntyre M, Hales S, Baker MG, French NP. Environmental change and enteric zoonoses in New Zealand: A systematic review of the evidence. *Aust N Z J Public Health.* 2015;39(1):63–8.
26. Léger E, Vourc'h G, Vial L, Chevillon C, McCoy KD. Changing distributions of ticks: Causes and consequences. *Exp Appl Acarol.* 2013;59(1–2):219–44.
27. Lindsay SW, Thomas CJ. Global warming and risk of vivax malaria in Great Britain. *Glob Chang Hum Heal.* 2001;2(1):80–4.
28. Moore S, Shrestha S, Tomlinson KW, Vuong H, Colwell RR, Pascual M, et al. Predicting the effect of climate change on African trypanosomiasis: integrating epidemiology with parasite and vector biology. *J R Soc Interface* [Internet]. 2012;9(70):817–30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22072451>
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3306657>
29. Patz JA, Olson SH, Uejio CK, Gibbs HK. Disease Emergence from Global Climate and Land Use Change. *Med Clin North Am.* 2008;92(6):1473–91.
30. Paz S, Bisharat N, Paz E, Kidar O, Cohen D. Climate change and the emergence of *Vibrio vulnificus* disease in Israel. *Environ Res.* 2007;103(3):390–6.
31. Sainz-Elipe S, Latorre JM, Escosa R, Masia M, Fuentes M V, Mas-Coma S, et al. Malaria resurgence risk in southern Europe: climate assessment in a historically endemic area of rice fields at the Mediterranean shore of Spain. *Malar J* [Internet]. 2010;9:221. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20673367>
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2924348/pdf/1475-2875-9-221.pdf>

32. Singh BB, Sharma R, Gill JPS, Aulakh RS, Banga HS. Climate change, zoonoses and India. *Rev Sci Tech* [Internet]. 2011;30(3):779–88. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22435190>
33. Tanga MC, Ngundu WI, Judith N, Mbuh J, Tendongfor N, Simard F, et al. Climate change and altitudinal structuring of malaria vectors in south-western Cameroon: Their relation to malaria transmission. *Trans R Soc Trop Med Hyg*. 2010;104(7):453–60.
34. World Health Organization. Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s [Internet]. Hales S, Kovats S, Lloyd S, Campbell-Lendrum D, editors. WHO Library Cataloguing-in-Publication Data. Geneva, Switzerland; 2014. Available from: www.who.int/about/licensing/copyright_form/en/index.html
35. Xue Z, Gebremichael M, Ahmad R, Weldu ML, Bagtzoglou AC. Impact of temperature and precipitation on propagation of intestinal schistosomiasis in an irrigated region in Ethiopia: Suitability of satellite datasets. *Trop Med Int Heal*. 2011;16(9):1104–11.
36. Yang D, Xu C, Wang J, Zhao Y. Spatiotemporal epidemic characteristics and risk factor analysis of malaria in Yunnan Province, China. *BMC Public Health* [Internet]. *BMC Public Health*; 2017;17(1):66. Available from: <http://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-016-3994-9>
37. Yusa A, Berry P, Cheng JJ, Ogden N, Bonsal B, Stewart R, et al. Climate change, drought and human health in Canada. *Int J Environ Res Public Health*. 2015;12(7):8359–412.