

Nutritional status, Vitamin D and glucose metabolism: relatives or colleagues?

Maurizio Mennini, Francesco Pallotti

“Sapienza” University of Rome, Italy

Dear Editor,

the general recommendation that breastfed infants require vitamin D (VitD) supplementations is widely accepted and many studies stress the need to start supplementation of 400 UI/d from birth (1), in order to maintain bone health in children as well in adults.

However, VitD deficiency is also called into question in many pathological inflammatory conditions and a possible influence on the nutritional status and on glucose metabolism seems also relevant. Esteghamati A et al. demonstrated that Serum 25-hydroxy vitamin D levels are lower in metabolically unhealthy obesity (MUO) compared to metabolically healthy obesity (MHO), and that reduced vitamin D concentrations are more strongly associated with cardio-metabolic and inflammatory markers in MUO than in MHO subjects (2).

We believe it is useful to recall the inverse relationship between parathyroid hormone (PTH) and VitD in children obesity: serum 25(OH)D levels are lower in obese, likely due to sequestration of VitD in the adipose tissue, in association with a PTH increase, thus eliciting risk of fracture (3).

Furthermore, VitD deficiency may be linked to the development of obesity-associated metabolic complications such as type 2 diabetes and metabolic syndrome, possibly by altering insulin secretion and reducing insulin sensitivity. It may be worthy to remember that effects on lipid profile and cardiovascular diseases are also noticeable. Many observational studies show increase in serum HDL cholesterol and reduction in total cholesterol, LDL and triglycerides in groups with higher serum 25(OH)D, which may be due to reduction of intestinal absorption and of lipolysis.

Kelishadi R et al. demonstrated such positive association in a pediatric age population in a recent metanalysis (4).

Kestenbaum B et al. (5) showed an increased incidence of myocardial infarction and cardiovascular mortality in adult patients with 25(OH)D deficiency (about 30% increased risk of mortality with VitD levels < 15ng/ml). In light of these findings, we want to promote in the field of pediatric endocrinology the need to recognize, assess and treat VitD deficiency since childhood, in order to reduce the risk of metabolic disorders as well as the risk of their related comorbidities.

References

1. Ziegler EE, Nelson SE, Jeter JM. Vitamin D supplementation of breastfed infants: A randomized dose-response trial. *Pediatr Res.* 2014 May 23.
2. Esteghamati A1, Aryan Z2, Esteghamati A1, Nakhjavani M3. Differences in vitamin D concentration between metabolically healthy and unhealthy obese adults: Associations with inflammatory and cardiometabolic markers in 4391 subjects. *Diabetes Metab.* 2014 May 5. pii: S1262-3636(14)00046-9.
3. Amini Z, Bryant S, Smith C, et al. Is the serum vitamin D-parathyroid hormone relationship influenced by obesity in children? *Horm Res Paediatr.* 2013;80(4):252-6.
4. Kelishadi R, Farajzadegan Z, Bahreynian M. Association between vitamin D status and lipid profile in children and adolescents: a systematic review and meta-analysis. *Int J Food Sci Nutr.* 2014 Jun;65:404-10.
5. Kestenbaum B, Katz R, de Boer I, Hoofnagle A, Sarnak MJ, Shlipak MG, Jenny NS, Siscovick DS. Vitamin D, parathyroid hormone, and cardiovascular events among older adults. *J Am Coll Cardiol.* 2011 Sep 27;58:1433-41.

Correspondence:

Maurizio Mennini, M.D.

“Sapienza” University of Rome – Italy

Viale Regina Elena, 324

00161 Rome Italy

E-mail: maurizio.mennini@gmail.com