

Effect of diverse low energy-dense versus healthy diet on metabolic outcomes in overweight/obese adolescents: a randomized controlled trial

*Niloofar Aghaei*¹, *Mohammad Hossein Rouhani*¹, *Fatemeh Tabatabaei*², *Bagher Larijani*³, *Leila Azadbakht*^{1,5,3}

¹Food Security Research Center and Department of Community Nutrition, School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan, Iran; ²Isfahan Endocrine and Metabolism Research Center, Isfahan University of Medical Sciences, Isfahan, Iran; ³Endocrinology and Metabolism Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran; ⁵Diabetes Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran, Email: azadbakht@hlth.mui.ac.ir

Summary. *Background:* Dyslipidemia and impaired glucose metabolism are prevalent among overweight/obese adolescents. Although several dietary interventions used to reduce these complications, no study has been conducted regarding the effect of a diverse low energy-dense diet (DLED) on metabolic outcomes. *Objective:* To assess the effect of diverse low energy-dense diet on anthropometric variables, lipid profile and fasting blood sugar in overweight/obese adolescents aged 8-18 years. *Methods:* For this parallel clinical trial, 40 overweight/obese adolescents were randomly assigned to a DLED or a healthy recommendation based diet for 10 weeks. Physical activity and dietary intakes were recorded during the study. Total energy expenditure was estimated by Harris Benedict equations for each subject. To reduce energy density, a low-fat (25% fat, 15% protein and 60% carbohydrate) diet rich in dietary fiber and water was prescribed. The dietary variety was increased by recommending adolescents to consume at least one-half of subgroups of each main food group daily. Healthy nutritional recommendations were focused on limiting fast foods, fatty foods, added sugar, sugar sweetened beverages, French fries and unhealthy fat intake. Also, these recommendations encourage adolescents to consume more fruits and vegetables, whole grains, low-fat dairy and water. Total cholesterol, low density lipoprotein, high density lipoprotein, triglyceride, fasting blood sugar and anthropometric variables were assessed at baseline and end-point. *Results:* The data of 40 adolescents were analyzed. Within groups analysis showed that there was a significant reduction in triglyceride ($p=0.04$) and low density lipoprotein ($p<0.01$) in DLED group. We could not find any significant difference in baseline relative to end-point measurements in both groups. Changes of anthropometric and biochemical variables were compared in a crude and adjusted models. We could not find any significant changes in anthropometric and biochemical variables between two diet groups. *Conclusion:* This study showed that compliance to a DLED diet may result in a significant decrease in TG and LDL after 10 weeks. However, we did not observe any significant difference in changes of variables between members of DLED group and those who consumed a healthy diet.

Keywords: dietary diversity, dietary energy density, lipid profile, adolescents

Introduction

During the recent decades, childhood obesity becomes a global concern (1). Obese children have greater risk for hypercholesterolemia and hypertension in adulthood (2). Also, elevated concentration of C-reactive protein (CRP) has been reported in obese children (3). The prevalence of obesity among American 2- to 19-year-olds children was 16.9% in 2011-2012 (4). In Iran, the range of the prevalence of obesity among children was estimated 3.2-11.9% (5). Dietary intervention is an effective choice for prevention of obesity complications. As low calorie diets are not recommended by some guidelines for individuals with childhood obesity (1), investigators focused on the other aspects of nutritional management such as manipulating dietary energy density (DED) and increase dietary diversity. DED is an index of diet quality which shows the ratio of energy content to gram of diet (6). It has an important role in nutritional intervention against obesity (7). Evidence showed a direct association between DED and obesity in different populations (6, 8). Also, it was observed that DED was related to dyslipidemia and cardiometabolic risk factors (9). Although some clinical trial studies have focused on the effects of low energy-dense diets on obesity in adults (10) and children (11), data regarding the effects of a low energy-dense diet on complications of obesity (e.g., hypertension, dyslipidemia and increased level of inflammatory markers) are scarce, especially among children.

Dietary diversity is another index of diet quality. Although the results of observational studies are not consistent (12), most studies have reported an indirect association between obesity and dietary variety (13, 14). Indeed, when diversity was in selective (not all) food groups, favorable association was observed (12). Moreover, studies reported that dietary diversity was inversely related to risk of hypertension, metabolic syndrome and cardiometabolic risk factors in adults (14, 15). However, we did not find any evidence regarding adolescents. Also, there is no clinical trial in which the effects of a diverse diet on metabolic markers have been assessed. Moreover, clinical trials which manipulated DED, overlooked dietary variety in their intervention and a diverse low energy-dense diet was

not prescribed in previous studies. Therefore, the aim of present study was to examine the effect of a diverse low energy-dense diet on complications of obesity among obese adolescents.

Methods

Subjects

From subjects referred to pediatric clinics (Nour and Ali Asghar Hospital Isfahan, Endocrinology and Metabolism Research Center, Isfahan Sedigheye Tahere) 40 adolescents (aged 10-18 years) with excess body weight were recruited for the current randomized parallel clinical study from subjects referred to the pediatric clinics. The individuals of BMI were between body mass index (BMI) 85th to 95th percentiles of BMI for age in accordance with guidelines by WHO (16). All participants signed informed written consent form. Individuals with poor compliance were excluded. Present study was conducted in 2015 in Isfahan, Iran. Data regarding general characteristics of subjects were collected by a questionnaire. We estimated sample size by following formula (type one and type two error were 0.05 and 0.20, respectively): $N=2[(Z_{1-\alpha/2}+Z_{1-\beta})^2 \times S^2]/d^2$ where S (variance of waist circumference) was 5.49 and d was 55 cm (17). According to this formula, 16 adolescents in each group were sufficient for present study. The study groups were matched for age of participants. Research Council and Ethical Committee of School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan, Iran and Food Security Research Center, Isfahan University of Medical Sciences, Isfahan, Iran approved this study. The Registration code of current clinical trial in IRCT was IRCT201109272839N4.

Study procedure

Eligible subjects were randomly assigned to consume a diverse low energy-dense diet or a diet based on healthy nutritional recommendations for 10 weeks. We used random number table for randomization. As this was a dietary intervention, we could not blind adolescents to the type of study intervention. Total energy expenditure (based on the age, sex and physical activity) was estimated by Harris Benedict equations

for each subject. According to the intervention groups, an individualized diet was prescribed to each subject. Exchange list for meal planning was taught to the all parents and adolescents. The macronutrient composition of prescribed diet in the healthy nutritional recommendations group was 35% of energy from fat, 15% of energy from protein and 50% energy from carbohydrate. To reduce DED, we prescribed a low-fat (25% of energy from fat, 15% of energy from protein and 60% energy from carbohydrate) diet rich in dietary fiber and water in diverse low energy-dense diet group. Also, a list of five main food groups (grains, dairies, fruits, vegetables and meats) and their subgroups was provided to members of this group. The dietary variety was increased by recommending adolescents to consume at least one-half of subgroups of each main food group daily. Parents were asked to contribute to the intervention. Healthy nutritional recommendations were focused on limiting fast foods, fatty foods, added sugar, sugar sweetened beverages, French fries and unhealthy fat intake. Also, these recommendations encourage adolescents to consume more fruits and vegetables, whole grains, low-fat dairies and water. Periodical individual visits (one visit per each 2 weeks) were conducted. All subjects were asked to fill in a one-day food diary and a one-day physical activity record and deliver them in each visit. Mean physical activity time multiplied by metabolic equivalent (MET) to calculate physical activity level.

Anthropometric measurements

Anthropometric variables were assessed at baseline and after 10 weeks. Body weight was measured to the nearest 0.1 kg while participants wore light clothes. We measured height to the nearest 1 cm while adolescents were without shoes and stand with his/her back against the wall. To calculate BMI, we divided weight (kg) by height squared (m^2). Waist circumference (WC) was measured at the line below the last rib by using standard type to the nearest 1 cm (18).

Biochemical measurements

Biochemical variables were measured at baseline and after 10 weeks of intervention. Blood specimens were collected after 12 hours overnight fasting. After coagulating and centrifuging at $3,000\times g$ for 10 min,

serum was separated. We run Enzymatic colorimetric tests to measure the concentration of triglyceride (TG) and total cholesterol (TC). After blocking low density lipoprotein (LDL), very low density lipoprotein and chylomicrons, the level of high density lipoprotein (HDL) was enzymatically measured. We blocked HDL, VLDL and chylomicrons by enzymatic colorimetric tests and then LDL concentration was assessed. To measure fasting blood sugar (FBS), a glucose oxidase method was used. All biochemical variables were measured by an inter- and intra-assay coefficients lower than 5%. We used biochemical kits provided by Pars Azmoon Inc (Tehran, Iran) for the current study. The measurement was repeated at 10th week of the trial.

Other variables

Education level of subjects was orally asked. Physical activity records analyzed and converted to MET/h (19).

Statistical Analysis

To test the normal distribution of variables, we run Kolmogorov–Smirnov test and drew a histogram curve. We used intention-to-treat analysis according to the last-observation-carried-forward (LOCF) method. Student t test was run to compare changes, baseline and endpoint measurements between two groups. Within groups differences between baseline and endpoint values was tested by paired t test. Change of each variable was calculated by this formula: “(endpoint value-baseline value)/baseline value”. We adjusted the effects of potential confounders by run analysis of the covariance (ANCOVA). All values were presented as means \pm SD. SPSS 20.0 (SPSS Inc) was used for statistical analysis.

Results

As illustrated in Figure 1, we enrolled a total of 40 volunteers and 38 subjects completed the study (from Jan 2013 to Aug 2015). Characteristics of participated adolescents at baseline in the table healthy diet are shown in Table 1. There was no significant difference in age ($P=0.53$), sex ($P=0.49$) ratio, height ($P=0.77$), weight ($P=0.42$), body mass index percen-




Figure 1. Study procedure

tile ($P=0.16$) and physical activity ($P=0.19$) between the two groups.

Dietary intake during study in two groups is shown in Table 2. There was no significant difference in intake of protein, carbohydrate, fat and several mineral and vitamins between the groups.

Baseline and endpoint measures of anthropometric and biochemical variables in adolescents are presented in Table 3. Between the groups analysis showed that subjects in the diverse low energy-dense group had higher level of TG at baseline in compared with healthy diet (100.78 ± 8.88 vs. 135.78 ± 10.80 mg/dl; $P=0.02$). Similar result was observed for end-point values of TG (90.94 ± 7.47 vs. 117.22 ± 9.84 mg/dl; $P=0.04$). between the groups analysis showed that there was a significant reduction in TG in diverse low energy-dense group

(135.78 ± 10.80 vs. 117.22 ± 9.84 mg/dl; $P=0.04$). A significant reduction was also observed for LDL in diverse low energy-dense group after 10 weeks of intervention (106.00 ± 4.90 vs. 89.56 ± 3.68 mg/dl; $P<0.01$). We could not find any significant results for other variables in both within and between groups analysis. Changes of anthropometric and biochemical variables in adolescents are displayed in Table 4. Results are presented in crude and adjusted models (adjusted for age, sex and education). We could not find any significant changes in anthropometric and biochemical variables between the two diet groups.

Discussion

The findings of the present study revealed that compliance to a diverse low energy-dense diet may result in a remarkable reduction in TG and LDL after 10 weeks. However, we did not observe any significant difference in changes of other variables (i.e., TC, HDL, FBS) between two diet groups. To the best of our knowledge, this is the first randomized clinical trial in which a diverse low energy-dense diet was prescribed.

Our previous observational studies conducted in Iran showed that there was a favorable association between dietary diversity score (DDS) and health outcomes in adults (13, 14, 20). Similar findings were reported from US, a developed country (21). These results have not been confirmed by a comprehensive

Table 1. Characteristics of participated adolescents at baseline in the two groups groups.

Variables	Healthy diet (n=20)	Diverse low energy-dense diet (n=20)	P value ¹
Age (yr)	10.97 \pm 2.54	11.44 \pm 2.10	0.53
Female (%)	66.7	10(55.6)	0.49
Height (cm)	147.86 \pm 16	146.46 \pm 13.63	0.77
Weight (Kg)	11(56.72 \pm 18.95%)	11(52.53 \pm 13%)	0.42
BMI percentile \geq 95	16(77.8%)	11(55.6%)	0.16
Physical activity (MET-h/wk)	1861.74 \pm 1097.30	1497.44 \pm 506.05	0.19

¹P for continuous variables generated from independent student t test and for categorical variables by chi-square test. As the distribution of physical activity was not normal, we used Mann-Whitney test for this variable.

Table 2. Reported dietary intake of energy and nutrients of participants.

Variables	Healthy diet (n=20)	Diverse low energy-dense diet (n=20)	P value ¹
Energy (kcal)	1225.39±1136.93 ²	930.35±773.79	0.34
Carbohydrate (g)	82.67±61.77	121.25±61.77	0.05
Protein (g)	50.87±22.10	50.00±22.11	
Fat (g)	61.59±26.33	45.47±26.35	0.06
Fiber (g)	7.50±6.19	9.51±6.20	0.31
Zinc (mg)	7.09±3.42	7.33±3.44	0.83
Calcium (mg)	856.19±900.37	999.61±900.37	0.62
Vitamin B ₃ (mg)	7.54±3.00	9.23±2.88	0.07
Folate (mcg)	172.78±74.86	176.77±74.58	0.87
Vitamin C (mg)	56.22±27.52	57.74±27.53	0.86
Potassium (mg)	1201.76±1351.40	1955.05±1352.85	0.09
Magnesium (mg)	133.03±83.59	177.86±83.54	0.10
Energy density (kcal/g)	1.31±0.35	1.19±0.35	0.22

¹ Independent T test (for energy and energy density) and Analysis of covariance were used for calculating P value.

² Variables are presented as mean±SD. All values were adjusted for total energy intake except for energy and energy density

systematic review and meta-analysis which reported that there was no significant relationship between DDS and BMI (22). Observational studies conducted in Iranian adolescents showed that there was no significant association between BMI, WC and DDS (23). Therefore, the findings regarding the association between DDS and health outcomes are controversial. It is possible that the duration of our study (10 weeks) may not be sufficient for significant change in anthropometric variables. Moreover, we compared two healthy diets (a diverse low energy-dense vs. a diet based on the healthy nutritional recommendations) and therefore, we did not find any significant changes between these two diets.

You conclude the studies related to DDS but your study did not include this score

The results of the current study showed a remarkable reduction in TG after 10 weeks of consuming a diverse low energy-dense diet. We could not find any clinical trial regarding the effect of dietary diversity on metabolic outcomes in adolescents, however, an observational study conducted in Iranian women revealed a

direct association between DED and serum concentration of TG (23). Another Iranian study reported that compliance to a high energy-dense diet may be associated with elevated LDL concentration (9).

In this study, we observed a significant reduction in TG and LDL after 10 weeks of consuming a diverse low energy-dense diet. A low energy-dense diet is limited in refined grains, added sugar, added fat, saturated fatty acids and *trans* fatty acids which may result in insulin resistance and metabolic abnormalities (24).

DED was introduced as a marker of diet quality in adolescents (25). The results regarding the association between DED and obesity in adolescents are similar. Although a systematic review suggested lowering DED as a strategy of obesity management in children and adolescents (26), O'Sullivan et al. (27) reported a non-significant association between DED and obesity in adolescents. Also, Kring et al. (28) did not find any significant relation between DED and subsequent change in BMI among children.

We could not find a clinical trial in which a diverse low energy-dense diet was prescribed and there-

Table 3. Baseline and endpoint measurements of anthropometric and biochemical variables in the two groups

Variables	Healthy diet (n=20)	Diverse low energy-dense diet (n=20)	P value ²
Weight (Kg)			
Baseline	56.74±18.86 ³	52.91±13.67	0.42
End	56.67±18.86	53.46±14.25	0.55
P value ¹	0.96	0.54	
Body mass index (kg/m²)			
Baseline	25.41±3.39	24.25±2.23	0.23
End	24.93±3.66	24.11±2.54	0.42
P value ¹	0.71		
Waist circumference (cm)			
Baseline	87.36±11.04	86.67±8.62	0.83
End	86.46±9.34	85.94±10.05	0.87
P value ¹	0.38		
Triglyceride (mg/dl)			
Baseline	100.78±39.69	135.78±48.28	0.02
End	90.94±33.39	117.22±43.98	0.04
P value ¹	0.40	0.04	
Total cholesterol (mg/dl)			
Baseline	198.33±32.58	177.56±31.78	0.45
End	162.61±34.77	171.61±47.47	0.46
P value ¹	0.38	0.52	
LDL (mg/dl)			
Baseline	95.28±19.98	106.00±21.90	0.11
End	88.31±19.35	89.56±16.45	0.83
P value ¹	0.02	<0.01	
HDL (mg/dl)			
Baseline	47.00±10.14	42.50±4.87	0.08
End	46.50±12.43	43.22±5.40	0.29
P value ¹	0.69	0.56	
FBS (mg/dl)			
Baseline	89.44±6.80	87.94±7.51	0.51
End	90.28±4.83	87.56±10.77	0.37
P value ¹	0.66	0.82	

FBS: fasting blood sugar, HDL: high density lipoprotein, LDL: low density lipoprotein

¹P values are resulted from paired t test to compare values within each group.

²P values are resulted from independent t test to compare values between groups

³Mean±SD

fore, we could not compare our findings with an absolutely similar study.

There are several potential reasons for different results obtained from researches: 1) There are several

Table 4. Changes of anthropometric and biochemical variables in adolescents.

Variables	Healthy diet (n=20)	Diverse low energy-dense diet (n=20)	P value ³
Weight (Kg)			
Crude	0.55±0.51 ¹	-0.05±0.23	0.65
Model I ²	-0.23±1.04	0.69±1.00	0.56
Body mass index (kg/m²)			
Crude	-0.14±0.24	-0.48±0.28	0.53
Model I ²	-0.45±0.41	-0.17±0.41	0.66
Waist circumference (cm)			
Crude	-0.72±0.79	-0.90±1.35	0.89
Model I ²	-1.47±1.18	-0.15±1.18	0.48
Triglyceride (mg/dl)			
Crude	18.55±8.27	-9.83±11.28	0.54
Model I ²	-9.90±8.94	-18.48±8.94	0.53
Total cholesterol (mg/dl)			
Crude	-5.94±10.43	-5.72±6.63	0.98
Model I ²	-5.54±8.83	-5.54±8.83	0.82
LDL (mg/dl)			
Crude	16.44±4.27	-6.94±2.82	0.07
Model I ²	-13.78±3.40	-9.63±3.40	0.42
HDL (mg/dl)			
Crude	0.72±1.22	-0.50±1.25	0.49
Model I ²	-0.75±1.33	0.53±1.33	0.52
FBS (mg/dl)			
Crude	-0.39±1.73	0.83±1.87	0.63
Model I ²	-0.86±1.89	1.31±1.89	0.44

FBS: fasting blood sugar, HDL: high density lipoprotein, LDL: low density lipoprotein

¹ All values are mean change ± SD

² Adjusted for sex, age, physical activity, education and baseline values

³ Calculated by ANCOVA

methods to assess dietary diversity (29). Therefore, these different methods may be responsible for a part of observed heterogeneity. 2) As dietary diversity, DED is measured by different methods through the studies (30). 3) Low compliance to prescribed diets is prevalent among adolescents (31, 32). This point may affect clinical trial conducted in adolescent. 4) The baseline values of biochemical variables were in the normal range in our study and several other researches. Therefore, we did not assess the effect of a diverse low energy-dense diet on subjects with unhealthy metabolic status.

In conclusion, this study showed that compliance to a diverse low energy-dense diet may result in a significant decrease in TG and LDL after 10 weeks. However, we did not observe any significant difference in changes of variables between members of diverse low energy-dense group and those who consumed a healthy diet.

This study had several limitations. We could not assess the compliance of subjects to the prescribed diets by a biochemical variable because there is no specific biochemical marker for dietary variety and DED. This is a short-term clinical trial and more long-term interventions should be designed in future.

There are several strength points in the current study. Although more previous studies which assessed dietary diversity had observational design, this was the first interventional study focused on dietary diversity in adolescents. This study was conducted in overweight/obese adolescents. In comparison with the studies conducted in adults, the number of researches which have focused on adolescents is limited.

Acknowledgment

Food Security Research Center and Department of Community Nutrition, School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan, Iran has granted this study.

Bibliografia

- Han JC, Lawlor DA, Kimm SY. Childhood obesity. *Lancet*. 2010 May 15;375(9727):1737-48. PubMed PMID: 20451244. Pubmed Central PMCID: 3073855.
- Freedman DS, Mei Z, Srinivasan SR, Berenson GS, Dietz WH. Cardiovascular risk factors and excess adiposity among overweight children and adolescents: the Bogalusa Heart Study. *J Pediatr*. 2007 Jan;150(1):12-7 e2. PubMed PMID: 17188605.
- Sacheck J. Pediatric obesity: an inflammatory condition? *JPEN J Parenter Enteral Nutr*. 2008 Nov-Dec;32(6):633-7. PubMed PMID: 18974243.
- Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011-2012. *JAMA*. 2014 Feb 26;311(8):806-14. PubMed PMID: 24570244.
- Jafari-Adli S, Jouyandeh Z, Qorbani M, Soroush A, Larijani B, Hasani-Ranjbar S. Prevalence of obesity and overweight in adults and children in Iran; a systematic review. *J Diabetes Metab Disord*. 2014;13(1):121. PubMed PMID: 25610814. Pubmed Central PMCID: 4301060.
- Murakami K, Sasaki S, Takahashi Y, Uenishi K, Japan Dietetic Students' Study for N, Biomarkers G. Dietary energy density is associated with body mass index and waist circumference, but not with other metabolic risk factors, in free-living young Japanese women. *Nutrition*. 2007 Nov-Dec;23(11-12):798-806. PubMed PMID: 17936194.
- Wilks DC, Mander AP, Jebb SA, Thompson SG, Sharp SJ, Turner RM, et al. Dietary energy density and adiposity: employing bias adjustments in a meta-analysis of prospective studies. *BMC public health*. 2011;11:48. PubMed PMID: 21255448. Pubmed Central PMCID: 3038903.
- Ledikwe JH, Blanck HM, Kettel Khan L, Serdula MK, Seymour JD, Tohill BC, et al. Dietary energy density is associated with energy intake and weight status in US adults. *Am J Clin Nutr*. 2006 Jun;83(6):1362-8. PubMed PMID: 16762948.
- Esmailzadeh A, Boroujeni HK, Azadbakht L. Consumption of energy-dense diets in relation to cardiometabolic abnormalities among Iranian women. *Public Health Nutr*. 2012 May;15(5):868-75. PubMed PMID: 22008550.
- Chang UJ, Hong YH, Suh HJ, Jung EY. Lowering the energy density of parboiled rice by adding water-rich vegetables can decrease total energy intake in a parboiled rice-based diet without reducing satiety on healthy women. *Appetite*. 2010 Oct;55(2):338-42. PubMed PMID: 20654665.
- Epstein LH, Paluch RA, Beecher MD, Roemmich JN. Increasing healthy eating vs. reducing high energy-dense foods to treat pediatric obesity. *Obesity*. 2008 Feb;16(2):318-26. PubMed PMID: 18239639. Pubmed Central PMCID: 2408744.
- Jayawardena R, Byrne NM, Soares MJ, Katulanda P, Yaddav B, Hills AP. High dietary diversity is associated with obesity in Sri Lankan adults: an evaluation of three dietary scores. *BMC Public Health*. 2013;13:314. PubMed PMID: 23566236. Pubmed Central PMCID: 3626879.
- Azadbakht L, Esmailzadeh A. Dietary diversity score is related to obesity and abdominal adiposity among Iranian female youth. *Public Health Nutr*. 2011 Jan;14(1):62-9. PubMed PMID: 20353617.
- Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi F. Dietary diversity score and cardiovascular risk factors in Tehranian adults. *Public Health Nutr*. 2006 Sep;9(6):728-36. PubMed PMID: 16925878.
- Vadiveloo M, Parkeh N, Mattei J. Greater healthful food variety as measured by the US Healthy Food Diversity index is associated with lower odds of metabolic syndrome and its components in US adults. *J Nutr*. 2015 Mar;145(3):564-71. PubMed PMID: 25733473. Pubmed Central PMCID: 4336534.
- Organization WH. BMI-for-age GIRLS/BOYS. Available from: http://www.who.int/entity/growthref/bmifa_girls_5_19years_per.pdf.
- Fleiss JL. *The Design and Analysis of Clinical Experiments*.

- London: John Wiley and Sons; 1986. p.263-271.
18. B. E. Ainsworth, W. L. Haskell, M. C. Whitt et al., "Compendium of physical activities: an update of activity codes and MET intensities," *Medicine and Science in Sports and Exercise*, vol. 32, no. 9, pp. S498-S504, 2000.
 19. Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ, et al. Compendium of physical activities: an update of activity codes and MET intensities. *Medicine and science in sports and exercise*. 2000 Sep;32(9 Suppl):S498-504. PubMed PMID: 10993420.
 20. Azadbakht L, Mirmiran P, Azizi F. Dietary diversity score is favorably associated with the metabolic syndrome in Tehranian adults. *Int J Obes (Lond)*. 2005 Nov;29(11):1361-7. PubMed PMID: 16116493.
 21. Vadiveloo M, Dixon LB, Mijanovich T, Elbel B, Parekh N. Dietary variety is inversely associated with body adiposity among US adults using a novel food diversity index. *J Nutr*. 2015 Mar;145(3):555-63. PubMed PMID: 25733472.
 22. Salehi-Abargouei A, Akbari F, Bellissimo N, Azadbakht L. Dietary diversity score and obesity: a systematic review and meta-analysis of observational studies. *Eur J Clin Nutr*. 2015 Jul 29. PubMed PMID: 26220567.
 23. Vakili M, Abedi P, Sharifi M, Hosseini M. Dietary diversity and its related factors among adolescents: a survey in Ahvaz-Iran. *Glob J Health Sci*. 2013 Mar;5(2):181-6. PubMed PMID: 23445707.
 24. Esmailzadeh A, Azadbakht L. Dietary energy density and the metabolic syndrome among Iranian women. *Eur J Clin Nutr*. 2011 May;65(5):598-605. PubMed PMID: 21224871.
 25. Patterson E, Warnberg J, Poortvliet E, Kearney JM, Sjostrom M. Dietary energy density as a marker of dietary quality in Swedish children and adolescents: the European Youth Heart Study. *Eur J Clin Nutr*. 2010 Apr;64(4):356-63. PubMed PMID: 20068584.
 26. Perez-Escamilla R, Obbagy JE, Altman JM, Essery EV, McGrane MM, Wong YP, et al. Dietary energy density and body weight in adults and children: a systematic review. *J Acad Nutr Diet*. 2012 May;112(5):671-84. PubMed PMID: 22480489.
 27. O'Sullivan TA, Bremner AP, Bremer HK, Seares ME, Beilin LJ, Mori TA, et al. Dairy product consumption, dietary nutrient and energy density and associations with obesity in Australian adolescents. *J Hum Nutr Diet*. 2014 Aug 26. PubMed PMID: 25157608.
 28. Kring SI, Heitmann BL. Fiber intake, not dietary energy density, is associated with subsequent change in BMI z-score among sub-groups of children. *Obes Facts*. 2008;1(6):331-8. PubMed PMID: 20054197.
 29. Ruel MT. Operationalizing dietary diversity: a review of measurement issues and research priorities. *J Nutr*. 2003 Nov;133(11 Suppl 2):3911S-26S. PubMed PMID: 14672290.
 30. Ledikwe JH, Blanck HM, Khan LK, Serdula MK, Seymour JD, Tohill BC, et al. Dietary energy density determined by eight calculation methods in a nationally representative United States population. *J Nutr*. 2005 Feb;135(2):273-8. PubMed PMID: 15671225.
 31. Johnson SB, Freund A, Silverstein J, Hansen CA, Malone J. Adherence-health status relationships in childhood diabetes. *Health Psychol*. 1990;9(5):606-31. PubMed PMID: 2226389.
 32. Kontogianni MD, Vidra N, Farmaki AE, Koinaki S, Belogianni K, Sofrona S, et al. Adherence rates to the Mediterranean diet are low in a representative sample of Greek children and adolescents. *J Nutr*. 2008 Oct;138(10):1951-6. PubMed PMID: 18806106.
-
- Correspondence:
Leila Azadbakht, PhD
Department of Nutrition. School of Public Health
Isfahan University of Medical Sciences - Isfahan, Iran
Tel: +98 3137922719
Fax: +98 3136682509
Email: azadbakht@hlth.mui.ac.ir