

EARLY HOSPITAL READMISSION INCREASES SHORT AND LONG – TERM MORTALITY IN PATIENTS WITH INTERSTITIAL LUNG DISEASE.

Diego Castillo^{1,2}, Silvia Barril^{1,3}, Ana Rodrigo – Troyano^{1,4}, Paloma Millan–Billi^{1,5}, Guillermo Suárez – Cuartín^{1,6}, Ana Alonso^{1,2}, Tomás Franquet^{2,7}, Laura López–Vilaró^{2,8}, Iván Castellví^{2,9}, Vicente Plaza^{1,2}, Oriol Sibila^{1,10}.

¹Respiratory Department. Hospital de la Santa Creu i Sant Pau. Barcelona (Spain)

²Sant Pau Biomedical Research Institute (IIB-SantPau). Barcelona (Spain)

³Respiratory Department. Hospital Universitari Arnau de Vilanova. Lleida (Spain)

⁴Respiratory Department. Hospital Universitari Son Espases, Palma, Illes Balears (Spain)

⁵Respiratory Department. Hospital Germans Trias I Pujol. Badalona (Spain)

⁶Respiratory Department. Hospital Universitari Bellvitge. L'Hospitalet de Llobregat (Spain)

⁷Radiology Department. Hospital de la Santa Creu i Sant Pau. Barcelona. (Spain)

⁸Histopathology Department. Hospital de la Santa Creu i Sant Pau. Barcelona. (Spain)

⁹Rheumatology Department. Hospital de la Santa Creu i Sant Pau. Barcelona (Spain)

¹⁰Thorax Institute. Hospital Clínic. Barcelona (Spain)

ABSTRACT.Objective: To investigate the prognostic impact of early readmission (30 days) on hospitalized patients with Interstitial Lung Disease (ILD). **Methods:** Observational study analysing a cohort of patients hospitalized in a respiratory ward at a University Hospital. Demographic, clinical data and survival status were collected from patients' records. Early readmission was defined as hospitalization within 30 days after patient's discharge. The primary outcome was 90-day and 1-year all-cause mortality. **Results:** Between 2013 to 2016, a total of 2.238 patients were admitted to the respiratory ward and 98 (%) had a diagnosis of ILD. Among them, 74 patients were discharged (25% in-hospital mortality). Early readmission was observed in 15 cases (20.2%). Early readmitted patients were more frequently current smokers (20% vs. 2%, p=0.02). After a multivariate analysis, early readmission was found to be independently associated with 90-day and 1 year mortality (Odds Ratio (OR) 17.6, 95% Confidence Interval (CI) 4.5-69.2, p=0.001 and OR 4.5; 95CI 1.3-15.2, p=0.01, respectively). **Conclusion:** In patients with ILD, early readmission after hospitalization increases both short-term and long term mortality. Thus, preventing early readmission after discharge from hospital admission may have an impact in the clinical course of ILD patients. Further studies are required to identify factors contributing to early readmission.

Key words: Interstitial lung disease, Hospitalization, Mortality, Readmission.

Received: 27 September 2020

Accepted after revision: 2 May 2021

Correspondence: Dr. Diego Castillo Villegas.
Respiratory Department. Hospital de la Santa Creu I Sant Pau.
C/ Sant Antoni M. Claret, 167.
08029 Barcelona (Spain)
Tel: +34935565972 e-mail: dcastillo@santpau.cat

INTRODUCTION

Interstitial lung disease (ILD) encompasses a group of rare lung disorders that share common clinical, radiological and pathological findings (1). The course of ILD is variable(2). Several disease progression markers have been identified, such as

age, lung function decline, extension of fibrotic features on lung imaging or underlying histology(3). Despite this, the disease behaviour remains highly unpredictable and more clinical tools to anticipate the evolution are needed.

Hospitalization has also been identified as a mortality risk factor in ILD. *Moua et al.* showed that in patients with fibrosing ILD, acute respiratory worsening leading to hospitalization were associated with significant in-hospital and post-discharge mortality(4). Although there are several factors linked to in-hospital mortality in ILD, such as clinical presentation, frailty, lung capacity or underlying disease, less is known about post-discharge mortality(5). In other chronic respiratory diseases, like Chronic Obstructive Pulmonary Disease (COPD), patients admitted with an exacerbation have an increased risk of future hospital readmissions(6,7), which is also related with a progressive increase in the risk of death(8). Nonetheless, COPD programs are focused on close follow-up after hospitalization(9,10). Hence, early readmission may influence the disease prognosis in ILD patients too, but to our knowledge, there are no previous studies evaluating this issue.

Therefore, the aim of this study was to investigate the impact of early readmission on mortality in a cohort of hospitalized patients with ILD.

METHODS

Study design

Observational study conducted at the Respiratory Department of Hospital de la Santa Creu i Sant Pau, Barcelona. The study was approved by Hospital de la Santa Creu I Sant Pau ethics committee (IIBSP-EPI-2018-108).

The study was designed following STROBE recommendations for observational studies(11). Using electronic on-site records, we included all consecutive patients admitted to the respiratory ward of our institution from January 2013 to December 2016. Study data collection was anonymous. Regardless of the purpose of this study, patients were managed following institutional protocols based on national and international guidelines. Post-discharge one-year follow-up data was obtained either from our institutional clinical records or from a database of the health-care system in our region.

Primary outcome was 90-day and 1-year all-cause mortality after the index hospitalization. Secondary outcomes were: readmission, length of stay, reason for hospitalization and main diagnosis.

Study population

Of all patients hospitalized during this period in the Respiratory Department ward, we identified those with a diagnosis of ILD stated in their discharge letter in accordance with current *International Classification of Diseases, 10th Revision (ICD-10)* (12). ILD classification was defined by current international and national guidelines based on clinical, radiologic and histologic findings(13,14). Those not fulfilling current guidelines were not included. Patients that died during their first admission were excluded from analysis.

Variables

We collected the following data from patients' records:

Hospitalization: Demographic and clinical data, including comorbid conditions, treatments, history of smoking, laboratory tests and radiology results were assessed at the time of hospital admission.

Early readmission (30 days) after discharge.

ILD diagnosis was specifically reviewed in each case to avoid any potential bias. We collected data regarding diagnosis, including lung function tests, high resolution CT scans, laboratory tests and histology. We ensured that diagnosis was based on current guidelines.

Statistical analyses

Univariate statistics were used to test the association of demographic and clinical characteristics with early hospital readmission. Categorical variables are presented by frequencies and percentages; statistical differences were analysed using a χ^2 test or a Fisher exact test when required. Continuous variables are presented as mean and SD or median and interquartile range (IQR) when data was not distributed normally. Statistical differences among continuous variables were analysed using a Student t test or the corresponding non-parametrical test when it was required. We performed logistic

regression analyses using clinical outcomes evaluated as dependent variables and early hospital readmission as an independent variable. We defined statistical significance as a 2-tailed $p < 0.05$. All analyses were performed with the SPSS 19.0 software program (SPSS Inc, Chicago, IL, USA).

RESULTS

A total of 2,238 patients were admitted to the respiratory ward. Of them, 98 (4,4%) were hospitalized with a diagnosis of ILD. Finally, 74 patients were discharged (25% inpatient mortality). These patients were included in the study (Figure 1).

There were no missing cases during follow-up.

Main demographic and clinical findings are summarized in Table 1. Note that patients were old (median age 74 ± 12), with a slight female predominance (55%) and were not smokers (54%). Chronic Hypersensitivity Pneumonitis and unclassifiable ILD were the most frequent diagnoses. The majority of patients had a high-resolution computed tomography (HRCT) inconsistent with usual interstitial pneumonia (UIP) patterns. Patients presented severe impairment in lung function.

Data regarding hospitalization is shown in Table 2. Mean length of admission was 8 ± 6.9 days, in the majority of cases due to a chest infection (63.5%). Acute respiratory failure was a common

finding among these patients (92%) but only a minority required intubation and mechanical ventilation (4.1%) or high flow oxygen (4.1%). Bronchoscopy was performed in 20% of the cases.

A significant proportion of patients died during the 12 month follow-up period (28, 37.8%). Early readmission occurred in 15 cases (20.2%). No differences in clinical, laboratory and microbiological data were found among patients with or without early hospital readmission, except a higher rate of current smokers in the first group (20% vs. 2%, $p=0.02$) (table 3). Those patients with early readmission were also older (79 ± 6 vs 73 ± 13 years old) with a trend towards significance ($p 0.07$). Besides, the majority presented evidence of acute or sub-acute progression during the first admission (8/15). The causes of readmission were worsening of symptoms due to an acute or sub-acute progression of the disease (8/15), following by hospital acquired pneumonia (2/15), heart failure (2/15) and other extrapulmonary complications (3/15).

Regarding the main objective of this study, early hospital readmission (30 days) was associated with an increased risk of mortality at 90 days, 6 months and 1 year (Table 4). Early readmission was identified as an independent risk factor for 90-day mortality (Odds Ratio (OR) 17.6, 95% Confidence Interval (CI) 4.5-69.2, $p=0.001$), and 1-year mortality (OR 4.5 95CI 1.3-15.2, $p=0.01$).

DISCUSSION

Early hospital readmission (30th days) has been linked to an increased risk of mortality in several respiratory diseases. To our knowledge, this is the first study addressing the prognostic impact of early readmission in ILD patients. Our data shows that early readmission after an index hospitalization is independently associated with short- and long-term mortality in these patients, thus highlighting the importance of post-discharge care and the need for specific clinical care pathways in ILD.

Reducing early readmission of chronic diseases is an objective of many healthcare systems in developed countries, as these events have a significant impact on the course of the disease but many can be prevented(15). In COPD, it is well known that early readmission increases morbimortality(6,16). However, one in five patients with COPD that presented




Figure 1. Study flow chart.

Table 1. Demographic and clinical data summary.

Characteristic	N = 74
Age at admission, mean (range)	74 ± 12 (33 - 90)
Sex, M/F (%/%)	33/41 (45/55)
Smoker status, No. (%)	
Current smoker	4 (5.4)
Ex - smoker	30 (40.5)
Never smoker	40 (54.1)
ILD-diagnoses, No. (%)	
Chronic HP	16 (21.6)
Unclassifiable	13 (17.5)
IPF	9 (12.1)
CTD - ILD	9 (12.1)
Drug / environmental	9 (12.1)
OP	8 (10.8)
Other ILD	10 (13.5)
HRCT Patterns, No. (%)	
UIP Pattern	10 (13.6)
Possible UIP Pattern	16 (21.6)
Inconsistent with UIP Pattern	48 (64.8)
PFT nearest to hospitalization	
FVC, % pred, mean ± SD	66.2 ± 21 (31 - 135) (n = 67)
FEV ₁ , % pred, mean ± SD	67.6 ± 21 (24 - 116) (n = 67)
TLC, % pred, mean ± SD	75.7 ± 21 (48 - 150) (n = 48)
D _{LCO} , % pred, mean ± SD	44.3 ± 19 (15 - 103) (n = 52)
Treatment, No. (%)	
None	47 (63.5)
Immunosuppression	11 (14.8)
Corticoid	16 (29.7)
Oxygen Therapy, Yes/No (%/%)	27/47 (36.5/63.5)

Data are presented as *n* (%) unless otherwise indicated.

HP: Hypersensitivity pneumonitis; IPF: idiopathic pulmonary fibrosis; CTD-ILD: connective tissue disease-interstitial lung disease; OP: organizing pneumonia; HRCT: High-resolution computed tomography; UIP: usual interstitial pneumonia; PFT: pulmonary functional test; % pred: percent predicted.

with an acute exacerbation will require rehospitalization within 30 days of discharge after an admission(17). It is estimated that around 10% to 55% of these episodes may be preventable as several factors have been identified as contributors of early readmission(17,18). To tackle the problem, many countries have implemented programs to reduce COPD early readmissions(19). Our study shows that in ILD, as in COPD, early readmission increases mortality. Thus, preventing early readmissions of these patients could have an impact on the course of the disease.

In our study, 24 patients died during the first hospitalization (25%). The course of ILD is variable,

but in fibrotic ILD-- especially Idiopathic Pulmonary Fibrosis (IPF)--there is a progressive loss of lung function disrupted by episodes of acute exacerbation (AE)(20). These events usually require hospitalization and have a poor prognosis(21). Patients with IPF and other ILD also require hospitalization related to other events (such as infection, cardiovascular or thromboembolic disease)(4). However, in any case, hospitalization has a significant impact on the natural history of the disease(22). *Durheim et al* reviewed the characteristics of 6.665 hospitalized patients with IPF identified from a representative US database(23). Overall, in-patient mortality

Table 2. Hospitalization Characteristics

Characteristic	N = 74
Distribution of admission diagnoses, No. (%)	
Infection	47 (63.5)
Acute exacerbation	13 (17.6)
Sub-acute progression	7 (9.5)
Heart failure	6 (8.1)
Others	1 (1.4)
Mean \pm SD duration of hospitalization (range)	8 \pm 6.9 (2 - 42)
Acute respiratory failure, No. (%)	68 (92)
Need for ICU admission, No. (%)	8 (10.8)
Intubation and mechanical ventilation, No. (%)	3 (4.1)
Need for nasal high flow oxygen, No. (%)	3 (4.1)
Bronchoscopy performed, No. (%)	15 (20.3)
High – dose IV steroids, No. (%)	3 (4.1)
Died within the first year after hospitalization, No. (%)	28 (37.8)

Data are presented as *n* (%) unless otherwise indicated.
ICU: Intensive care unit. IV: intravenous.

was 14.3%, and a total of 684 patients (10.3%) died during the first hospitalization. This percentage is close to a study conducted in Spain in 2014-2015 (14.94%)(24). In an acute setting, these numbers are exacerbated: *Moua et al.* analysed all consecutive patients with fibrotic ILD, hospitalized due to acute respiratory worsening in the institution from 2000 to 2014(4). Interestingly, in-hospital mortality during their first hospitalization accounts for 47% of IPF patients compared with 33% of patients without IPF. In-patient mortality in our study shifts between these studies because the nature of the hospitalization in our cohort includes both patients with AE and other causes of acute worsening, mainly chest infection.

Post discharge mortality, in our whole cohort, was 37.8 % at 12 months. There are few studies that have investigated the factors influencing post discharge mortality in ILD. In the study of *Moua T et al.* it was intriguing that post discharge survival was not related to the cause of hospitalization (AE versus no AE) in either IPF or non-IPF patients(4). More recently, *Yamazaki et al.* showed in a cohort of patients with chronic Idiopathic Interstitial Pneumonia that repeated hospitalization was associated with poor survival (25). But, interestingly, although survival rate after the first hospitalization was significantly better than those with multiple hospitalizations, there were no differences between the second

hospitalization and each subsequent. In our cohort, smoking was the only factor identified as a predictor of readmission. This is in accordance with data from COPD, which shows a smoke-free policy reduces rates of early readmission(26). Although it was not statistically significant, age seems to also have an impact on readmission. No other factors related to the lung condition were identified.

This study has limitations that need to be taken into account when interpreting the results. We recognize the possible bias due to its single-centre design and the retrospective nature itself. Conversely, the strength of our project is that the study analyses, for the first time, the risk factors for mortality associated with early readmission in ILD patients.

In conclusion, our study shows that in patients with ILD, early readmission (30 days) after a hospitalization increases mortality in the short (90 days) and long term (12 months). We found smoking as a possible contributor to early readmission but further research is needed to identify other risk factors. Implementing adequate post discharge programs in ILD to prevent early readmission may have an impact in the course of the disease.

Acknowledgement: Thanks to Mrs. Emily Broe for her assistance with the English edition.

Table 3. Predictors of readmission.

Variable	30 th day Readmission N = 15	No 30 th day Readmission N = 59	P Value
Age at admission, mean \pm SD (range)	79 \pm 6 (67 – 87)	73 \pm 13 (33 – 90)	0.07
Sex M, No. (%)	9/15 (60)	24/59 (41)	0.25
Smoker Status (current) No. (%)	3/15 (20)	1/59 (2)	0.026
HRCT (UIP) No. (%)	7/15 (47)	19/59 (32)	0.37
Baseline PFR (% pred)			
FVC < 50%	2/15 (15)	16/54 (30)	0.5
D _{LCO} < 35%	3/9 (33)	20/43 (46)	0.7
Baseline treatment (immunosuppression) No. (%)	3/15 (20)	8/59 (14)	0.68
Baseline oxygen therapy, No. (%)	7/15 (47)	20/59 (34)	0.38
Admission type (infection), No. (%)	9/15 (60)	38/59 (64)	0.77
Duration of hospitalization (days), mean \pm SD (range)	8 \pm 3 (3 – 15)	8 \pm 7 (2 – 42)	0.9
Acute respiratory failure, No. (%)	15/15 (100)	53/59 (90)	0.33
Need for mechanical ventilation, No. (%)	0/15 (0)	3/59 (5)	0.99
High – dose steroids	0/15 (0)	3/59	0.99

Data are presented as *n* (%) unless otherwise indicated.

HRCT: High-resolution computed tomography; UIP: usual interstitial pneumonia, PFT: pulmonary functional tests; % pred: percent predicted;

Contributions: DC, SB, ART and OS were responsible for study design, analysis of data and manuscript preparation. SB and PM did the literature search and data collection. All authors reviewed the final version of the manuscript. All authors contributed to and approved the final version of the manuscript. DC confirms that he had full access to all the data and had final responsibility for the decision to submit for publication.

Declaration of interest: DC reports personal fees and non-financial support from Roche, personal fees and non-financial support from Boehringer-Ingelheim, grants from Fujirebio, outside the submitted work. In the last three years, VP has received honoraria for speaking at sponsored meetings from Astrazeneca, Boehringer-Ingelheim, MSD and Chiesi, received help assistance to meeting travel from Astrazeneca, Chiesi and Novartis, act as a consultant

Table 4. Multivariate analysis of early hospital readmission (30 th day) as mortality risk factor.

Variable	Multivariable (OR, CI)	P Value
90 - days mortality	17.6 (4.5 – 69.2)	< 0.001
6 - months mortality	7.0 (2.0 – 24.3)	0.002
1 - year mortality	4.5 (1.3 – 15.2)	0.001

for ALK, Astrazeneca, Boehringer, MSD, MundiPharma and Sanofi and received funding/grant support for research projects from a variety of Government agencies and not-for-profit foundations, as well as AstraZeneca, Chiesi and Menarini. Other authors have no conflicts of interest to report.

REFERENCES

- Bradley B, Branley HM, Egan JJ, et al. Interstitial lung disease guideline: the British Thoracic Society in collaboration with the Thoracic Society of Australia and New Zealand and the Irish Thoracic Society. *Thorax*. 2008;63 Suppl 5:v1-58.
- Kolb M, Vasakova M. The natural history of progressive fibrosing interstitial lung diseases. *Respiratory research*. 2019;20(1):57.
- Adegunsoye A, Oldham JM, Chung JH, et al. Phenotypic Clusters Predict Outcomes in a Longitudinal Interstitial Lung Disease Cohort. *Chest*. 2017.
- Moua T, Westerly BD, Dulohery MM, Daniels CE, Ryu JH, Lim KG. Patients With Fibrotic Interstitial Lung Disease Hospitalized for Acute Respiratory Worsening: A Large Cohort Analysis. *Chest*. 2016;149(5):1205-1214.
- Brown AW, Fischer CP, Shlobin OA, et al. Outcomes after hospitalization in idiopathic pulmonary fibrosis: a cohort study. *Chest*. 2015;147(1):173-179.
- Mullerova H, Maselli DJ, Locantore N, et al. Hospitalized exacerbations of COPD: risk factors and outcomes in the ECLIPSE cohort. *Chest*. 2015;147(4):999-1007.
- Roberts MH, Clerisme-Beaty E, Kozma CM, Paris A, Slaton T, Mapel DW. A retrospective analysis to identify predictors of COPD-related rehospitalization. *BMC pulmonary medicine*. 2016;16(1):68.
- Guerrero M, Crisafulli E, Liapikou A, et al. Readmission for Acute Exacerbation within 30 Days of Discharge Is Associated with a Subsequent Progressive Increase in Mortality Risk in COPD Patients: A Long-Term Observational Study. *PloS one*. 2016;11(3):e0150737.
- Shah T, Press VG, Huisingh-Scheetz M, White SR. COPD Readmissions: Addressing COPD in the Era of Value-based Health Care. *Chest*. 2016;150(4):916-926.
- Marcos PJ, Represas Represas C, Ramos C, et al. Impact of a Home Telehealth Program After a Hospitalized COPD Exacerbation: A Propensity Score Analysis. *Archivos de bronconeumologia*. 2020.
- von Elm E, Altman DG, Egger M, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ*. 2007;335(7624):806-808.
- International Statistical Classification of Diseases and Related Health Problems 10th Revision. <https://icd.who.int/browse10/2019/en#/J80-J84>.
- Xaubet A. Diagnóstico y tratamiento de las enfermedades pulmonares intersticiales difusas *Archivos de bronconeumologia*. 2003;39:580-600.
- Travis WD, Costabel U, Hansell DM, et al. An official American Thoracic Society/European Respiratory Society statement: Update of the international multidisciplinary classification of the idiopathic interstitial pneumonias. *American journal of respiratory and critical care medicine*. 2013;188(6):733-748.
- Quality AfHRA. Guide to Prevention Quality Indicators: Hospital Admission for Ambulatory Care Sensitive Conditions. . <https://www.ahrq.gov/downloads/pub/ahrqqi/pqguide.pdf>.
- Soler-Cataluna JJ, Martinez-Garcia MA, Roman Sanchez P, Salcedo E, Navarro M, Ochando R. Severe acute exacerbations and mortality in patients with chronic obstructive pulmonary disease. *Thorax*. 2005;60(11):925-931.
- Shah T, Churpek MM, Coca Perrillon M, Konetzka RT. Understanding why patients with COPD get readmitted: a large national study to delineate the Medicare population for the readmissions penalty expansion. *Chest*. 2015;147(5):1219-1226.
- Auerbach AD, Kripalani S, Vasilevskis EE, et al. Preventability and Causes of Readmissions in a National Cohort of General Medicine Patients. *JAMA Intern Med*. 2016;176(4):484-493.
- Kristensen SR, Bech M, Quentin W. A roadmap for comparing readmission policies with application to Denmark, England, Germany and the United States. *Health Policy*. 2015;119(3):264-273.
- Collard HR, Ryerson CJ, Corte TJ, et al. Acute Exacerbation of Idiopathic Pulmonary Fibrosis. An International Working Group Report. *American journal of respiratory and critical care medicine*. 2016;194(3):265-275.
- Sakamoto S, Shimizu H, Isshiki T, Kurosaki A, Homma S. Pharmacological treatment of acute exacerbation of idiopathic pulmonary fibrosis: a retrospective study of 88 patients. *Sarcoidosis Vasc Diffuse Lung Dis*. 2019;36(3):176-184.
- Spagnolo P, Wuyts W. Acute exacerbations of interstitial lung disease: lessons from idiopathic pulmonary fibrosis. *Curr Opin Pulm Med*. 2017;23(5):411-417.
- Durheim MT, Judy J, Bender S, et al. In-Hospital Mortality in Patients with Idiopathic Pulmonary Fibrosis: A US Cohort Study. *Lung*. 2019;197(6):699-707.

-
24. Pedraza-Serrano F, Jimenez-Garcia R, Lopez-de-Andres A, et al. Characteristics and outcomes of patients hospitalized with interstitial lung diseases in Spain, 2014 to 2015. *Medicine (Baltimore)*. 2019;98(21):e15779.
 25. Yamazaki R, Nishiyama O, Saeki S, Sano H, Iwanaga T, Tohda Y. Characteristics of patients with chronic idiopathic interstitial pneumonia undergoing repeated respiratory-related hospitalizations: A retrospective cohort study. *PloS one*. 2020;15(4):e0232212.
 26. Stallings-Smith SH, H.Y.; Peterson, B.N.; Apatu, E.J.I.; Spaulding, A.C. Smoke-Free Policies and 30-Day Readmission Rates for Chronic Obstructive Pulmonary Disease. *Am J Prev Med*. 2019;57(5).